

Simon, E. 1901. Descriptions d'arachnides nouveaux de la famille des Attidae (suite). Ann. Soc. ent. Belg. 45: 151-152.

Natta chionogaster (Simon, 1901)

C. chionogaster sp. nov. — ♂. Long. 3,5-4 mill. — Cephalothorax niger postice rufesceni-tinctus, supra squamulis fuscis postice, versus marginem, sensim albidioribus vestitus, utrinque, in declivitate fere glaber sed linea marginali nivea cinctus. Pili oculorum breves, rubri. Pili clypei densi niveti. Abdomen oblongum, supra nigrum, parce tenuiter et longe fulvo-pilosum et squamulis grossis subrotundis obscure cinereo-violaceis vestitum, subtus fulvum albo-argenteo-squamulatum. Chelæ longæ. parallelæ, nigritidæ, minute et parce rugosæ. Partes oris nigræ. Sternum nigrum vel fuscum, albo-squamulatum. Pedes 4ⁱ paris robusti, fusco-castanei vel nigricantes, patella paulo dilutiore, metatarso tarsoque fulvis, tibia longe ovata et supra et subtus longe et crebre sed inordinate nigro-crinita. Reliqui pedes fulvi, patellis tibiis metatarsisque posticis utrinque fusco-lineatis. Pedes-maxillares fusti apice fulvi, albo-squamulati, tibia patella breviore, extus apophysi longa, gracili, antice directa et apice oblique truncata armata, tarso angusto, longissimo et tereti, bulbo rufulo, ad basin ovato, ad apicem longissime producio et acuto.

♀. Long. 4 mill. — A mari distert abdome supra obscure cinereo violaceo-squamulato, maculis sinuosis biseriatis aureo vel rubro squamulatis maculaque apicali parva alba notato, subtus albo-argenteo-squamulato, chelis fulvo rufulis levibus et nitidis, pedibus anticis fuscis, annulis latis albo-squamulatis ornatis, tibiis haud sinuatis, pedibus-maxillaribus luteis ad basin infuscatis, tarso longissimo acuminato. Area genitalis antice depressa, postice plagula semicirculari fulvo-nitida, stria tenue secta, notata. — Africa austr. : Prom. Bonac Spei, Kimberley, Transvaal! — Madagascar : Diego-Suarez (C. Alluaud).



L. *C. chionogaster* E. Sim. P. M. ♂.
[Simon, E. 1901. Histoire naturelle des araignées. Paris. 2: 541].