



Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by Far East Branch
of the Russian Entomological Society
and Laboratory of Entomology,
Institute of Biology and Soil Science,
Vladivostok

Number 247: 1-7

ISSN 1026-051X

May 2012

NEW SPECIES OF THE GENUS *ARGYRA* MACQUART, 1834 (DIPTERA: DOLICHOPODIDAE) FROM THE RUSSIAN FAR EAST AND JAPAN

O. P. Negrobov¹⁾, M. Satô²⁾ and O.V. Selivanova¹⁾

1) Voronezh State University, Voronezh, 394006, Russia. E-mail: negrobov@list.ru

2) Rishiri Town Museum, Senhoshi, Rishiri Island, Hokkaido 097-0311, Japan.
E-mail: dolicho@d1.dion.ne.jp

Two species of the genus *Argyra* Macquart, 1834 are described from the Russian Far East and Japan. *A. igori* Negrobov, Satô et Selivanova **sp. n.** is closely related to *A. spoliata* Kowarz, 1879, and *A. zlobini* Negrobov, Satô et Selivanova **sp. n.** is similar with *A. atriceps* Loew, 1857.

Key words: Dolichopodidae, *Argyra*, new species, Russia, Japan.

О. П. Негрбов, М. Сато, О. В. Селиванова. Новые виды рода *Argyra* Macquart, 1834 (Diptera: Dolichopodidae) с Дальнего Востока России и из Японии // Дальневосточный энтомолог. 2012. N 247. С. 1-7.

С Дальнего Востока России и из Японии описаны два новых вида рода *Argyra* Macquart, 1834. *A. igori* Negrobov, Satô, et Selivanova **sp. n.** близок к *A. spoliata* Kowarz, 1879, а *A. zlobini* Negrobov, Satô et Selivanova **sp. n.** – к *A. atriceps* Loew, 1857.

1) Воронежский государственный университет, Воронеж, 394006, Россия.

2) Музей г. Рисири, Сенхоши, Хоккайдо, 097-0311, Япония.

INTRODUCTION

Most Palaearctic species of the genus *Argyra* are known from Europe (Negrobov, 1991). The following species of the genus *Argyra* were described from Asian

SHORT COMMUNICATION

P. G. Nemkov. DIGGER WASPS OF THE GENUS *STIZOIDES* GUÉRIN-MÉNEVILLE (HYMENOPTERA, CRABRONIDAE, BEMBICINAE) OF THE FAUNA OF RUSSIA AND NEIGHBORING COUNTRIES. – Far Eastern Entomologist. 2012. N 247: 8-13.

Summary. Five *Stizoides* species of the fauna of Russia and neighboring countries are reviewed. A key to the species is provided.

Key words: Digger wasps, Hymenoptera, Crabronidae, Bembicinae, *Stizoides*, Russia.

П. Г. Немков. Роющие осы рода *Stizoides* Guérin-Méneville (Hymenoptera, Crabronidae, Bembicinae) фауны России и сопредельных стран // Дальневосточный энтомолог. 2012. N 247. С. 8-13.

Резюме. Дается обзор 5 видов рода *Stizoides* фауны России и сопредельных стран. Приводится оригинальная определительная таблица видов.

INTRODUCTION

Stizoides Guérin-Méneville, 1844 is a not numerous digger wasp genus included 30 species which inhabit all continents except Oriental, Australian, and Neotropical regions (Pulawski, 2012). The species of *Stizoides* are cleptoparasites in the nests of Orthoptera-hunting digger wasps of the genera *Sphex* Linnaeus, 1758, *Palmodes* Kohl, 1890, *Prionyx* Vander Linden, 1827 (Sphecidae), and *Stizus* Latreille, 1802 (Crabronidae) (Bohart & Menke, 1976; Ohl, 1999).

This genus was not especially studied in Russia, but there are species keys of European part of USSR (Pulawski, 1978), Kazakhstan and Middle Asia (Kazenas, 1978), and world-wide fauna (Ohl, 1999). The material used in this study is from the collections of Zoological Institute of Russian Academy of Sciences (St-Petersburg) and Institute of Biology and Soil Science of Russian Academy of Sciences (Vladivostok).

Genus *Stizoides* Guérin-Méneville, 1844

Stizoides Guérin-Méneville, 1844: 438 (as subgenus of *Stizus*). Type species: *Larra fasciata* Fabricius, 1798, junior secondary homonym of *Stizus fasciatus* (Fabricius, 1798) (= *Sphex assimilis* Fabricius, 1787), designated by J. Parker, 1929: 10.

Five species known in Russia and neighboring countries are included in the key.

Key to the species

1. Forewing base hyaline. Sternite II with macropunctures sparser medially than laterally, basomedially with densely pubescent micropunctate patch. 10.0–23.0 mm *S. assimilis* (Fabricius)
- Forewing base infumate. Sternite II with regular macropunctures, without pubescent patch 2
2. Antennal base and legs black 3
- Antennal base and legs for the most part reddish-yellow 4

3. Forewing uniformly infumate. 12.0–22.0 mm *S. melanopterus* (Dahlbom)
 – Forewing apex with broad hyaline area. 11.5–21.0 mm *S. tridentatus* (Fabricius)
 4. All tergites yellow or with broad yellow band. 12.5–16.0 mm
 *S. cyanopterus* (Gussakovskij)
 – Only tergites I-III yellow, remaining tergites black. 8.5–19.5 mm
 *S. crassicornis* (Fabricius)

List of the species

Stizoides assimilis (Fabricius, 1787)

Sphex assimilis Fabricius, 1787: 276, sex not indicated (lectotype [designated by van der Vecht, 1961: 54] – ♂, India, Tamil Nadu, Tranquebar, [Copenhagen]). – *Stizus egregius* Gussakovskij, 1928: 11, ♀ (syntypes – ♀ ♀, Turkmenistan, Yolatan station near Mary, [depository unknown]), synonymized with *Stizoides assimilis* by Ohl, 1999: 81; Myartseva, 1972: 89, 1976: 74; Kazenas, 1978: 83; Nazarova, 1998: 41; Ohl, 1999: 81; Kazenas, 2001: 49, 232.

MATERIAL. Tajikistan: 1 ♀, 1 ♂, Aivadz, outfall of Kafirnigan River, 30.VII 1934 (Gussakovskij).

DISTRIBUTION. Israel, Palestine, Saudi Arabia, Yemen, UAE, Kyrgyzstan, Uzbekistan, Turkmenistan, Tajikistan, Iran, Pakistan, India, Nepal, Morocco, Algeria, Egypt, Mauritania, Senegal, Mali, Sudan.

Stizoides crassicornis (Fabricius, 1787)

Tiphia crassicornis Fabricius, 1787: 278, sex not indicated (lectotype [designated by van der Vecht, 1961: 54] – ♀, Tunisia, Porto Farina, [Copenhagen]). – *Stizus fulvipes* Eversmann, 1846: 440, ♀, ♂ (lectotype [designated by Ohl, 1999: 112] – ♂, Russia, lower Volga area, [Kraków]), synonymized with *Stizus crassicornis* by Handlirsch, 1892: 104; Eversmann, 1849: 392; Radoszkowski, 1877: 38, 1892: 578. – *Pompilus alienus* Fischer de Waldheim, 1843: 3, sex not indicated (lectotype [designated by Harttig, 1932: 146] – sex not indicated, southern Russia, no specific locality, [Dresden, destroyed in World War II]), synonymized with *Stizus crassicornis* by Harttig, 1932: 146. – *Stizus crassicornis*: Becker, 1880: 153; F. Morawitz, 1891: 225. – *Stizoides crassicornis*: Kazenas, 1972: 147; Myartseva, 1972: 89; Myartseva, 1976: 74; Kazenas, 1978: 83; Pulawski, 1978: 206; Ohl, 1999: 112; Kazenas, 2001: 49, 232, 2002: 128, 2004: 111; Shoreenko, 2005: 168.

MATERIAL. Turkmenistan: 1 ♀, 4 ♂, Imam-Baba, 1932 (without other data) (Shchestakov).

DISTRIBUTION. France, Spain, Italy, Greece, Cyprus, Ukraine (Crimea), Turkey, Israel, Palestine, Russia (Saratov, Volgograd, and Astrakhan regions), Azerbaijan, Kazakhstan, Kyrgyzstan, Uzbekistan, Turkmenistan, Tajikistan, Iran, Algeria, Tunisia, Egypt.

Stizoides cyanopterus (Gussakovskij, 1928)

Stizus cyanopterus Gussakovskij, 1928: 10, ♀ (holotype – ♀, Uzbekistan, Khiva, [depository unknown]). – *Stizoides cyanopterus*: Myartseva, 1972: 89; Myartseva, 1976: 74; Kazenas, 1978: 83; Ohl, 1999: 115; Kazenas, 2001: 49, 232.

MATERIAL. Turkmenistan: 1 ♂, Akhcha-Kuima, 6.VI 1988 (Lelej).

DISTRIBUTION. Georgia, Uzbekistan, Turkmenistan, Iran.

***Stizoides melanopterus* (Dahlbom, 1845)**

Stizus melanopterus Dahlbom, 1845: 478, ♂, actually ♀ (holotype – ♀, Greece, Rhodes Island, [Lund]). – *Stizus concolor* Eversmann, 1849: 392, ♀ (holotype [or syntypes] – ♀, Kazakhstan, no specific locality, [depository unknown]), synonymized with *Stizoides melanopterus* by Handlirsch, 1892: 102; Radoszkowski, 1877: 39; Handlirsch, 1892: 103. – *Stizoides melanopterus*: Gussakovskij, 1935: 440; Kazenas, 1978: 83; Pulawski, 1978: 206; Nazarova, 1998: 41; Ohl, 1999: 127; Kazenas, 2001: 49, 2002: 128, 2004: 111; Shorenko, 2005: 168; Kazenas, 2008: 105.

MATERIAL. Turkmenistan: 1 ♀, Kerki, 15.VI 1933 (Ushchinskii).

DISTRIBUTION. Croatia, Greece, Ukraine (Crimea), Turkey, Lebanon, Israel, Palestine, Russia (Volgograd region), Armenia, Kazakhstan, Kyrgyzstan, Uzbekistan, Turkmenistan, Tajikistan, Iraq.

***Stizoides tridentatus* (Fabricius, 1775)**

Crabro tridentatus Fabricius, 1775: 373, sex not indicated (holotype – ♀, southern Europe, no specific locality, [Copenhagen]). – *Larra bifasciata* Fabricius, 1798: 252, sex not indicated (lectotype [designated by van der Vecht, 1961: 55] – ♂, Morocco, Tanger, [Copenhagen]), synonymized with *Larra tridentata* by F. Smith, 1856: 351. – *Stizus bifasciatus*: Eversmann, 1849: 392; Becker, 1880: 153; Morawitz, 1891: 225. – *Stizus unifasciatus* Radoszkowski, 1877: 39, ♀, ♂ (lectotype [designated by Ohl, 1999: 141] – ♀, Uzbekistan, Samarkand, [Moscow]), synonymized with *Stizus tridentatus* by Handlirsch, 1892: 98; Radoszkowski, 1893: 63; Gussakovskij, 1935: 440; Islamov, 1970: 63; Myartseva, 1972: 89, 1976: 74. – *Stizus tridentatus*: Kohl & Handlirsch, 1889: 281; F. Morawitz, 1893: 421; Myartseva, 1963: 60. – *Stizoides tridentatus*: Myartseva, 1965: 90; Tsuneki, 1971: 207; Kazenas, 1972: 147, 1978: 83; Pulawski, 1978: 206; Islamov, 1986: 523; Shkuratov, 1998: 97; Ohl, 1999: 136; Kazenas, 2001: 49, 232, 2002: 129; Shkuratov, 2002: 140; Kazenas, 2004: 111; Nazarova, 2004: 107; Shorenko, 2005: 167; Kazenas, 2007: 92.

MATERIAL. Azerbaijan: 1 ♀, Altan, 30.VI 1927 (Zapolskii).

DISTRIBUTION. France, Portugal, Spain, Switzerland, Italy (including Sicilia and Sardinia), Croatia, Serbia, Macedonia, Albania, Greece (including Crete), Hungary, Romania, Bulgaria, Cyprus, Ukraine, Turkey, Israel, Russia (Saratov, Volgograd, Rostov, Astrakhan, and Orenburg regions), Azerbaijan, Kazakhstan, Kyrgyzstan, Uzbekistan, Turkmenistan, Tajikistan, Iran, Mongolia, Morocco, Algeria, Tunisia, Libya, Egypt.

ACKNOWLEDGEMENTS

My great thanks are due to S.A. Belokobylskij for kindly loaned specimens as well as A.S. Lelej for valuable comments. The work was supported by grants of the Far Eastern Branch of the Russian Academy of Sciences No 12-III-A-06-074, No 12-I-OBH-02, and No 12-I-II30-03.

REFERENCES

- Becker, A. 1880. Beiträge zu meinem Verzeichnissen der um Sarepta und am Bogdo vorkommenden Pflanzen und Insekten, und Beschreibung einer *Mylabris*-Larve. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 55: 145–156.
- Bohart, R.M. & Menke, A.S. 1976. *Sphecid Wasps of the World. A generic revision*. Berkeley, Los Angeles, London: University of California Press: ix + 695 p.

- Dahlbom, A.G. 1845. *Hymenoptera Europaea praecipue borealia; formis typicis nonnullis Specierum Generumve Exoticorum aut Extraneorum propter nexum systematicus associatis; per Familias, Genera, Species et Varietates disposita atque descripta. Tomus: Sphecx in sensu Linneano. Fasc. 3.* Lund: Officina Lundbergiana: 353–528.
- Eversmann, E. 1846. Hymenopterorum Rossicorum species novæ vel parum cognitæ, descriptæ et ex parte depictæ. Decas prima. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 19: 436–443, pl. IV–V.
- Eversmann, E. 1849. Fauna Hymenopterologica Volgo-Uralensis. Fam. III. Sphegidae Latr. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 22: 359–436.
- Fabricius, J.Ch. 1775. *Systema Entomologiae, sistens Insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus.* Flensburgi et Lipsiae: Kortii: 832 p.
- Fabricius, J.Ch. 1787. *Mantissa Insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiispecificis, emendationibus, observationibus, Vol. 1.* Hafniae: C.G. Proft.: xx + 348 p.
- Fabricius, J.Ch. 1798. *Supplementum Entomologiae Systematicae.* Hafniae: C.G. Proft et Storch: 572 p.
- Fischer de Waldheim, G. 1843. Observata quaedam de Hymenopteris Rossicis. *Magasin de Zoologie, d'Anatomie Comparée et de Paléontologie (Deuxième Série)*, 5: 1–4, pl. 122.
- Guérin-Méneville, F.E. 1844. *Iconographie du règne animal de G. Cuvier. Tome III. Insectes.* Paris: Imprimerie de Fain et Thunot: 576 p.
- Gussakovskij, V.V. 1928. Novyie vidy Sphecidae iz Zakaspiya i Khivy. *Izvestiya Kursov Prikladnoy Zoologii i Fitopatologii*, 4: 3–19. (In Russian).
- Gussakovskij, V.V. 1935. K faune os (Hymenoptera, Sphecodea et Vespodea) Tadzhiqistana. *Trudy Tadzhiqskoy Bazy Akademii Nauk SSSR*, 5: 409–467. (In Russian).
- Handlirsch, A. 1892. Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen VI. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I*, 101: 25–205, pl. I–III.
- Hartig, G. 1932. Was ist *Pompilus alienus* Fisch.-Waldh.? (Hym.). *Konowia*, 11: 146.
- Islamov, Sh.D. 1970. K poznaniyu os gornykh chastey Chirchikskogo basseina. *Tashkentskiy Gosudarstvennyi Universitet im. V.I. Lenina. Nauchnyie Trudy*, 378: 61–66.
- Islamov, Sh.D. 1986. Royushchiye osy (Hymenoptera, Sphecidae) gornykh rayonov Uzbekistana. *Entomologicheskoye Obozreniye*, 65: 513–534. (In Russian).
- Kazenas, V.L. 1972. Royushchiye osy (Hymenoptera, Sphecidae) yugo-vostochnogo Kazakhstana. *Trudy Vsesoyuznogo Entomologicheskogo Obshchestva*, 55: 93–186. (In Russian).
- Kazenas, V.L. 1977. Novye vidy royushchikh os roda *Cerceris* Latr. (Hymenoptera, Sphecidae) iz Kazakhstana. *Entomologicheskoye Obozreniye*, 56: 431–434. (In Russian).
- Kazenas, V.L. 1978. *Royushchiye osy Kazakhstana i Sredney Azii (Hymenoptera, Sphecidae). Opredelitel.* Alma Ata: Izdatelstvo Nauka Kazakhskoy SSR: 172 p. (In Russian).
- Kazenas, V.L. 2001. *Fauna i biologiya royushchikh os (Hymenoptera, Sphecidae) Kazakhstana i Sredney Azii.* Almaty: KazgosINTI: 333 p. (In Russian).
- Kazenas, V.L. 2002. Royushchiye osy (Hymenoptera, Sphecidae) Kazakhstana. *Tethys Entomological Research*, 4: 1–174. (In Russian).
- Kazenas, V.L. 2004. Royushchiye osy (Hymenoptera: Apoidea: Sphecidae, Crabronidae) Zapadnogo Tyan-Shanya. *Tethys Entomological Research*, 10: 97–116. (In Russian).
- Kazenas, V.L. 2007. Faunisticheskiy obzor royushchikh os (Hymenoptera, Sphecidae i Crabronidae) Kurgaldzhinskogo zapovednika i privileyushchikh territoriy (Tsentralnyi Kazakhstan). *Selevinia*, 2006: 89–94. (In Russian).

- Kazenas, V.L. 2008. Redkiye vidy royushchikh os (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) Yugo-Vostochnogo Kazakhstana i ikh okhrana. *Tethys Entomological Research*, 16: 97–108. (In Russian).
- Kohl, F.F. & Handlirsch, A. 1889. Transcaspische Hymenopteren. *Verhandlungen der kaiserlich-königlichen Zoologisch-Botanischen Gesellschaft in Wien*, 39: 267–286, pl. VII.
- Morawitz, F. 1891. Ueber Astrachan'sche Fossorien. *Horae Societatis Entomologicae Rossicae*, 25: 175–233.
- Morawitz, F. 1893. Catalog der von D. Glasunov in Turkestan gesammelten Hymenoptera Fossoria. *Horae Societatis Entomologicae Rossicae*, 27: 391–428.
- Myartseva, S.N. 1963. Ekologicheskoye raspredeleniye royushchikh os (Hymenoptera, Sphecidae) v nizoviyakh Murgaba. *Izvestiya Akademii Nauk Turkmenskoy SSR. Seriya Biologicheskikh Nauk*, 4: 56–63. (In Russian).
- Myartseva, S.N. 1965. Royushchiye osy (Hymenoptera, Sphecidae) nizoviy Murgaba. In: Medvedev, G.S., Luppova, A.N. (eds.). *Nasekomye nizoviy Murgaba (vostochnaya Turkmeniya). Fauna, ekologiya, khozaystvennoye znacheniye*. Ashhabad: Turkmenskoye Izdatel'stvo: 147 p.
- Myartseva, S.N. 1972. Fauna sfetsid yuzhnoy Turkmenii. In: Tokgayev, T.B., Myartseva, S.N. (eds.). *Nasekomye yuzhnoy Turkmenii*. Ashhabad: Ylym: 75–100. (In Russian).
- Myartseva, S.N. 1976. Zаметки по биологии maloizuchennykh sfetsid Turkmenii (Hymenoptera, Sphecidae). In: Tashliev, A.O., Myartseva, S.N. (eds.). *Ekologiya i khozaystvennoye znachenye nasekomykh Turkmenii*. Ashhabad: Ylym: 72–82. (In Russian).
- Nazarova, Sh.D. 1998. O royushchikh osakh (Sphecidae) zapovednika "Tigrovaya Balka". *Izvestiya Akademii Nauk Respubliki Tadjikistan. Otdeleniye Biologicheskikh i Meditsinskikh Nauk*, 139: 38–43. (In Russian).
- Nazarova, Sh.D. 2004. O royushchikh osakh fauny Badakhshana. In: Abdusalyamov, I.A. (ed.). *Fauna i ekologiya zhivotnykh Tadjikistana*. Dushanbe: Akademiya Nauk Respubliki Tadjikistan, Institut Zoologii i Parazitologii im. E.N. Pavlovskogo: 103–109.
- Ohl, M. 1999. A revision of *Stizoides* Guérin-Méneville, 1844: taxonomy, phylogenetic relationships, biogeography, and evolution (Hymenoptera: Apoidea: "Sphecidae"). *Mitteilungen aus dem Museum für Naturkunde in Berlin, Zoologische Reihe*, 75: 63–169.
- Parker, J.B. 1929. A generic revision of the fossorial wasps of the tribes Stizini and Bembicini, with notes and descriptions of new species. *Proceedings of the United States National Museum*, 75: 1–203, pl. I–XV.
- Pulawski, W.J. 1978. Nadsem. Sphecoidea. In: Medvedev, G.S. (ed.). *Opredelitel nasekomykh evropeyskoi chasti SSSR. Tom III. Pereponchatokrylye. Pervaya chast*. Leningrad: Nauka: 584 p. (In Russian).
- Pulawski, W.J. 2012. *Catalog of Sphecidae sensu lato*. California Academy of Sciences, Golden Gate Park, San Francisco, California, USA. Available from: http://research.calacademy.org/ent/catalog_sphecidae/ (accessed 15 April 2012).
- Radoszkowski, O.I. 1877. Sphegidae. *Izvestiya Imperatorskogo Obshchestva Lyubiteley Estestvoznaniya, Antropologii i Etnografii pri Imperatorskom Moskovskom Universitete*, 26: 1–87, pl. I–VIII. (Voyage au Turkestan d'A.P. Fedtchenko, fasc. 14, tome 2, part. 5).
- Radoszkowski, O.I. 1892. Essai sur une classification des Sphegides in sensu Linneano d'après la structure des armures copulatrices. *Bulletin de la Société Impériale des Naturalistes de Moscou (Nouvelle Série)*, 5: 571–596, pl. XIX–XXIII.
- Radoszkowski, O.I. 1893. Faune hyménoptérologique transcaspienne (Suite et fin). *Horae Societatis Entomologicae Rossicae*, 27: 38–81.

- Shkuratov, A.V. 1998. Dopolneniya k faune royushchikh os (Hymenoptera, Sphecidae) Rostovskoy oblasti . *Izvestiya Vysshikh Uchebnykh Zavedeniy. Severo-Kavkazskiy Regyon. Estestvennyie Nauki*, 3: 97–99. (In Russian).
- Shkuratov, A.V. 2002. Fauna royushchikh os (Hymenoptera, Sphecidae) stepnogo zapovednika "Rostovskiy" i eyo osobennosti po sravneniyu s faunoy raznotravo-tipchakovykh stepy Rostovskoy oblasti. *Trudy Gosudarstvennogo Zapovednika "Rostovskiy"*, 1: 138–156. (In Russian).
- Shorenko, K.I. 2005. K faune royushchikh os (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) Krymskogo poluostrova. *Kavkazskiy Éntomolgicheskiy Byulleten*, 1: 161–170. (In Russian).
- Smith F. 1856. *Catalogue of hymenopterous insects in the collection of the British Museum. Part IV. Sphegidae, Larridae and Crabronidae*. London: Taylor and Francis. P. 207–497.
- Tsuneki, K. 1971. Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 239. Sphecidae (Hymenoptera). I–II. *Acta Zoologica Academiae Scientiarum Hungaricae*, 17: 139–217.
- van der Vecht, J. 1961. Hymenoptera Sphecoidea Fabriciana. *Zoologische Verhandelingen*, 48: 1–85.

Author's address:

Institute of Biology and Soil Science,
 Far Eastern Branch of
 Russian Academy of Sciences,
 Vladivostok, 690022, Russia
 E-mail: p_nemkov@ibss.dvo.ru