PROCEEDINGS

of the

ENTOMOLOGICAL SOCIETY

of WASHINGTON



DEPARTMENT OF ENTOMOLOGY SMITHSONIAN INSTITUTION WASHINGTON, D.C. 20560

PUBLISHED QUARTERLY

CONTENTS

BURGER, J. F. and F. C. THOMPSON—The Tabanus striatus complex (Diptera:

	Tabanidae): A revision of some Oriental horse fly vectors of surra	339
C	ARROLL, J. F., J. W. KIMBROUGH, and W. H. WHITCOMB—Mycophagy by Aphaenogaster spp. (Hymenoptera: Formicidae)	326
С	LARK, W. E. and D. M. ANDERSON—The genus <i>Chionanthobius</i> Pierce (Coleoptera: Curculionidae): Descriptions of a new species on <i>Forestiera</i> (Oleaceae) and of the larva and pupa of <i>C. schwarzi</i> Pierce	185
F	OOTE, B. A.—Biology and immature stages of <i>Lytogaster excavata</i> , a grazer of bluegreen algae (Diptera: Ephydridae)	304
G	AGNÉ, R. J.—A new species of <i>Endaphis</i> (Diptera: Cecidomyiidae) endoparasitic in aphids in British Columbia	222
K	APLAN, M. and F. C. THOMPSON—New Syrphidae from Israel (Diptera)	198
K	NUTSON, L.—New combinations and synonymies in Palearctic and Nearctic Sciomyzidae (Diptera)	332
K	ONDRATIEFF, B. C., J. W. S. FOSTER, III, and J. R. VOSHELL, JR.—Description of the adult of <i>Ephemerella berneri</i> Allen and Edmunds (Ephemeroptera: Ephemerellidae) with biological notes	300
K	ORMILEV, N. A.—A new micropterous species of <i>Carventus</i> Stal from Chile (Hemiptera: Aradidae)	296
N	AKAHARA, S.—The proper placements of the Nearctic soft scale species assigned to the genus <i>Lecanium</i> Burmeister (Homoptera: Coccidae)	283
R.	ADOVIĆ, I. T.—Anatomy and function of the sting apparatus of stingless bees (Hymenoptera: Apidae: Apinae: Meliponini)	269

(Continued on back cover)

CONTENTS

(Continued from front cover)

RAY, C. H., JR. and M. L. WILLIAMS—Redescription and lectotype designation of the tessellated scale, <i>Eucalymnatus tessellatus</i> (Signoret) (Homoptera: Coccidae)	230
SCARBROUGH, A. G.—Ethology of <i>Eudioctria tibialis</i> Banks (Diptera: Asilidae) in Maryland: Seasonal distribution, abundance, diurnal movements, and behaviors	245
SCARBROUGH, A. G.—Ethology of <i>Eudioctria tibialis</i> Banks (Diptera: Asilidae) in Maryland: Prey, predator behavior, and enemies	258
SHOLES, O. D. V.—Herbivory by species of <i>Trirhabda</i> (Coleoptera: Chrysomelidae) on <i>Solidago altissima</i> (Asteraceae): Variation between years	274
SPANGLER, P. J.—A new water beetle, <i>Troglochares ashmolei</i> , n. gen., n. sp., from Ecuador; the first known eyeless cavernicolous hydrophilid beetle (Coleoptera:	21/
Hydrophilidae)	316
STEYSKAL, G. C.—A bibliography of the Malaise trap	225
TODD, E. L.—The noctuoid moths of the Antilles—Part I (Lepidoptera: Dioptidae)	324
WIRTH, W. W. and N. C. RATANAWORABHAN—The Oriental species of the genus Dibezzia Kieffer (Diptera: Ceratopogonidae)	287
YONKE, T. R.—Descriptions of two new species of Neotropical <i>Leptoglossus</i> Guérin (Hemiptera: Coreidae)	213
NOTES:	
CRABILL, R. E., JR.—Synonymy by way of teratology (Chilopoda: Lithobiomorpha: Lithobiidae)	359
CRABILL, R. E., JR.—On the true identity of <i>Zygethobius pontis</i> Chamberlin (Chilopoda: Lithobiomorpha: Henicopidae)	360
VOEGTLIN, D.—Notes on a European aphid (Homoptera: Aphididae) new to North America	361
BOOK REVIEW:	
KOSZTARAB, M.—Coccoids of the Far East USSR (Homoptera, Coccinea) With a Phylogenetic Analysis of the Coccoid Fauna of the World (Evelyna M. Danzig)	363
SUMMARY REPORTS OF SOCIETY OFFICERS FOR 1980	366
SOCIETY MEETINGS	367

A BIBLIOGRAPHY OF THE MALAISE TRAP

GEORGE C. STEYSKAL

Cooperating Scientist, Systematic Entomology Laboratory, IIBIII, Agric. Res., Sci. and Educ. Admin., USDA, % U.S. National Museum of Natural History, Washington, D.C. 20560.

Abstract.—The flight trap designed by Malaise in 1937 was slow to attract attention but it has now become a major tool in general collecting and surveillance of insects. Its principles of operation are discussed, with mention of a few similar devices. All published notes that have come to the author's attention are listed chronologically.

The Swedish entomologist René Malaise published in 1937 the description of a new kind of trap that had performed well in general collecting for several years. It was many years before other entomologists began to use this kind of trap, but all who did became enthusiastic about its general efficiency in capturing all kinds of flying insects at all times of the day, even night-flying insects that are not attracted to light. Modifications in the design have been made, and the trap has been adapted to the use of carbon dioxide gas as an attractant. It is now considered to be one of the major advances in collecting methods in this century.

The Malaise trap is essentially an interceptive device in which screens of non-opaque fine-meshed fabric netting serve as baffles to impede the flight of insects. The tendency of flying insects to go over or around objects in their path is then used to direct them upward into a closed chamber. The chamber is generally a killing bottle, but may contain a killing and preserving fluid. The original Malaise design and those now in use do not include an attractant, but modifications for use of a lure, notably carbon dioxide gas, have been made.

A few citations to devices that are not strictly Malaise traps have also been included. The window box trap designed by C. F. Hodge and described by Bishopp (1916) operates on much the same principle as the Malaise trap. The traps designed by Burgess (1935) consist of opaque baffles and use an attractant to lure insects into an open-sided 'lean-to.' A bright area near the top of the structure is provided with a glass jar into which the insects are attracted when they attempt to escape upwards from the semi-enclosure.

The interceptive trap described by Leech (1955) is very similar to the Malaise trap but does not include a killing chamber; the operator must collect the intercepted insects by other means. The traps based upon the eeltrap principle, called a Reuse in German, are not included here, although the killing chamber used with the Malaise trap is often constructed on this principle, that of a funnel set in the mouth of a jar. The McPhail trap (Steyskal, 1977) and many emergence traps and cone traps set over a lure also use this principle.

Malaise's design was apparently independent of any previous design. He says, "During my extensive travels I have repeatedly found that insects happened to enter my tent, and that they always accumulated at the ceiling-corners in vain attempts to escape at that place without paying any attention to the open tent-door. On one occasion one of the upper tent-corners happened to have a small hole torn in the fabric, and through this hole all the insects pressed their way and escaped. Later on the idea occurred to me, that, if insects could enter a tent and not find their way out, and always persistently tried to reach the ceiling, a trap, made as invisible as possible and put up at a place where insects are wont to patrol back and forth, might catch them better than any tent and perhaps better than a man with a net, as a trap could catch all the time, by night as well as by day " Thus, the whole idea was original.

BIBLIOGRAPHY

The following bibliography is arranged chronologically by years and alphabetically under each year. Citations not specifically referring to Malaise traps are enclosed in parentheses. No claim is made for completeness. It is hoped that supplementary citations, both by me and by others, will be published.

- (1916 and later editions. Bishopp, F. C. Flytraps and their operation. U.S. Dep. Agric. Farmers' Bull. 734, 14 pp. Describes the Hodge trap, a screen box trap to be placed in a window; operates similarly to the Malaise trap.)
- (1935. Burgess, R. W. The eye gnat in the Coachella Valley, California, U.S. Dep. Agr. Bur. Entomol. Plant Quarantine Bul. E-335. See also 1978, Rogoff.)
- 1937. Malaise, R. A new insect-trap. Entomol. Tidskr. 58: 148-160.
- 1955. Beirne, B. P. Collecting, preparing and preserving insects. Can. Dep. Agric. Sci. Serv. Entomol. Div. 133 pp. Reference, pp. 19–20.
- (——. Leech, H. B. Cheesecloth flight trap for insects. Can. Entomol. 87; 200.)
- 1957. Roos, T. Studies on upstream migration in adult stream-dwelling insects. Rep. Inst. Freshw. Res. Drottningholm 38: 167–193.
- 1958. Oldroyd, H. Collecting, preserving and studying insects. Hutchinson Sci. and Technol., London; Macmillan, New York, 327 pp. Reference, pp. 57–59.
- 1960. Nielsen, E. T. A note on stationary nets. Ecology 41: 375–376.
- Provost, M. W. The dispersal of Aedes taeniorhynchus. III. Study methods for migrating exodus. Mosq. News 20: 148–161.

- 1962. Gressitt, J. L. and M. K. Gressitt. An improved Malaise trap. Pac. Insects 4: 87–90.

 ——. Townes, H. Design for a Malaise trap. Proc. Entomol. Soc. Wash. 64: 253–262.
- 1963. Juillet, J. A. A comparison of 4 types of traps used for capturing flying insects. Can. J. Zool. 41: 219–223.
- 1965. Breeland, S. G. and E. Pickard. The Malaise trap—an efficient and unbiased mosquito collecting device. Mosq. News 25: 19–21.
- ——. Butler, G. D., Jr. A modified Malaise insect trap. Pan-Pac. Entomol. 41: 51–53.
- ——. Chanter, D. O. The Malaise trap. Entomol. Rec. 77: 224–226, pl. 3.
- ——. Evans, R. C. and D. F. Owen. Measuring insect flight activity with a Malaise trap. Mich. Acad. Sci. Arts Lett. 50: 89–94.
- ——. Marston, N. Recent modifications in the design of Malaise insect traps with a summary of the insects represented in collections. J. Kans. Entomol. Soc. 38: 154–162.
- Moczar, L. Trap automatic for capture of flying insects. Folia Entomol. Hung. 20: 213–221.
- Smith, G. E., S. G. Breeland, and E. Pickard. The Malaise trap—a survey tool in medical entomology. Mosq. News 25: 398–400.
- 1966. Pruess, K. P. and N. C. Pruess. Note on a Malaise trap for determining flight direction of insects. J. Kans. Entomol. Soc. 39: 98–102.
- Southwood, T. R. E. Ecological methods with particular reference to the study of insect populations. Methuen, London, xviii, 391 pp. Reference on p. 192. See also 2nd edition, 1978.
- 1967. Gunstream, S. E. and R. M. Chew. A comparison of mosquito collection by Malaise and miniature light traps. J. Med. Entomol. 4: 495–496.
- Olkowski, W. and J. R. Anderson. Relationships between host attack rates and CO₂-baited Malaise trap catches of certain tabanid species. Proc. Annu. Meet. Am. Mosq. Control Assoc. 35: 77.
- 1968. Easton, E. R., M. A. Price, and O. H. Graham. The collection of biting flies in West Texas with Malaise and animal-baited traps. Mosq. News 28: 465–469.
- Evans, F. C. and W. W. Murdoch. Taxonomic composition, trophic structure, and seasonal occurrence in a grassland insect community. J. Animal Ecol. 37: 259–273.
- Geijskes, D. C. Insect collecting in Suriname with the help of "Malaise" traps. *In* Studies on the fauna of Suriname and other Guyanas, No. 39. Natuurwet. Studierkring Suriname Ned. Antillen 48: 101–109.
- ——. Owen, D. F. Species diversity and seasonal abundance in tropical Sphingidae (Lepidoptera). Proc. R. Entomol. Soc. Lond. (A) 44: 162–168.
- (1969. Gillies, M. T. The ramp-trap, an unbaited device for flight studies of mosquitoes. Mosq. News 29: 189–193.)
- Graham, P. A comparison of sampling methods for adult mosquito populations in central Alberta, Canada. Quaest. Entomol. 5: 217–261.
- (——. Herting, B. Tent window traps used for collecting tachinids (Dipt.) at Delémont, Switzerland. Techn. Bul. Commonw. Inst. Biol. Control 12: 1–19.)
- 1970. Hoy, J. B. Trapping the stable fly by using CO₂ or CO as attractants. J. Econ. Entomol. 63: 792–795.
- Matthews, R. W. and J. R. Matthews. Malaise trap studies of flying insects in a New York mesic forest. I. Ordinal composition and seasonal abundance. J. N.Y. Entomol. Soc. 78: 52–59.
- Nishida, T. et al. A handbook of field methods for research on rice-stem-borers and their natural enemies. Int. Biol. Programme Handb. No. 14. Blackwell Sci. Publ., Oxford.
- ——. Roberts, R. H. Tabanidae collected in a Malaise trap baited with CO₂. Mosq. News 30: 52-53.

- -----. Roberts, R. H. Color of Malaise trap and the collection of Tabanidae. Mosq. News 30: 567–571.
- 1971. Matthews, R. W. and J. R. Matthews. The Malaise trap: Its utility and potential for sampling insect populations. Mich. Entomol. 4: 117–122. Imprint as of 1971, but appeared in 1972.
- ——. Roberts, R. H. The seasonal appearance of Tabanidae as determined by Malaise trap collections. Mosq. News 31: 509–512.
- ——. Roberts, R. H. Effect of amount of CO₂ on collection of Tabanidae in Malaise traps. Mosq. News 31: 551–558.
- 1972. Anderson, J. R. and J. B. Hoy. Relationship between host attack rates and CO₂-baited insect flight trap catches of certain *Symphoromyia* species. J. Med. Entomol. 9: 373–393.
- Blume, R. R. et al. Trapping tabanids with modified Malaise traps baited with CO₂. Mosq. News 32: 90–95.
- ——. Roberts, R. H. The effectiveness of several types of Malaise traps for the collection of Tabanidae and Culicidae. Mosq. News 32: 542–547.
 - ——. Townes, H. A light-weight Malaise trap. Entomol. News 83: 239–247.
- 1973. Bailly-Choumara, H. Étude comparative de differentes techniques de récolté de moustiques adultes (Diptera, Culicidae) faite au Maroc, en zone rurale. Bul. Soc. Sci. Nat. Phys. Maroc 53: 135–187.
- ——. Corbet, P. S. and H. V. Danks. Seasonal emergence and activity of mosquitoes (Diptera: Culicidae) in a high-arctic locality. Can. Entomol. 105: 837–872.
- 1974. Anderson, J. R., W. Olkowski, and J. B. Hoy. The response of tabanid species to carbon dioxide baited insect flight traps in northern California USA (Diptera: Tabanidae). Pan-Pac. Entomol. 50: 255–268.
- Roberts, R. H. Relationship between the amount of CO₂ and the collection of Tabanidae in Malaise traps. Mosq. News 35: 220–223.
- 1975. Roberts, R. H. Relationship between the amount of CO₂ and the collection of Tabanidae in Malaise traps. Mosq. News 35: 150–154.
- Roberts, R. H. Influence of trap screen age on collections of tabanids in Malaise traps. Mosq. News 35: 538–539.
- Schroeder, M. H., J. C. Mitchell, and J. M. Schmid. Modifications in the Malaise insect trap. U.S. For. Serv., Rocky Mt. For. Range Exp. Stn., Res. Note RM-299, 2 pp.
- Ticehurst, M. and R. Reardon. Malaise trap: A survey tool for collecting the adult stage of gypsy moth, *Lymantria dispar*. Melsheimer Entomol. Ser. 18, 4 pp.
- ——. Yano, K. et al. Preliminary evaluation on the use of a modified Malaise trap in paddy fields. Mushi 48: 125–144.
- 1976. Acuff, V. R. Trap biases influencing mosquito collecting. Mosq. News 36: 173-176.
- ——. Roberts, R. H. The comparative efficiency of six trap types for the collection of Tabanidae (Diptera). Mosq. News 36: 530–535.
- ——. Service, M. W. Mosquito ecology: Field sampling methods. Applied Sci. Publ., London, xii, 583 pp. References on pp. 177–181, 284, 285, 308, 317–319.
- Tallamy, D. W., E. J. Hansens, and R. F. Denno. A comparison of Malaise trapping and aerial netting for sampling a horsefly and deerfly community. Environ. Entomol. 5: 788–792.
- 1977. Martin, J. E. H. The insects and arachnids of Canada. Part 1. Collecting, preparing, and preserving insects, mites, and spiders. Agriculture Canada, Ottawa, 182 pp.
- (———. Steyskal, G. C. History and use of the McPhail trap. Fla. Entomol. 60: 11–16. A line on p. 14, after line 15, was omitted: "record that I have seen of the use of the trap for other".)

- . Thompson, P. H. and J. W. Holmes, Jr. The Gressitt trap for collecting Tabanidae (Diptera). Proc. Entomol. Soc. Wash. 79: 350–353. This actually refers to the Malaise trap as modified by Gressitt and Gressitt (1962). The change of name is confusing; it is commonly referred to by most users as a Malaise trap.
- 1978. Bram, R. A., ed. Surveillance and collection of arthropods of veterinary importance. U.S. Dep. Agric. Agric. Handb. No. 518, v, 125 p. References, pp. 40, 47–48, 78.
- . Merritt, R. W. et al. Collecting, sampling, and rearing methods for aquatic insects. *In Merritt*, R. W. and K. W. Cummins, An introduction to the aquatic insects of North America. Kendall/Hunt Publ. Co., Dubuque, Iowa. xii, 441 pp. Reference on pp. 13–28, a modified Malaise trap is shown in illustration (Fig. 3.26) only (p. 25).
- (——. Rogoff, W. M. Methods for collecting eye gnats (family Chloropidae). U.S. Dep. Agric. Sci. Educ. Admin. Agric. Rev. and Man. ARM-W-2 [iii], 27 pp. Does not mention Malaise trap, but describes a 'lean-to' trap used by Burgess, 1935.)
- Walker, T. J. Migration and re-migration of butterflies through north peninsular Florida: quantification with Malaise traps. J. Lepid. Soc. 32: 178–190.