

Last updated: 12 August 2025

ALYSSON

Alysson Panzer, 1806:169. Type species: *Pompilus spinosus* Panzer, 1801, designated by Morice and Durrant, 1915:406.

Alysson Jurine, 1807:195. Type species: *Pompilus spinosus* Panzer, 1801, by monotypy.

Keys: Handlirsch, 1895:836 (Nearctic species); Kokujev, 1906 (Russia); Tsuneki, 1968h:4 (South and East Asia); Wu and Zhou, 1987:214 (China); Nemkov, 2014a (Russia and adjacent countries); Girish Kumar, Sheela, and Sharma, 2020 (India); Zou, Ma, Chen, and Li, 2025 (China).

Review of biology: Kazenas, 2001b:216.

1. *annulipes* Cameron

Alysson annulipes Cameron, 1897:21, "♀" = ♂. Holotype: ♂, India: Maharashtra: Pune (OXUM). – Rothney, 1903:105 (India: West Bengal: Barrackpore); R. Turner, 1917e:184 (India; Cameron's type is a male); Tsuneki, 1968h:5 (in key to *Alysson* of southern and eastern Asia); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Girish Kumar, Sheela, and Sharma, 2020:271 (in key to Indian *Alysson*).

2. *attenuatus* Wu and Zhou

Alysson attenuatus Wu and Zhou, 1987:218 and 221, ♂. Holotype: ♂, China: Yunnan: Xishuangbanna: Menglun (Inst. Zool. Beijing). – Wu and Zhou, 1987:215 (in key to *Alysson* of China), 1996a:132 (in revision in Economic Insect Fauna of China); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Zou, Ma, Chen, and Li, 2025:586 (in key to *Alysson* of China).

3. *bengalensis* Girish Kumar

Alysson bengalensis Girish Kumar in Girish Kumar, Sheela, and Sharma, 2020:273, ♀. Holotype: ♀, India: West Bengal: Neora Valley National Park at 27°04'56"N 88°42'02"E (Zoological Survey of India, Western Ghat Region, Kozhikode, Kerala, India). – Girish Kumar, Sheela, and Sharma, 2020:271 (in key to Indian *Alysson*)

4. *caeruleus* Wu and Zhou

Alysson caeruleus Wu and Zhou, 1987:216 and 221, ♀. Holotype: ♀, China: Yunnan: Xishuangbanna: Menglun (Inst. Zool. Beijing). – Wu and Zhou, 1987:215 (in key to *Alysson* of China), 1996a:131 (in revision in Economic Insect Fauna of China); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Zou, Ma, Chen, and Li, 2025:587 (in key to *Alysson* of China).

5. *cameroni* Yasumatsu and Masuda

Alysson cameroni Yasumatsu and Masuda, 1932:53, ♀, ♂. Holotype: ♂, Japan, Honshu: Mount Mitsustôge near Kai (Entomol. Lab., Kyushu Univ., Fukuoka, Japan). – Evans, 1959b:150 (description of larva by Yasumatsu and Masuda, 1932); Iwata, 1960:209 (number of ovarioles and oocytes, egg size); Tsuneki, Iida, Hirai, and Sabi, 1964:60 (Japan: Aomori Prefecture: Towada); Tsuneki, 1965c:26 (in key to Bembicinae of Japan and Korea); Evans, 1966d:29 (review of known behavioral data); Itami, 1968a:16 (Japan: Yamagata Prefecture: foot of Mount Iide); Tsuneki, 1968h:6 (in key to *Alysson* of South and East Asia); Haneda, 1968a:46 (Japan); Tsuneki, 1969d:16 (nesting habits), 1969e:26 (Japan: specimens in Osaka Museum of Natural History); Haneda, 1970:13 (Japan: Akita Prefecture); Tsuneki, 1970b:9 (central Japan); Suda, 1973:123 (Japan: Yamanashi Prefecture); Nambu, 1975b:71 (Ja-

pan: Saitama Prefecture); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Kazenas, 1980e:85 (Russia: Far East); Nemkov *in* Nemkov, Kazenas, Budrys, et Antropov, 1995:455 (in key to Sphecidae of Russian Far East); Miyatake, 1996:103 (specimens in Hiroshi Aoki collection); Haneda, Nosaka, Tano, Kurokawa, and Murota, 2004:37 (Japan: Gifu Prefecture); Suda, 2004:46 (Japan: Yamanashi Prefecture); Haneda, Nozaka, Tano, Kurokawa, and Murota, 2005:54 (Japan: Gifu Prefecture); Negoro, Haneda, and Tano, 2005:49 (Japan: Toyama Prefecture); Haneda, Nozaka, Tano, Kurokawa, and Murota, 2006a:29 (Japan: Toyama Prefecture); Terayama, 2006:2, 4 (in key to Japanese Bembicinae); Evans and O'Neil, 2007:23 (summary of information on nesting habits); Ohkusa, 2007:35 (Japan: central Honshu: Aichi Prefecture); Nemkov, 2008b:23 (in catalog of Crabronidae of Asiatic Russia); Ohkusa and Tajima, 2008:75 (Japan: central Honshu: Aichi Prefecture); Nemkov, 2009b:127 (in new catalog of Sphecidae and Crabronidae of Asiatic Russia); Ohkusa and Tajima, 2010:13 (Japan: Honshu Island: Shiga Prefecture); Nemkov, 2012b:119 (review of known prey), 2012c:445 (in catalog of Crabronidae of Russian Far East), 2014a:277, 278 (in key to *Alysson* of Russia and adjacent countries), 278 (in revision of *Alysson* of Russia and adjacent countries); Mokrousov and Danilov, 2017:57 (Russia: Kuril Islands: Island of Kunashir); Nemkov 2017a:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae), 2017b:219 (in catalog of Bembicinae of Russia); Itami, 2018:7 (Japan: Niigata Prefecture), 12 (color photographs); Mita, 2021:43 (Japan: Kumamoto Prefecture), 44-45 (comparison with *Alysson japonicas*, *A. monticola*, *A. pertheesi*, *A. ratzeburgi*, *A. spinosus*, *A. taiwanus*, and *A. verhoeffi*).

6. *carinatus* Wu and Zhou

Alysson carinatus Wu and Zhou, 1987:219 and 222, ♂. Holotype: ♂, China: Yunnan: Weixi: Pantiange (Inst. Zool. Beijing). – Wu and Zhou, 1987:215 (in key to *Alysson* of China, as *Carinatus*), 1996a:135 (in revision in Economic Insect Fauna of China); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Zou, Ma, Chen, and Li, 2025:587 (in key to *Alysson* of China).

7. *conicus* Provancher

Alysson conicus Provancher, 1889:271, ♀. Lectotype: ♂ [sic], Canada: Ontario: Ottawa (Laval Univ.), designated by Gahan and Rohwer, 1917:305. – W. Fox, 1894b:88 (in revision of North American *Alysson*); Handlirsch, 1895:832 (original description and Fox's redescription translated into German); Dalla Torre, 1897:563 (in catalog of world Sphecidae); Ashmead, 1899d:329 (in checklist of North American Crabronidae); Harrington, 1902:223 (Canada: Ontario: Ottawa); Bradley, 1920:130 (New York, Ontario); Mickel, 1918b:348 (in catalog of Nebraskan Sphecidae); J.Ch. Bradley, 1928:1014 (in catalog of New York Sphecidae); Krombein *in* Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera); Krombein, 1963f:276 (Maryland: Plummers Island near Washington, D.C.); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1979b:1685 (in catalog of North American Hymenoptera); O'Brien and Kurczewski, 1979:435 (nesting behavior); Finnamore, 1982:202 (in Sphecidae Fauna of southern Quebec); O'Brien and Kurczewski, 1982b:225 (nesting behavior); Radović, 1985:65 (sting apparatus analyzed); Steiner, 1986:109 (references to publications on nesting habits); Kurczewski and Miller, 1991:254 (Pennsylvania: Erie County); Buck, 2004:27 (Canada: in checklist of Crabronidae of Ontario); Evans and O'Neil, 2007:24 (summary of information on nesting habits); Buck and Bennett, 2025:207 (in checklist of Sphecidae *sensu lato* of Canada, Alaska, and Greenland: Canada: Manitoba, New Brunswick, Ontario, and Quebec: no specific locality).

As *Euspongia bipunctata* (now *Ceropales bipunctata* Say, Pompilidae): A. Brown, 1934:263 (Canada: Ontario: Timagami), corrected to *Alysson conicus* by Buck, 2004:77.

8. *costai* de Beaumont

As *Alysson ratzeburgi*: Costa, 1867c:77 and 1869b:93 (in revision of Italian Sphecidae: Piemonte), corrected to *Alysson costai* by de Beaumont, 1953i:8; possibly also De Stefani Perez, 1895:228 (in catalog of Sicilian Hymenoptera).

Alyson sp.?: de Beaumont, 1951g:317 (Italy: Emilia), corrected to *Alysson costai* by de Beaumont, 1953i:8.

Alysson costai de Beaumont, 1953i:8, ♀, ♂. Holotype: ♀, Italy: Piemonte: no specific locality (Napoli). – de Beaumont and Bytinski-Salz, 1959:108 (Israel); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Pagliano, 1990:67 (in catalog of Italian Sphecidae), 1992:31 (in key to Italian Nyssoninae); Negrisola *in* Minelli, Ruffo, and La Posta, 1995b:3 (in catalog of Italian fauna); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:123 (in Sphecidae Fauna of Western Europe); Pagliano and Negrisola, 2005:112 (in Sphecidae Fauna of Italy); Strumia, Pagliano, and Gayubo, 2012:56 (Italy: Toscana: Riserva di San Rossore); Nemkov, 2017:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae); Gayubo, Bitsch, Barbier, and Jacobs *in* Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:193 (in Sphecidae Fauna of Europe).

9. *cyanotaeniatu*s Zou and Chen

*Alysson cyanotaeniatu*s Zou and Chen *in* Zou, Ma, Chen, and Li, 2025:587, ♂. Holotype: ♂. China: Guizhou Province: Fanjingshan Mountain: Jinding (Yunnan Agricultural University, Kunming, China). – Zou, Ma, Chen, and Li, 2025:587 (in key to *Alysson* of China).

10. *erythrothorax* Cameron

Alysson erythrothorax Cameron, 1902c:291, ♀. Holotype or syntypes: ♀, India: Himachal Pradesh: Simla (BMNH). – Tsuneki, 1968h:5 (in key to *Alysson* of South and East Asia); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Girish Kumar, Sheela, and Sharma, 2020:271 (in key to Indian *Alysson*).

11. *flavomaculatus* Cameron

Alysson flavomaculatus Cameron, 1901c:314, ♂. Holotype or syntypes: ♂, USA: New Mexico: Santa Fe Mountains (depository?). – R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1979b:1685 (in catalog of North American Hymenoptera). – *As Solierella flavomaculata*?: Krombein *in* Muesebeck, Krombein, and Townes, 1951:942 (tentative new combination, in catalog of North American Hymenoptera).

12. *guichardi* Arnold

Alysson guichardi Arnold, 1951:120, ♀, ♂. Syntypes: Ethiopia: Ghedo (BMNH). – R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae).

13. *guignardi* Provancher

Alysson guignardi Provancher, 1887:271, ♀, ♂. Lectotype: ♀, Canada: Quebec: Bécancour (Laval Univ.), designated by Gahan and Rohwer, 1917:305. – W. Fox, 1894b:88 (in revision of North American *Alysson*); Handlirsch, 1895:833 (original description translated into German); Dalla Torre, 1897:564 (in catalog of world Sphecidae); Ashmead, 1899d:329 (in checklist of North American Crabronidae); Harrington, 1902:223 (Canada: Ontario: Ottawa); J.Ch. Bradley, 1926:175 (New York: Lloyd-Cornell Reservation near McLean), 1928:1014 (in catalog of New York Sphecidae); Krombein *in* Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1979b:1685 (in catalog of North American Hymenoptera); Finnamore, 1982:203 (in Sphecidae Fauna of southern Quebec); Radović, 1985:65 (sting apparatus analyzed); Ahlstrom, 1995:112 (in checklist of insects of North Carolina); Finnamore, 1997:892 (Canada: Yukon); Sugar *et al.*, 1998:16 (Canada: southern Ontario); Buck, 2004:27 (in checklist of Crabronidae of Ontario, Canada); Ratzlaff, 2016:23 (Canada: British Columbia, after Buck, 2004); Buck and Bennett, 2025:207 (in checklist of Sphecidae *sensu lato* of Canada, Alaska, and Greenland: Canada: Nova Scotia, Ontario, Quebec, first records from Manitoba, New Brunswick, and Prince Edward Island: no specific locality).

Alysson interstitialis Cameron, 1902e:375, ♂. Holotype or syntypes: ♂, USA: New Mexico: Santa Fé (depository?). Synonymized with *Alysson guignardi* by R. Bohart *in* R. Bohart and Menke, 1976:458. – Krombein *in* Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera).

Alysson petiolatus Cameron, 1902e:374, ♂. Holotype or syntypes: ♂, USA: New Mexico: Santa Fé (depository?). Synonymized with *Alysson guignardi* by R. Bohart *in* R. Bohart and Menke, 1976:458. – Krombein *in* Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera).

14. *guillarmodi* Arnold

Alysson guillarmodi Arnold, 1944:20, ♀ (as *Guillarmodi*, incorrect original capitalization). Holotype: ♀, Lesotho: Mamathes (SAM). – R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae).

15. *himachalensis* Girish Kumar

Alysson himachalensis Girish Kumar in Girish Kumar, Sheela, and Sharma, 2020:273, ♀. Holotype: ♀, India: Himachal Pradesh: Catchment Ala at 32°31'28"N 76°00'07"E (Zoological Survey of India, Western Ghat Region, Kozhikode, Kerala, India). – Girish Kumar, Sheela, and Sharma, 2020:271 (in key to Indian *Alysson*)

16. *japonicus* Tsuneki

Alysson japonicus Tsuneki, 1977b:26, ♀, ♂. Holotype: ♂, Japan: Ishikawa Prefecture: Mount Haku near Ichinose (originally K. Tsuneki coll., now Hyogo Mus.). – Tano, 1994a:7 (holotype in K. Tsuneki coll.); Hashimoto and Nakanishi, 1997:23 (holotype transferred to Hyogo Mus.); Suda, 2004:46 (Japan: Yamanashi Prefecture); Haneda, Nozaka, Tano, Kurokawa, and Murota, 2005:54 (Japan: Gifu Prefecture), 2006a:29 (Japan: Toyama Prefecture); Nambu, 2006:247 (Japan: Tokyo: Imperial Palace garden); Terayama, 2006:2, 4 (in key to Japanese Bembicinae); Nemkov, 2014a:277, 278 (in key to *Alysson* of Russia and adjacent countries), 280 (in revision of *Alysson* of Russia and adjacent countries), 2017:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae); Itami, 2018:7 (Japan: Niigata Prefecture), 12 (color photographs).

17. *labrisectus* Zou and Li

Alysson labrisectus Zou and Li in Zou, Ma, Chen, and Li, 2025:587, ♂. Holotype: ♂. China: Ningxia Autonomous Region: Erlonghe Forest Farm: Botanical Garden (Yunnan Agricultural University, Kunming, China). – Zou, Ma, Chen, and Li, 2025:587 (in key to *Alysson* of China).

18. *madecassus* Arnold

Alysson madecassus Arnold, 1945:50, ♀. Holotype: ♀, Madagascar: Behara (MNHH). – Leclercq, 1960a:100 (Madagascar); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Pulawski, 2003b:795 (in checklist of Malagasy Sphecidae); Madl, 2014a:939 (in catalog of Ampulicidae, Crabronidae, and Sphecidae of Malagasy sub-region, with synonymy and locality records).

19. *maracandensis* Radoszkowski

Alysson maracandensis Radoszkowski, 1877:34, ♀ (as *Maracandensis*, incorrect original capitalization). Holotype or syntypes: ♀, Uzbekistan: Samarkand (ZMMU). – Handlirsch, 1888a:252 (original description copied); Dalla Torre, 1897:564 (in catalog of world Sphecidae); Kokujev, 1906:213 (in revision of Russian Alyssontini); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Kazenas, 1978b:94 (in key to Sphecidae of Kazakhstan and Central Asia), 2001b:43 (in checklist of Sphecidae of Kazakhstan and Central Asia); Nemkov, 2014a:277 (in key to *Alysson* of Russia and adjacent countries), 280 (in revision of *Alysson* of Russia and adjacent countries), 2017:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae).

Alysson incertus Radoszkowski, 1877:34, ♂. Holotype or syntypes: ♂, Uzbekistan: Peishambe (ZMMU). Synonymized with *Alysson maracandensis* by Handlirsch, 1888a:244 (as tentative synonym of *Alysson tricolor*), 252 (as tentative synonym of *Alysson maracandensis*); Dalla Torre, 1897:564 (in catalog of world Sphecidae); Kokujev, 1906:212 (in revision of Russian Alyssontini).

20. *melleus* Say

Alyson melleus Say, 1837:380, ♀. Holotype or syntypes: ♀, USA: Indiana: no specific locality (lost). – F. Smith, 1856:373 (in catalog of Hymenoptera in British Museum); Le Conte, 1859b:762 (original description copied); Cresson, 1862:235 (in catalog of North American Hymenoptera); Packard, 1867:422 (revision); Cresson, 1887:280 (in catalog of North American Hymenoptera); Handlirsch, 1888a:253 (in revision of world Bembicinae); Cameron, 1890a:70 (Mexico: Huastec); W. Fox, 1894b:87 (in revision of North American *Alysson*); Handlirsch,

1895:831 (additional bibliography); Dalla Torre, 1897:564 (in catalog of world Sphecidae); Ashmead, 1899d:329 (in checklist of North American Crabronidae); Bridwell, 1899:209 (Kansas: no specific locality); J. Smith, 1900:521 (in list of insects of New Jersey); Harrington, 1902:223 (Canada: Ontario: Ottawa); Hartman, 1905:41 (nesting habits); H. Smith, 1908b:342 (in revision of Nebraskan Sphecidae); J. Smith, 1910:680 (in new list of insects of New Jersey); Rohwer, 1916b:654 (in catalog of Hymenoptera of Connecticut); Mickel, 1918b:347 (in catalog of Nebraskan Sphecidae); Rau and Rau, 1918:140 (nesting habits); Washburn, 1919:216 (in list of Hymenoptera of Minnesota); J.Ch. Bradley, 1920:130 (New York, Tennessee, Texas, Quebec); Rau, 1922:18 (USA: Missouri: St. Louis; nesting in the ground); Rohwer *in* Viereck, 1925:654 (in key to Sphecidae of Connecticut); J.Ch. Bradley, 1928:1014 (in catalog of New York Sphecidae); Brimley, 1938:446 (North Carolina: Bryson City, Lake Waccamaw, Raleigh, Tarboro); K. Cooper, 1950:105 (Massachusetts: Cape Code: Woods Hole); Krombein, 1950a:269 (North Carolina: Dare County), 1951a:143 (Virginia: Dunn Loring; visiting tulip-tree honeydew); Krombein *in* Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera); Krombein, 1952a:94 (USA: Virginia: Westmoreland State Park), 1953a:333 (North Carolina: Kill Devil Hills); Krombein and Evans, 1955:233 (Florida); Evans and Lin, 1956b:41 (description of larva); Krombein, 1958f:192 (in supplement to catalog of North American Hymenoptera: description of larva by Evans and Lin, 1956a, reported); Evans, 1959b:150 (additional larval characters); Krombein, 1963f:276 (Maryland: Plummers Island near Washington, D.C.); Evans, 1966d:17 (nesting habits), 1968:1344 (prey records); Kurczewski and Kurczewski, 1971b:335 (prey record); Thorp, 1973:89 (Oklahoma: Lake Texoma region, used as prey by asilid *Promachus bastardii* (Macquart)); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); L. Davis, 1978:216 (North Carolina: Kill Devil Hills, data from Krombein, 1953a); Krombein, 1979b:1685 (in catalog of North American Hymenoptera, nesting sites, list of prey); Finnamore, 1982:204 (in Sphecid Fauna of southern Quebec); Piek and Spanjer, 1986:179 (in list of Sphecidae with known prey); Steiner, 1986:109 (references to publications on nesting habits); Kurczewski and Acciavatti, 1990:61 (New York: Cayuga County); Spofford, Kurczewski, and Downes, 1989:257 (nest parasite: *Phrosinella fulvicornis* (Coquillett), a miltogrammine fly); Spofford and Kurczewski, 1992:999 (species of miltogrammine parasites listed); Ahlstrom, 1995:112 (in checklist of insects of North Carolina); Kurczewski, 1998d:250 (pine barrens in upstate New York); Amarante, 2002:7 (in catalog of Neotropical Crabronidae); Buck, 2004:27 (in checklist of Crabronidae of Ontario, Canada); Evans and O'Neil, 2007:25 (summary of information on nesting habits); Nemkov, 2012b:119 (review of known prey); Buck and Bennett, 2025:207 (in checklist of Sphecidae *sensu lato* of Canada, Alaska, and Greenland: Canada: Manitoba, Ontario, Quebec: no specific locality).

21. *monticola* Tsuneki

Alysson monticola Tsuneki, 1977b:29, ♂, ♀. Holotype: ♂, Japan: Saitama Prefecture: Karisaka Pass (originally K. Tsuneki coll., now Hyogo Mus.). – Tano, 1994a:7 (holotype in K. Tsuneki coll., as *monticola monticola*); Hashimoto and Nakanishi, 1997:23 (holotype transferred to Hyogo Mus.); Haneda, Nosaka, Tano, Kurokawa, and Murota, 2004:37 (Japan: Gifu Prefecture, as *monticola monticola*); Haneda, Nozaka, Tano, Kurokawa, and Murota, 2005:54 (Japan: Gifu Prefecture, as *monticola monticola*), 2006a:29 (Japan: Toyama Prefecture, as *monticola monticola*); Nagase, 2006a:55 (Japan: Kanagawa Prefecture: Mount Komagatake near Hakone, as *monticola monticola*); Terayama, 2006:2, 4 (in key to Japanese Bembicinae); Nemkov, 2007b:74 (first record from Russia: Kuril Islands: Island of Kunashir), 2007c:997 (in Appendix to Key to identification of Sphecidae of Far East of Russia), 2008b:24 (in catalog of Crabronidae of Asiatic Russia), 2009b:127 (in new catalog of Sphecidae and Crabronidae of Asiatic Russia), 2011:95 (known only from Kuril Islands and Japan), 2012c:445 (in catalog of Crabronidae of Russian Far East), 2014a:277, 278 (in key to *Alysson* of Russia and adjacent countries), 280 (in revision of *Alysson* of Russia and adjacent countries), 2017a:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae); Itami, 2018:8 (Japan: Niigata Prefecture), 12 (color photographs); J.-K. Kim, 2020a:2 (South Korea: locality records).

As *Alysson ratzeburgi*: Nemkov in Nemkov, Kazenas, Budrys, and Antropov, 1995:455 (in key to Sphecidae of Russian Far East), corrected to *Alysson monticola* by Nemkov, 2007:74.

ssp. ezoensis Tsuneki

Alysson monticola ezoensis Tsuneki, 1977b:32, ♀, ♂. Holotype: ♀, Japan: Hokkaido: Sapporo (originally K. Tsuneki coll., now Hyogo Mus.). – Tano, 1994a:7 (holotype in K. Tsuneki coll.); Hashimoto and Nakanishi, 1997:23 (holotype transferred to Hyogo Mus.); Nemkov, 2017b:219 (in catalog of Bembicinae of Russia).

22. nigrilabius Wu and Zhou

Alysson nigrilabius Wu and Zhou, 1987:219 and 222, ♂. Holotype: ♂: China: Sichuan: Emei (Inst. Zool. Beijing). – Wu and Zhou, 1987:215 (in key to *Alysson* of China), 1996a:134; Hua, 2006:277 (in list of Chinese insects, geographic distribution); Nemkov, 2017:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae); Zou, Ma, Chen, and Li, 2025:587 (in key to *Alysson* of China).

23. ocellatus de Beaumont

Alysson ocellatus de Beaumont, 1967a:327, ♀, ♂. Holotype: ♀, Turkey: Amasya (BMNH). – R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Ljubomirov and Yildirim, 2008:62 (in catalog of Crabronidae of Turkey); Yildirim, Ljubomirov, and Lelej, 2014:5 (distribution in Turkey by biogeographic provinces), 19 (is an endemic of Turkey); Koçak and Kemal, 2015:136 (in checklist of Hymenoptera of Turkey); Nemkov, 2017:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae); Kaplan and Yildirim, 2021:22 (in catalog of Crabronidae of Turkey), 2023:1679 (in checklist of Sphecidae *sensu lato* of Turkey).

24. oppositus Say

Alyson oppositus Say, 1837:380, ♀, ♂. Syntypes: USA: Indiana: no specific locality (lost). – F. Smith, 1856:373 (in catalog of Hymenoptera in British Museum); Le Conte, 1859b:761 (reprint of Say, 1837:380); Cresson, 1862:235 (in catalog of North American Hymenoptera); Packard, 1867:421 (in revision of North American Bembicinae); Provancher, 1883b:636 (in revision of Canadian Hymenoptera); Cresson, 1887:280 (in catalog of North American Hymenoptera); Handlirsch, 1888a:249 (in revision of world Bembicinae); Ashmead, 1890:32 (in checklist of Hymenoptera of Colorado); W. Fox, 1894b:87 (in revision of North American *Alysson*); Handlirsch, 1895:831 (additional bibliography); Dalla Torre, 1897:564 (in catalog of world Sphecidae); Ashmead, 1899d:329 (in checklist of North American Crabronidae); Bridwell, 1899:209 (Kansas: no specific locality); J. Smith, 1900:521 (in list of insects of New Jersey); Harrington, 1902:223 (Canada: Ontario: Ottawa); J. Smith, 1910:680 (in new list of insects of New Jersey); Rohwer, 1916b:654 (in catalog of Hymenoptera of Connecticut); Mickel, 1918b:348 (in catalog of Nebraskan Sphecidae); Washburn, 1919:216 (in list of Hymenoptera of Minnesota); J.Ch. Bradley, 1920:130 (New York, Quebec); Britton, 1920:338 (in checklist of insects of Connecticut); Rohwer in Viereck, 1925:654 (in key to Sphecidae of Connecticut; Thompson, Connecticut); J.Ch Bradley, 1928:1015 (in catalog of New York Sphecidae); Brimley, 1938:446 (North Carolina: Raleigh, Swannanoa); Krombein in Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera); Krombein, 1952a:94 (USA: Virginia: Westmoreland State Park), 1963f:276 (Maryland: Plummers Island near Washington, D.C.); Evans, 1966d:28 (habitat, prey); Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1979b:1685 (in catalog of North American Hymenoptera); Finnamore, 1982:205 (in Sphecid Fauna of southern Quebec); Ahlstrom, 1995:112 (in checklist of insects of North Carolina); Amarante, 2002:7 (in catalog of Neotropical Crabronidae); Buck, 2004:27 (in checklist of Crabronidae of Ontario, Canada); Evans and O'Neil, 2007:26 (summary of information on nesting habits); Buck and Bennett, 2025:207 (in checklist of Sphecidae *sensu lato* of Canada, Alaska, and Greenland: Canada: Manitoba, New Brunswick, Ontario, Quebec, first record from Nova Scotia: no specific locality).

25. pertheesi Gorski

Alysson pertheesi Gorski, 1852:178, ♀ (as *Perthéesi*, incorrect original capitalization and diacritic mark). Syntypes: Lithuania: Vilnius (mainly lost except one specimen in ZMHU). – Handlirsch, 1888a:251 (in revision of world

Bembicinae); Kohl, 1893a:28 (Austria: Niederösterreich: Wiener Prater and Donauauen); Dalla Torre, 1897:564 (in catalog of world Sphecidae); Mocsáry, 1897:81 (Hungary: Budapest); Kokujev, 1906:216 (in revision of Russian Alyssontini); Schmiedeknecht, 1907:225 (in key to Hymenoptera of Central Europe); de Gaulle, 1908:120 (in catalog of French Hymenoptera); Noskiewicz, 1918:137 (Galicia, now western Ukraine: Brzuchowice and Hołosko near Lwów, Janów); Berland, 1925d:96 (in Sphecid Fauna of France); Koornneef, 1925:364 (first record from Netherlands: Gelderland: Dieren); Bouwman, 1927:94 (in revision of Sphecidae of Netherlands); Schmiedeknecht, 1930:686, 687 (in keys to Hymenoptera of North and Central Europe); Szulczewski, 1947:112 (Poland: Wielkopolski National Park 15 km south of Poznań); Šnoflák in Zavadil and Šnoflák, 1948:78, 79 (in key to Sphecidae of Czechoslovakia, not yet found in Czechoslovakia); Szulczewski, 1950:107 (Poland: Wielkopolski National Park 15 km south of Poznań); Bajári, 1957a:94, 95 (in key to Hungarian Sphecidae, as *Parthéesi*); Benno, 1958b:10 (in review of Dutch Sphecidae); Noskiewicz and Puławski, 1960:83, 84 (in key to Polish Sphecidae); Tsuneki, 1965c:25 (in key to Bembicinae of Japan and Korea; first record from Asia); Benno, 1967:34 (in checklist of Sphecidae of Netherlands); Itami, 1967:4 (nesting habits), 1968b:17 (Japan); Nambu, 1968b:27 (Japan: Saitama Prefecture); Tsuneki, 1968h:5 (in key to *Alysson* of South and East Asia); Itami, 1969:46 (Japan: Niigata Prefecture); Oehlke, 1970:705 (not recorded from Democratic Germany); Tsuneki, 1970b:9 (central Japan); Tumšs, 1970:77 (first record from Latvia); Balthasar, 1972:352 (in Sphecid Fauna of Czechoslovakia: not yet found in country); Tsuneki and Itami, 1973:13 (Japan: Niigata Prefecture; color variation); Tsuneki, 1974a:361 (Korea); Nambu, 1975b:72 (Japan: Saitama Prefecture); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Kolesnikov, 1977:318 (Russia: Bryansk Oblast'); Pulawski, 1978:212 (in key to Sphecidae of European part of USSR); Lefebvre, 1979:11 (in checklist of Sphecidae of Netherlands), 56 (Netherlands: distribution map); Kazenas, 1980e:85 (Russia: Far East); Mityaev, Matesova, Kulenova, Kazenas, and Zhdanko, 1980:119 (Russia: Siberia: Altayskiy Kray: basin of Aley River); Tsuneki, 1982b:18 (known from Korea); Dollfuss, 1983a:75 (Austria: endangered in Niederösterreich), 80 (Austria: endangered in Tirol), 1983b:20 (in catalog of Sphecidae of Austria); Paik, 1985:203 (in list of Sphecidae of Korea); Nemkov, 1986:106 (Russia: Irkutsk Oblast'); Tyrner, 1987:282 (Slovakia); Pádr in Šedivý, 1989a:167 (in checklist of Czechoslovakian Sphecidae); Jacobs and Oehlke, 1990:204 (unknown from Democratic Germany); Schwammberger and Priesner, 1990:550 (Austria: Kärnten); Day, 1991:xii (in summary of European Endangered Hymenoptera Lists); Dollfuss, 1991:181 (in key to Sphecidae of North and Central Europe); Tsuneki, 1991:199 (Korea); Budrys, 1992:28 (in list of Sphecidae of Lithuania); Dollfuss, 1994:97 (endangered in Austria); Nemkov in Nemkov, Kazenas, Budrys, et Antropov, 1995:455 (in key to Sphecidae of Russian Far East); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:124 (in Sphecid Fauna of Western Europe); Celary, 1997:60 (in checklist of Animals of Poland); Shlyakhtenok, Blinov, and Belyavskaya, 1997:152 (Belorussia: Pripyatskiy National Park); Zettel, Gross, and Mazzucco, 2001:79 (Austria: Wien, not collected after 1870); Shlyakhtenok and Skibinska, 2002:34 (Belorussia: no specific locality); Klein and Lefebvre in Peeters, van Achterberg, et al., 2004:357 (in Wasp and Ant Fauna of Netherlands); Skibińska in Bogdanowicz, Chudzicka, Pilipiuk, and Skibińska, 2004:354 (in catalog of Polish Crabronidae); van Achterberg and Peeters, 2004:32 (in checklist of Sphecidae of Netherlands); Wiśniowski, 2004:50 and 62 (in checklist of Polish Sphecidae); Nemkov, 2005a:158 (Russia: Island of Sakhalin); Terayama, 2006:2, 4 (in key to Japanese Bembicinae); Evans and O'Neil, 2007:26 (summary of information on nesting habits); Jacobs, 2007:159, 160 (in key to Crabronidae of Germany, not yet found in Germany); Nemkov, 2007b:74 (Russia: Kuril Islands: Island of Kunashir), 2007c:997 (in Appendix to Key to identification of Sphecidae of Far East of Russia); Vepřek and Straka, 2007:207 (in catalog of Sphecidae of Czech Republic and Slovakia, known from Slovakia only); Nemkov, 2008b:24 (in catalog of Crabronidae of Asiatic Russia); Mokrousov, 2009:278 (Russia: Republic Mari El); Nemkov, 2009b:127 (in new catalog of Sphecidae and Crabronidae of Asiatic Russia); Bitsch, 2010:108 (in supplement to vol. II of Faune de France, 1997: recent records reviewed; the only French specimen comes from Var: La Roquebrussanne); Mokrousov, 2010a:61 (Russia: Kostroma and Nizhniy Novgorod oblast's, Mari El Republic: no specific localities); Baghirov, 2011b:141 (Russia: recorded from Altayskiy Kray by Kazenas, Mityaev, Matesova, Kulenova, and Zhdanko, 1979); Józán, 2011:183 (in checklist of Sphecidae

sensu lato of Hungary); Nemkov, 2012c:445 (in catalog of Crabronidae of Russian Far East); Shlyakhtenok, 2012b:91 (Belorussia: Beresina National Reserve: overgrowing dry meadow), 2013:221 (in annotated catalog of aculeate wasps of Belorussia); J.-K. Kim, 2014:420 (in catalog of Sphecidae *sensu lato* of Korean Peninsula); Nemkov, 2014a:277, 278 (in key to *Alysson* of Russia and adjacent countries), 280 (in revision of *Alysson* of Russia and adjacent countries), 2017a:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae), 2017b:219 (in catalog of Bembicinae of Russia, as *pertheesi pertheesi*); Itami, 2018:8 (Japan: Niigata Prefecture), 13 (color photographs); Jacobs and Liebig, 2018:136 (Russia: Irkutsk Oblast': Sarma); Gayubo, Bitsch, Barbier, and Jacobs in Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:194 (in Sphecidae Fauna of Europe); J.-K. Kim, 2020a:3 (South Korea: locality records); Olszewski, Wiśniowski, and Ljubomirov, 2021:83 (in commented list of Sphecidae *sensu lato* of Poland); Zou, Ma, Chen, and Li, 2025:586 (in key to *Alysson* of China).

Alysson festivus Mocsáry, 1879:129, ♀ (as *festivum*, incorrect original termination). Holotype or syntypes: ♀, Hungary: Budapest (TMB). Synonymized with *Alysson pertheesi* by Handlirsch, 1888a:250. – Haris, 2016:16 (in list of species described by S. Mocsáry from Carpathian Basin).

Alysson fuscatus var. *jaroslavensis* Kokujev, 1906:216, ♀. Holotype: ♀, Russia: Yaroslavl' (originally N. Kokujev coll., now ZIN). Synonymized with *Alysson pertheesi* by Nemkov, 2014a:280. – **As *Alysson spinosus jaroslavensis***: R. Bohart and Menke, 1976:458 (new status, in checklist of world Sphecidae); Shlyakhtenok and Skibinska, 2002:34 (new status, Belorussia: no specific locality). – **As *Alysson jaroslavensis***: Pulawski, 1978:212, 213 (new status, in key to Sphecidae of European part of USSR); Pankevich, Belyavskaya, Molchanova, Shlyakhtenok, and Blinov, 1989:95 (Belorussia: Berezinskiy Nature Reserve); Blagoveshchenskaya, 1994:91 (Russia: Ul'yanovsk Oblast', misspelled *jaroslavensis*); Voblenko, Gorobchishin, and Nesterov, 1996:15 (Ukraine: Polesye Region); Shlyakhtenok, Blinov, and Belyavskaya, 1997:152 (Belorussia: Pripyatskiy National Park); Mokrousov, 2002:150 (Russia: Nizhniy Novgorod Oblast': Kerzhenskiy Nature Reserve); Pankov, 2013:228 (Russia: Ivanovo Oblast').

Alysson harbinensis Tsuneki, 1967e:7, ♀. Holotype: ♀, China: Heilongjiang: Harbin (RMNH). Synonymized with *Alysson pertheesi* by Nemkov, 2014a:280. – Tsuneki, 1968h:5 (in key to *Alysson* of South and East Asia); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Wu and Zhou, 1987:214 (in key to *Alysson* of China), 1996a:136 (in revision in Economic Insect Fauna of China); Hua, 2006:277 (in list of Chinese insects, geographic distribution).

Alysson pertheesi forma *typica* Tsuneki and Itami, 1973:13. Not available (Article 16).

Alysson pertheesi forma *nigra* Tsuneki and Itami, 1973:13, ♀. Syntypes: ♀, Japan: Niigata Prefecture: Shidaihana (Itami coll., K. Tsuneki coll.). Not available (Article 16).

ssp. *formosanus* Tsuneki

Alysson pertheesi formosanus Tsuneki, 1968h:4, ♀. Holotype: ♀, Taiwan: Nantou Prefecture: Puli (originally K. Tsuneki coll., now USNM). – Tsuneki, 1977c:6 (Taiwan: Chiayi Prefecture: Chuchi; comparison with Japanese populations of *Alysson pertheesi*); Zou, Ma, Chen, and Li, 2025:586 (in key to *Alysson* of China). – **As *Alysson formosanus***: R. Bohart and Menke, 1976:458 (new status, in checklist of world Sphecidae); Wu and Zhou, 1987:214 (in key to *Alysson* of China); Nuhn and Menke, 1994:25 (holotype transferred to USNM); Porter, Stange, and Wang, 1999:13 (in checklist of Sphecidae of Taiwan); Hua, 2006:277 (in list of Chinese insects, geographic distribution).

26. *picteti* Handlirsch

Alysson picteti Handlirsch, 1895:830, ♂ (as *Picteti*, incorrect original capitalization). Syntypes: Algeria: Constantine Province: Col des Oliviers in El Kantour Mountains (NHMW). – Dalla Torre, 1897:564 (in catalog of world Sphecidae); Schmiedeknecht, 1900:233 (Tunisia, description of ♀); Graeffe, 1906:455 (Tunisia: Tunis area); von Schulthess, 1926b:216 (Tunisia); de Beaumont, 1950f:403 (Algeria: Maison Carrée, now El Harrach, a suburb of Alger); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Nemkov, 2017:Table p. 1 (in zoogeographic

analysis of Palearctic Bembicinae); Ben Khedher, Yildirim, and Braham, 2022:2 (in checklist of Bembecinae of Tunisia).

27. *radiatus* W. Fox

Alysson radiatus W. Fox, 1894b:87, ♀, ♂. Lectotype: ♀, USA: Nevada: no specific locality (ANSP), designated by Cresson, 1928:48. – Handlirsch, 1895:832 (original description translated into German); Dalla Torre, 1897:564 (in catalog of world Sphecidae); Ashmead, 1899d:329 (in checklist of North American Crabronidae); H. Smith, 1908b:341 (in revision of Nebraskan Sphecidae); Mickel, 1918b:348 (in catalog of Nebraskan Sphecidae); Brimley, 1938:446 (North Carolina: Raleigh, Shining Rock); Krombein *in* Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera); Evans, 1970:493 (Wyoming: Jackson Hole); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1979b:1685 (in catalog of North American Hymenoptera); Ahlstrom, 1995:112 (in checklist of insects of North Carolina); Ratzlaff, 2016:23 (first record from Canada: British Columbia: Southern Interior), 30 (British Columbia: Oliver, Osoyoos, Penticton); Buck and Bennett, 2025:207 (in checklist of Sphecidae *sensu lato* of Canada, Alaska, and Greenland: Canada: British Columbia, first record from Alberta: no specific locality).

28. *ratzeburgi* Dahlbom

Alyson ratzeburgi Dahlbom, 1843:145, ♀, ♂ (as *Ratzeburgi*, incorrect original capitalization). Lectotype: ♂, Norway: Ostre Naess Vaerdaliae (Lund), designated by de Beaumont, 1953h:197. – Dahlbom, 1845:474 (in key to European Nyssoninae), 1847:279 (nest and prey); Erichson, 1848:137 (Dahlbom's biological observations reported); Eversmann, 1849:388 (Russia: Kazan' Province); Taschenberg, 1858:86 (in key to Sphecidae of Germany); Brischke, 1862:100 (Germany: Preussen Province, now Poland: Pomorze Region); Taschenberg, 1866:199 (in revision of Hymenoptera of Germany); nec Costa, 1867c:77 and 1869b:93 (= *Alysson costai*); Kirchner, 1867:219 (in catalog of European Hymenoptera); Thomson, 1870:249 (in revision of Swedish wasps), 1874:249 (in revision of Sphecidae of Scandinavia); nec Marquet, 1875:204 (= *Alysson spinosus*); Siebke, 1880:67 (Norway); Friese, 1883:202 (Germany: Thüringen: Rothensteiner Felsen); Brischke, 1887:87 (Germany: Seeresen near Karthaus, now Poland: Dzierżąno near Kartuzy), 1888b:97 (Germany: Preussen Province, now Poland: Pomorze Region); Handlirsch, 1888a:242 (in revision of world Bembicinae); F. Morawitz, 1893a:112 (Russia: Karelian Isthmus); Westerlund, 1893:26 (Russia: Karelian Republic: northern shore of Lake Ladoga); De Stefani Perez, 1895:228 (in catalog of Sicilian Hymenoptera, may actually be *Alysson costai*); Handlirsch, 1895:830 (one of Dahlbom's specimens examined); Dalla Torre, 1897:564 (in catalog of world Sphecidae); Strand, 1898b:95 (Norway); Ferton, 1901a:104 (prey in Algeria, determination probably incorrect), 107 (preying on Fulgoridae); Adlerz, 1904:138 (known prey: cicadellids); Aurivillius, 1904:275 (in revision of Swedish Sphecidae); Kokujev, 1906:214 (in revision of Russian Alyssontini); Schmiedeknecht, 1907:224, 225 (in key to Hymenoptera of Central Europe); de Gaulle, 1908:120 (in catalog of French Hymenoptera); Adlerz, 1910:42 (Sweden, nesting habits); nec Dittrich, 1911b:24 (= *Alysson spinosus*); Morice, 1911:10 (Algeria: Constantine); Kohl, 1912a:63 (Austria: Tirol: Innsbruck; and Südtirol: Bad Ratzes, now Italy: Trentino-Alto Adige: Razzes); Stöckhert, 1919:40 (Germany: Franken); Enslin, 1922c:246 (Germany: Bayern: Fürth); Lundblad, 1924:22 (Sweden: Lappland: Lycksele); Berland, 1925d:97 (in Sphecidae Fauna of France); Schütze, 1925:113 (Germany: Lausitz Region); Bouwman, 1927:94 (in key to Sphecidae of Netherlands, not yet found in Netherlands); Barovskij, 1929:149 (Russia: north of Sankt Petersburg); Gaunitz, 1929:199 (Sweden: Lapland); Schmiedeknecht, 1930:686, 687 (in keys to Hymenoptera of North and Central Europe); Pulkkinen, 1931:66 (in revision of Sphecidae of Finland); Giordani Soika, 1932a:22 (Italy: Lido di Venezia, determination probably incorrect); E. Nielsen, 1933:302 (summary of nesting habits); Jansson, 1934:290 (Sweden); Forsius *in* Forsius and Hellén, 1935:12 (in checklist of Hymenoptera of Finland); Grandi, 1935a:110 (Italy); Móczár, 1938:64 (Hungary), 1939:5 (Hungary: Jászberény); Zavdil, 1939:129 (Czech Republic: Morava: Ubušin); Schneid, 1941:1042 (Germany: Bayern: Bamberg area); Giner Marí, 1943a:153 (in Sphecidae Fauna of Spain, determination probably incorrect); de Beaumont, 1945:475 (Switzerland: Valais); Szulczewski, 1947:112 (Poland: Wielkopolski

National Park 15 km south of Poznań); Šnoflák in Zavadil and Šnoflák, 1948:68 (in key to Sphecidae of Czechoslovakia); Aerts, 1950a:51 (Germany: Nordrhein-Westfalen: Rheidter Wertchens near Köln); Szulczewski, 1950:107 (Poland: Wielkopolski National Park 15 km south of Poznań); Leininger, 1953:18 (Germany: Baden-Württemberg: Hinterzarten); Grandi, 1954:234 (Italy); Valkeila, 1954:87 (Finland: Lappeenranta); Hellén, 1955a:125 (Finland: 16 localities, including Terijoki, now Russia: Zelenogorsk); Vogrin, 1955:36 (Yugoslavia, determination probably incorrect); Bajári, 1957a:94 (in key to Hungarian Sphecidae); Wolf, 1959b:29 (Germany: Nassau); Noskiewicz and Puławski, 1960:83, 84 (in key to Polish Sphecidae, not yet found in Poland); Valkeila, 1962b:67 (southwestern Finland); de Beaumont, 1964c:63 (in Sphecidae Fauna of Switzerland); Valkeila, Brander, and Käpylä, 1964:75 (Finland: southern Finland: Forssa); Tsuneki, 1965c:25 (in key to Bembicinae of Japan and Korea; first record from Asia); Bonelli, 1966c:216 (Italy: Trentino–Alto Adige: Razzes); Evans, 1966d:28 (review of known behavioral data); Gauss, 1967:553 (Germany: Baden-Württemberg: südlicher Schwarzwald); J. Heinrich, 1967:74 (Germany: Bayern: Kahl, Lohr, Obernau); Niedl and Deneš, 1967:83 (Czech Republic: Třeboň area); T. Iida, 1968:21 (Japan); Nambu, 1968:22 (Japan); Tano, 1968:33 (Japan); Tsuneki, 1968h:5 (in key to *Alysson* of South and East Asia); Itami, 1969:46 (Japan: Niigata Prefecture); Schmidt, 1969:163 (Germany: Rhein-Main area: Wilbrands-Schneiß south of Darmstadt); Oehlke, 1970:705 (Democratic Germany); Tsuneki, 1970b:9 (central Japan); Tumšs, 1970:77 (first record from Latvia); Balthasar, 1972:351 (in Sphecidae Fauna of Czechoslovakia); Kofler, 1972:106 (Austria: eastern Tirol); Niedl and Deneš, 1972:153 (Czech Republic: Třeboň area); Suda, 1973:123 (Japan: Yamanashi Prefecture); Lomholdt, 1975c:221 (in Sphecidae Fauna of Fennoscandia and Denmark); Nambu, 1975b:72 (Japan: Saitama Prefecture); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Eck, 1976:90 (Democratic Germany); Pulawski, 1978:212 (in key to Sphecidae of European part of USSR); Schmidt, 1979b:308 (Germany: Baden-Württemberg); Casolari and Casolari Moreno, 1980:125 (specimens in M. Spinola collection, Torino); Dollfuss and Ressler, 1981:317, 324 (Austria: Niederösterreich: Scheibbs District); Axelsson, 1982:79 (Sweden); Leclercq, 1982e:100 (Norway: Nord-Trøndelag and Hedmark provinces); Tsuneki, 1982b:18 (known from Korea); Dollfuss, 1983a:76 (Austria: endangered in Niederösterreich), 78 (Austria: endangered in Oberösterreich), 80 (Austria: endangered in Tirol), 1983b:20 (in catalog of Sphecidae of Austria); Paik, 1985:203 (in list of Sphecidae of Korea); Westrich and Schmidt, 1985b:110 (Germany: extinct in Baden-Württemberg); Nemkov, 1986:106 (Russia: Irkutsk Oblast); Vikberg, 1986:78 (Finland); Dollfuss, 1987:23 (Austria); Wu and Zhou, 1987:215 (in key to *Alysson* of China, as *rozenburgii*); Nilssen, 1989:48 (Sweden); Pádr in Šedivý, 1989a:167 (in checklist of Czechoslovakian Sphecidae); Pankevich, Belyavskaya, Molchanova, Shlyakhtenok, and Blinov, 1989:95 (Belorussia: Berezinskiy Nature Reserve); Scobiola-Palade, 1989:88 (Romania: delta of Danube); Jacobs and Oehlke, 1990:122 and 204 (not collected in Democratic Germany after 1960); Pagliano, 1990:67 (in catalog of Italian Sphecidae); Schwammbberger and Priesner, 1990:550 (Austria: Kärnten); Day, 1991:xii (in summary of European Endangered Hymenoptera Lists); Dollfuss, 1991:181 (in key to Sphecidae of North and Central Europe); Pagliano, 1992:31 (in key to Italian Nyssoninae); K. Weber, 1992:86 (Germany: Bayern: Bamberg District); Dollfuss, 1994:101 (endangered in Austria); Janssen and Kaluza, 1995:& (Germany: extinct in Sachsen); Negrisolo in Minelli, Ruffo, and La Posta, 1995b:3 (in catalog of Italian fauna); nec Nemkov in Nemkov, Kazenas, Budrys, et Antropov, 1995:455 (= *Alysson monticola*); Shlyakhtenok, 1995:193 (Belorussia: Berezinskiy Nature Reserve); Jakubzik, 1996:251 (Germany: Köln); Minoranskiy and Shkuratov, 1996:81 (Russia: Rostov Oblast); Schmid-Egger, Schmidt, and Doczkal, 1996:376 (Germany: Baden-Württemberg: last collected in 1952); Wu and Zhou, 1996a:136 (in revision in Economic Insect Fauna of China); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:125 (in Sphecidae Fauna of Western Europe); Pagliano and Alma, 1997:179 (known prey listed); Schmidt and Schmid-Egger, 1997:5 (in checklist of German Sphecidae); Schnee, 1997:100 (Germany: Sachsen: Nature Reserve Heidewald- und Moorgebiet; first collected in Sachsen since 1919); Shlyakhtenok, Blinov, and Belyavskaya, 1997:151 (Belorussia: Pripyatskiy National Park); F. Burger, Saure, and Oehlke, 1998:26 (Germany: in list of Sphecidae of Brandenburg), 36 (probably endangered); K. Weber, 1998:133 (Germany: Bayern: Bamberg area); Mandery, 1999:104 (Germany: Bayern: Coburg area); Pauly, 1999:111 (in catalog of Belgian

aculeates); M. Schwarz, 1999:88 (Austria: Oberösterreich: city of Linz area); Blösch, 2000:359 (in Sphecidae Fauna of Germany); Ohl et al., 2001:137 (distribution in Germany by provinces); Shlyakhtenok and Skibińska, 2002:34 (Belorussia: no specific locality); Söderman and Vikberg, 2002:53 (in list of Aculeata of Finland); Mandery, Kraus, Voith, Wickl, Scheuchl, Schubert, and Warncke, 2003:63 (in catalog of wasps and bees of Bayern, Germany); Wickl, Voith, Mandery, Weber, Kraus, Bausenwein, and Blösch, 2003:195 (Germany: Bayern: no longer endangered); Esser, Jakubzik, and Sonnenburg, 2004:18 (Germany: Nordrhein-Westfalen); Esser, Jakubzik, Sonnenburg, and Woydak, 2004:261 (Germany: in checklist of Sphecidae of Nordrhein-Westfalen); Shkuratov, 2004a:76 (Russia: Rostov Oblast': Migulinskaya), 2004b:166 (Russia: Rostov Oblast': Gosudarstvennyi Muzei-Zapovednik M.A. Sholokhova); Skibińska in Bogdanowicz, Chudzicka, Pilipiuk, and Skibińska, 2004:354 (in catalog of Polish Crabronidae); Wiśniowski, 2004:50 and 62 (in checklist of Polish Sphecidae); Burger, 2005:31 (Germany: Thüringen); Kofler, 2005:182 (Austria: Tirol: Lienz District: Tessenberg in Heinfels Common); Pagliano and Negrisolo, 2005:112 (in Sphecidae Fauna of Italy); Straka, 2005a:393 (extinct in Czech Republic); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Liebig, 2006:37 (Germany: Sachsen: Muskauer Heide); Jacobs, 2007:159, 160 (in key to Crabronidae of Germany); Pettersson, Hellqvist, and Gabrielsson, 2007:7 (Sweden: Västerbotten: Ammarnäs); Saure, 2007:81 (Germany: in checklist of Aculeata of Land Brandenburg); Vepřek and Straka, 2007:207 (in catalog of Sphecidae of Czech Republic and Slovakia, known from Bohemia and Moravia only); Nemkov, 2008b:24 (in catalog of Crabronidae of Asiatic Russia); U. Frommer, 2009a:26 (Germany: in checklist of Sphecidae of central Hessen, locality records, 30 (currently not occurring in Hessen); Humala and Polevoi, 2009:63 (Russia: Republic of Karelia east of Lake Onega); Nemkov, 2009b:127 (in new catalog of Sphecidae and Crabronidae of Asiatic Russia); Ødegaard, Sverdrup-Thygeson, Hansen, Hanssen, and Öberg, 2009:70 (Norway: locality records); Bitsch, 2010:108 (in supplement to vol. II of Faune de France, 1997: revue of recent records from Europe); Mokrousov, 2010a:61 (Russia: Kirov and Nizhniy Novgorod oblast's, Tatarstan: no specific localities), 2010b:100 (Russia: Kirov and Nizhniy Novgorod oblast's, Tatarstan: no specific locality); Mokrousov and Zryanin, 2010:98 (Russia: needs to be protected, Novgorod Oblast'); Rudoiskatel', 2010:152 (Russia: central Ural Mountains); Schmid-Egger, 2010a:13 (in red list of Aculeata of Germany: very rare); Shorenko and Konovalov, 2010:13 (Ukraine: Donetsk); Burger, 2011a:286 (Germany: in red list of Sphecidae *sensu lato* of Thüringen); Józán, 2011:183 (in checklist of Sphecidae *sensu lato* of Hungary); Tischendorf, Frommer, and Flügel et al., 2011:40 (Germany: extinct in Hessen: last collected in 1961); Nemkov, 2012c:445 (in catalog of Crabronidae of Russian Far East); Saure, 2012:30 (Germany: Brandenburg: Döberitzer Heide at 52.521°N 13.064°E); Shlyakhtenok, 2012b:91 (Belorussia: Beresina National Reserve: overgrowing dry meadow), 2013:221 (in annotated catalog of aculeate wasps of Belorussia); Çubuk and Gülmez, 2013:34 (first record from Turkey: Tokat Province: 10 km of Ormandibi road); Danilov and Odintseva, 2013:333 (Russia: Western Siberia: 65 km north-northeast of Tobol'sk at 58°45'N 68°74'E); A. Scholz and Liebig, 2013:16 (Germany: Sachsen: in list of Sphecidae *sensu lato*); Shlyakhtenok, 2013:221 (in annotated catalog of aculeate wasps of Belorussia); Bitsch, 2014:398 (record by Saure, 2012 reported); Hellqvist, Abenius, and Norén, 2014:90 (in catalog of Ampulicidae, Sphecidae, and Crabronidae of Sweden, by provinces); J.-K. Kim, 2014:421 (in catalog of Sphecidae *sensu lato* of Korean Peninsula); Mokrousov and Vafin, 2014:55 (Russia: Republic of Tatarstan); Nemkov, 2014a:277, 278 (in key to Alysson of Russia and adjacent countries), 281 (in revision of Alysson of Russia and adjacent countries); Pagliano, 2014:118 (specimens in M. Spinola collection, Torino, are *Alysson ratzeburgi*); Yıldırım, Ljubomirov, and Lelej, 2014:5 (distribution in Turkey by biogeographic provinces); Koçak and Kemal, 2015:136 (in checklist of Hymenoptera of Turkey, possibly a misdetermination of *Alysson costai*); Paukkunen and Kozlov, 2015:62 (Russia: Murmansk Oblast'); Blažej, Macek, and Trýzna, 2016:111 (Czech Republic: České Švýcarsko = Bohemian Switzerland National Park); Nemkov, 2017a:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae), 2017b:219 (in catalog of Bembicinae of Russia); Gayubo, Bitsch, Barbier, and Jacobs in Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:195 (in Sphecidae Fauna of Europe); Kejval, Blažej, and Erhart, 2020:91 (western Czech Republic); J.-K. Kim, 2020a:3 (South Korea: previous locality records cited); Kaplan and Yildirim, 2021:22 (in catalog of Crabronidae of Turkey,

possibly a misdetermination of *Alysson costai*; Olszewski, Wiśniowski, and Ljubomirov, 2021:83 (in commented list of Sphecidae *sensu lato* of Poland); Ben Khedher, Yildirim, and Braham, 2022:2 (first record from Tunisia: Ben Arous: Hammam Lif); Kaplan and Yildirim, 2023:1679 (in checklist of Sphecidae *sensu lato* of Turkey); Zou, Ma, Chen, and Li, 2025:587 (in key to *Alysson* of China).

29. *ruficollis* Cameron

Alyson [sic] *ruficollis* Cameron, 1898b :25, "♂" = ♀ (as *ruficolle*, incorrect original termination). Holotype: ♀, Sri Lanka: Kandy (OXUM). – Tsuneki, 1968h:5 (in key to *Alysson* of South and East Asia); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1985a:11 (Sri Lanka; redescription, description of ♂); Girish Kumar, Sheela, and Sharma, 2020:271 (in key to Indian *Alysson*).

30. *seyrigi* Arnold

Alysson seyrigi Arnold, 1945:49, ♀ (as *Seyrigi*, incorrect original capitalization). Holotype: ♀, Madagascar: Perinet (MNHN). – R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Pulawski, 2003b:795 (in checklist of Malagasy Sphecidae); Madl, 2014a:939 (in catalog of Ampulicidae, Crabronidae, and Sphecidae of Malagasy subregion, with synonymy and locality records).

31. *sichuanensis* Wu and Zhou

Alysson sichuanensis Wu and Zhou, 1987:219 and 221, ♀. Holotype: ♀, China: Sichuan: Wenchuan Wolun (Inst. Zool. Beijing). – Wu and Zhou, 1987:215 (in key to *Alysson* of China), 1996:134 (in revision in Economic Insect Fauna of China); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Nemkov, 2017:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae); Zou, Ma, Chen, and Li, 2025:587 (in key to *Alysson* of China).

32. *spinosus* (Panzer)

Pompilus spinosus Panzer, 1801, Heft 80:17, [♀]. Holotype or syntypes: ♀, Germany: no specific locality (depository?). Synonymized with *Alyson bimaculatus* by Jurine, 1807:196. – Peets, 1909-1911:67 (*Pompilus spinosus* of Panzer = *Alyson fuscatus*). – **As *Alyson spinosus***: Jurine, 1807:196 (new combination, the only species of *Alysson* listed) and pl. 10, Fig. 21 (illustration of habitus); Lepeletier de Saint Fargeau and Audinet-Serville, 1825:50 (redescription); Stephens, 1829:362 (in catalog of British insects), 1829b:33 (in checklist of British insects); Vander Linden, 1829:91 (bibliographic references, possibly a synonym of *Alysson bimaculatus*, Belgium: Bruxelles, Germany); Lepeletier de Saint Fargeau, 1845:87 (in revision of world Hymenoptera); A. Costa, 1864b:104 (specimen from Torino, Italy, in Napoli Museum), 1869a:17 (Italy: Piemonte); Dours, 1874:142 (in catalog of Hymenoptera of France, as *spinosum*); Frei-Gessner, Kohl, and Kriechbaumer, 1882:393 (Jurine's specimens are *Alyson bimaculatus*); Acloque, 1897:84 (in Sphecidae Fauna of France and Algeria); Szczepko and Kowalczyk, 2001:189 (Poland: Kampinos National Park near Warszawa); Kowalczyk, Szczepko, and Świątek, 2002:155 (Poland: Kampinos National Park near Warszawa). – **As *Alysson spinosus***: J.Ch. Bradley, 1919a:64 (new combination, type species of *Alysson* Jurine in Panzer, 1801, and of *Alyson* Jurine, 1807; first of these names was published in paper suppressed in 1939 by International Commission on Zoological Nomenclature in Opinion 135); Lomholdt, 1975c:220 (in Sphecidae Fauna of Fennoscandia and Denmark); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Haeseler, 1977a:23 (northwestern Germany); Leclercq, 1978b:map 1168 (distribution in Belgium); Lefebvre, 1978:137 (Netherlands: Schaesberg); Petit, 1978:86 (Belgium: Vance); Pulawski, 1978:212, 213 (in key to Sphecidae of European part of USSR); Lefebvre, 1979:11 (in checklist of Sphecidae of Netherlands), 57 (Netherlands: distribution map); Schmidt, 1979b:308 (Germany: Baden-Württemberg); Casolari and Casolari Moreno, 1980:125 (specimens in M. Spinola collection, Torino); Dollfuss and Ressler, 1981:324 (Austria: Niederösterreich: Scheibbs District); Gayubo, 1982b:34 (Spain: Salamanca: Valdefuentes de Sangusín); Dollfuss, 1983b:20 (in catalog of Sphecidae of Austria); Gayubo, 1983e:232 (Spain: Salamanca Province); Haack, Tschamtker, and Vidal, 1984:124 (Germany: Hamburg area); Józán, 1985b:60 (Hungary south of Lake Balaton), 81 (floral records), 84 (ecological and zoogeographic characteristics); Radović, 1985:65 (sting apparatus analyzed); Scobiola-Palade, 1985:96 (Ro-

mania: delta of Danube); Brechtel, 1986:60 (Germany: Südpfalz); Gayubo and Heras, 1986:60 (Spain: Segovia Province); Gayubo and Sanza, 1986:72 (Spain: Burgos); Józán, 1986:373 (Hungary: Kiskunság National Park); Skibińska, 1986b:154 (Poland: suburbs of Warszawa), 157 (rural and suburban parks); Riemann, 1987b:32 (Germany: Niedersachsen: Nature Reserve Dünengebiet near Neumühlen); Schmidt and Westrich, 1987:362 (Germany: Mainz area); Wu and Zhou, 1987:215 (in key to *Alysson* of China); Dollfuss, 1988:18, 29 (Austria: Bisamberg near Vienna; Niederösterreich: Oberweiden); Karsai, 1988:101 (Hungary: Kiskunság National Park); Jacobs, 1989:25 (Germany: Mecklenburg-Vorpommern); Pankevich, Belyavskaya, Molchanova, Shlyakhtenok, and Blinov, 1989:95 (Belorussia: Berezinskiy Nature Reserve); Scobiola-Palade, 1989:88 (Romania: delta of Danube); Skibińska, 1989a:204 (Poland: Warszawa area); Gayubo, Asís, and Tormos, 1990b:77 (Spain: Toledo Province); Jacobs and Oehlke, 1990:204 (new records from Democratic Germany); Kuznetzova, 1990:20 (Russia: Voronezh Oblast': Galich'ya Gora Nature Reserve); Pagliano, 1990:67 (in catalog of Italian Sphecidae); Schwammberger and Priesner, 1990:550 (Austria: Kärnten); Barbier, Rasmont, and Wahis, 1990:32 (Belgium: Hainaut); Day, 1991:xii (in summary of European Endangered Hymenoptera Lists); Dollfuss, 1991:181 (in key to Sphecidae of North and Central Europe); Gayubo, Borsato, and Osella, 1991:412 (Italy); Gayubo and Torres, 1991:Table I (Spain: Salamanca; effects of urban pressure); Jacobs, 1991:46 (Germany: Mecklenburg-Vorpommern: Nature Reserve Elbtalldünen); Jakubzik, 1991:75 (Germany: Köln); Józán, 1991:604 (Hungary: Bátorliget Nature Reserve); Kowalczyk, 1991a:89, 93, 97, 99, 101, 107 (Poland: Łódź), 1991b:120 (Poland: Augustów Forest); Negrisolo, 1991:322 (Italy: Treviso and Udine Provinces); Scobiola-Palade, 1991:220 (Romania: Danube delta); Budrys, 1992:28 (in list of Sphecidae of Lithuania); Józán, 1992b:175 (Hungary: Boronka-melléki Protected Area), 1992c:289 (Hungary: Zselic Hills); Pagliano, 1992:31 (in key to Italian Nyssoninae); K. Weber, 1992:86 (Germany: Bayern: Bamberg District); Kuhlmann, 1993:75 (Germany: known from Westfalen); Sorg and Wolf, 1993:59 (Germany: Rheinland-Westfalen: Nature Reserve Die Spey); Torregrosa, Gayubo, Tormos, and Asís, 1993:20 (Spain: Alicante Province); Blagoveshchenskaya, 1994:91 (Russia: Ul'yanovsk Oblast'); Kowalczyk, 1994:73 (Poland: Łagiewniki forest in Łódź); Theunert, 1994c:32 (Germany: summary of records from Niedersachsen and Bremen); Gorobchishin, 1995:18 (Ukraine: Kanev Nature Reserve); Jansen and Kaluza, 1995:7 (Germany: endangered in Sachsen); Józán, 1995:105 (Hungary: projected Duna-Dráva National Park); Krasnobayev et al., 1995:138 (Russia: Zhiguli Hills northwest of Samara); Negrisolo in Minelli, Ruffo, and La Posta, 1995b:3 (in catalog of Italian fauna); Saure and Dürrenfeld, 1995:29 (Germany: Brandenburg: Gabower Hänge near Bad Freienwalde); Scharer, 1995:28 (Austria: Niederösterreich: Donau-Auen); Schmid-Egger, Risch, and Niehuis, 1995:163 (summary of records from Germany: Rheinland-Pfalz State); Gorobchishin, 1996:53 (Ukraine: Kanev Nature Reserve); Jakubzik, 1996:251 (Germany: Köln); Jakubzik and Cölln, 1996b:398 (Germany: Rheinland-Pfalz); Minoranskiy and Shkuratov, 1996:81 (Russia: Rostov Oblast'); Saure, 1996:503 (Germany: Berlin-Pankow: sewage irrigation field); Schmid-Egger, Schmidt, and Doczkal, 1996:376 (Germany: Baden-Württemberg: potentially endangered); Voblenko, Gorobchishin, and Nesterov, 1996:15 (Ukraine: Polesye Region); Witt, 1996:105 (Germany: Brandenburg), 106 (locality records); Woydak, 1996:89 (Germany: Westfalen: summary of nesting habits and distribution); Wu and Zhou, 1996a:136 (in revision in Economic Insect Fauna of China); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:126 (in Sphecid Fauna of Western Europe); Celary, 1997:60 (in checklist of Animals of Poland); Lauterbach, 1997b:64 (Germany: Nordrhein-Westfalen: Bielefeld); Pagliano and Alma, 1997:178, 179 (prey: *Cicadella viridis* (L.), Cicadellidae); K. Schmidt and Schmid-Egger, 1997:5 (in checklist of German Sphecidae); Shlyakhtenok, Blinov, and Belyavskaya, 1997:152 (Belorussia: Pripyatskiy National Park); F. Burger, Saure, and Oehlke, 1998:26 (Germany: in list of Sphecidae of Brandenburg); Dollfuss, Gusenleitner, and Bregant, 1998:521 (Austria: summary of collecting records from Burgenland); Flügel, 1998:59 (Germany: Brandenburg: Unteres Odertal National Park); Gorobchishin, 1998a:50 (Ukraine: Kiev and surroundings); Józán, 1998:312 (Hungary: Duna-Dráva National Park); N. Schneider and Wahis, 1998:19 (Luxemburg: sandstone quarries); H.-J. Schulz, 1998c:112 (Germany: Lausitz: former military training area near Mücka); van der Smissen, 1998:53 (Germany: Schleswig-Holstein, Niedersachsen); K. Weber, 1998:133 (Germany: Bayern: Bamberg area); Gayubo, García, Torres, and González, 1999:91 (Spain: Soria Prov-

ince); Kuhlmann, Wolf, and Woydak, 1999:569 (Germany: endangered in Nordrhein-Westfalen); Ljubomirov, 1999a:22 (Bulgaria); Majzlan, Rychlík, and Deván, 1999:145 (Slovakia: Čenkovská Step Nature Reserve); Mandery, 1999:104 (Germany: Bayern: Coburg area); Neumayer, Schwarz, and Bregant, 1999:223 (Austria: in checklist of Sphecidae of Land Kärnten); Pauly, 1999:111 (in catalog of Belgian aculeates); Saure, 1999:33 (Germany: Berlin-Köpenick); M. Schwarz, 1999:88 (Austria: Oberösterreich: city of Linz area); Blösch, 2000:359 (in Sphecidae Fauna of Germany); Dressler, 2000:30 (Germany: Hessen: Darmstadt area); Esenbekova and Kazenas, 2000:57 (southeastern Kazakhstan: no specific locality); Giachino, Grosso, Marchetti, Pagliano, Scaramozzino, and Vailati, 2000:108 (Greece); Jacobs, 2000:6 (Germany: Land Mecklenburg-Vorpommern: not endangered); Józán, 2000:112 (Hungary: Bakony Mountains); Ljubomirov, 2000:12 (Bulgaria: specimens in N. Nedelkov collection); Pijfers, 2000:4 (Netherlands: Twente Region); Schmid-Egger, 2000b:275 (Germany: Baden-Württemberg: shores of Rhine River); Vicidomini, 2000b:22 (Italy: previously recorded from Campania); Józán, 2001:280 (Hungary: Somogy County); Kazenas, 2001b:43 (in checklist of Sphecidae of Kazakhstan and Central Asia); Ohl et al., 2001:137 (distribution in Germany by provinces); Reemer and van der Meer, 2001:16 (Netherlands: along Almelo-Nordhorn canal); Skibińska, 2001:220 (Poland: Białowieża Primeval Forest); van der Smitten, 2001:58 (Germany: Schleswig-Holstein: not endangered); Zettel, Gross, and Mazzucco, 2001:79 (Austria: Wien); Józán, 2002:54 (Hungary: Mecsek Mountains); Kazenas, 2002a:113 (geographic distribution, collecting localities in Kazakhstan); Reemer and van der Meer, 2002:Bijlage 5 (Netherlands: Gelderland: Zaltbommel area); Shlyakhtenok and Skibinska, 2002:34 (Belorussia: no specific locality); Gorobchishin, 2003:48 (Ukraine: Kiev: botanical garden); Józán, 2003:228 (Hungary: Látványi Puszta Nature Conservation Area); Lukáš, 2003:73 (Slovakia: Bratislava: old fruit orchard); Mandery, Kraus, Voith, Wickl, Scheuchl, Schubert, and Warncke, 2003:63 (Germany: in catalog of wasps and bees of Bayern); Reemer, 2003:62 (Netherlands: Gelderland: Zaltbommel area); von der Heide and Metscher, 2003:101 (Germany: Niedersachsen: Emsland area); Wiśniowski in Dylewska and Wiśniowski, 2003:174 (Poland: Ojców National Park); Beron, 2004:839 (Bulgaria: eastern Rhodope Mountains: Arda Hut, citation of Ljubomirov, 1999:22); Esser, Jakubzik, Sonnenburg, and Woydak, 2004:261 (Germany: in checklist of Sphecidae *sensu lato* of Nordrhein-Westfalen); Klein and Lefebvre in Peeters, van Achterberg, et al., 2004:357 (in Wasp and Ant Fauna of Netherlands); Shkuratov, 2004a:76 (Russia: Rostov Oblast'), 2004b:166 (Russia: Rostov Oblast': Gosudarstvennyi Muzej-Zapovednik M.A. Sholokhova); Skibińska in Bogdanowicz, Chudzicka, Pilipiuk, and Skibińska, 2004:354 (in catalog of Polish Crabronidae); Szczepko and Kowalczyk, 2004:127 (Poland: Kampinos National Park near Warszawa); van Achterberg and Peeters, 2004:32 (in checklist of Sphecidae of Netherlands); Wiśniowski, 2004:50 and 62 (in checklist of Polish Sphecidae); Dubitzky, Blank, and Schönlitzer, 2005:271 (Germany: Bayern: area north of Dachau); Jacobs, 2005a:438 (Bulgaria); Kofler, 2005:183 (Austria: Tirol: Lienz District: Lesach in Kals Common, Lavant, Tschule Alm in Thurn Common); Kowalczyk and Kurzac, 2005:104 (Poland: Botanical Garden in Łódź); Mandery, 2005:173 (Germany: Bayern: Heinberg Nature Reserve near Nürnberg); Pagliano and Negrisola, 2005:113 (in Sphecidae Fauna of Italy); Riemann and Hohmann, 2005:543 (Germany: Niedersachsen: city of Bremen and adjacent area); Saure, 2005:13 (Germany: Berlin); Theunert, 2005a:16 (Germany: in checklist of Sphecidae of Niedersachsen and Bremen); Venne and Bleidorn, 2005:42 (Germany: Nordrhein-Westfalen: Moosheide Nature Reserve); Ziska and Saure, 2005:80 (Germany: Berlin: Biesenhorster Sand); Bock, 2006:41 (Germany: Bayern: Kelheim District: Abensberg sands); Gorobchishin, 2006:135 (Ukraine: several localities); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Liebig, 2006:37 (Germany: Sachsen: Muskauer Heide); Liška, 2006:88 (Slovakia: Váh river valley near Trenčín); Shlyakhtenok, 2006:113 (Belorussia: Polesye Radiological Nature Reserve); Evans and O'Neil, 2007:23 (summary of information on nesting habits); Guldemand, Pijfers, and den Belder, 2007:202 (Netherlands: Gelderland: dairy farm in de Graafschap); Jacobs, 2007:159, 160 (in key to Crabronidae of Germany); Jansen and Kaluza, 2007:22 (Germany: northwestern Sachsen); Józán, 2007:180 (Hungary: Zselic hills); Saure, 2007:81 (Germany: in checklist of Aculeata of Land Brandenburg); Sipos and Móczár, 2007:205 (Hungary: Bács-Kiskun megye: vicinity of Foktő); Smit and Megens, 2007:17 (Netherlands: Noort Brabant: vicinity of Leende); Vago, 2007:174 (France: départements d'Aisne and Pas-de-Calais);

Venne, Bleidorn, and Lauterbach, 2007:300 (Germany: Nordrhein-Westfalen: Bielefeld); Vepřek and Straka, 2007:207 (in catalog of Sphecidae of Czech Republic and Slovakia); Kazenas, 2008a:104 (southeastern Kazakhstan: 5 km east of Chemolgan); Livory, Chevin, Lair, Sagot, and Baldock, 2008:30 (France: Département de la Manche); Ljubomirov and Ban-Calefariu, 2008:203 (Romania: Maramureş District: Repedea, Vişeu Valley); Ljubomirov and Yildirim, 2008:62 (in catalog of Crabronidae of Turkey); Theunert, 2008:7 (Germany: map of distribution in Niedersachsen and Bremen); Baños-Picón, Asís, Gayubo, and Tormos, 2009:310 (Spain: frequency of specimens collected with hand nets and Malaise traps); Debuysse and Vago, 2009:2 (in list of Sphecidae *sensu lato* of northern France); U. Frommer, 2009a:26 (Germany: in checklist of Sphecidae of central Hessen); González, Gayubo, Asís, and Tormos, 2009:625 (Spain: Salamanca and Zamora provinces: Arribes del Duero Natural Park); Straka, Dvořák, and Bogusch, 2009:257 (Czech Republic: Jizerské Mountains and Frýdlant region); van Diemen, 2009:49 (Netherlands: airbase Gilze-Rijen); Bitsch, 2010:108 (in supplement to vol. II of Faune de France, 1997: prey record of Pagliano and Alma reported); Burguet in Durand and Burguet, 2010:32 (France Département d'Ardèche: Ardèche Gorges); Esser, Fuhrmann, and Venne, 2010:35 (Germany: in checklist of aculeates of Nordrhein-Westfalen); Hable, Kraus, von der Dunk, and Wickl, 2010:26 (Germany: Bayern: Sulz- and Ottmaringer Tal); Mokrousov, 2010a:61 (Russia: Kostroma, Nizhniy Novgorod, and Yaroslavl' oblast's, Chuvash Republic, Tatarstan: no specific localities); Saure, 2010b:174 (Germany: Berlin-Reinickendorf: airport Tempel and Flughafensee Protected Area); Schmid-Egger, 2010a:13 (in red list of Aculeata of Germany: relatively common); Schmidt, 2010:131 (Germany: Bayern: collected along Main-Donau canal); Shorenko and Konovalov, 2010:13 (Ukraine: Khar'kov Oblast': Kupyansk); Smetana, Roller, Beneš, Bogusch, Dvořák, Holý, Karas, Macek, Straka, Šima, Tyrner, Vepřek, and Zeman, 2010:100 (Western Slovakia: recorded from Borská Nížina); van der Smissen, 2010a:332 (Germany: Schleswig-Holstein: locality records); Gogala, 2011:26 (Slovenia: Podčetrtek); Józán, 2011:183 (in checklist of Sphecidae *sensu lato* of Hungary); Mokrousov, Berezin, and Egorov, 2011:66 (Russia: Chuvash Republic); Saure, 2011a:5 (Germany: Berlin: former airport Tempelhof), 2011c:197 (Germany: Berlin-Spandau: Fort Hahneberg); Schmid-Egger, 2011c:555 (Austria: Tirol: Tiroler Lech Nature Park); Theunert, 2011a:18 (Germany: in checklist of aculeates of Niedersachsen including city of Bremen); Tischendorf, Frommer, Flügel et al., 2011:66 (Germany: Hessen: endangered); Tripotin, 2011:23 (France: Seine-Maritime: artificial sand site near Sandouville); Ziska and Saure, 2011:108 (Germany: Berlin: Tegeler Fließ valley); Bogusch and Straka, 2012:87 (in Czech Republic nesting in postindustrial habitats); Cölln, Jakubzik, and Kinkler, 2012:24 (Germany: Nordrhein-Westfalen: Leverkusen); Nemkov, 2012b:119 (review of known prey); Prisniy, 2012:50 (Russia: Belgorod Oblast'); Saure, 2012b:228 (Germany: Berlin-Spandau: Spandauer Forst); Shlyakhtenok, 2012b:91 (Belorussia: Beresina National Reserve: overgrowing dry meadow); Strumia, Pagliano, and Gayubo, 2012:56 (Italy: Toscana: Riserva di San Rossore); Tripotin, 2012:23 (France: Seine-Maritime: estuary of Seine river); Dyuzhaeva, 2013:116 (Russia: Samara Oblast': Zhiguli Nature Reserve); A. Scholz and Liebig, 2013:16 (Germany: Sachsen: in list of Sphecidae *sensu lato*); Shlyakhtenok, 2013:222 (in annotated catalog of aculeate wasps of Belorussia); Wiśniowski and Szczepko, 2013:23 (Poland: Kampinos National Park near Warszawa); Bitsch, 2014:398 (France: Ardèche: Saint-Remèze; record from France by Vago, 2007 reported); Mokrousov and Vafin, 2014:55 (Russia: Republic of Tatarstan); Nemkov, 2014a:277, 278 (in key to *Alysson* of Russia and adjacent countries), 282 (in revision of *Alysson* of Russia and adjacent countries); Pagliano, 2014:118 (specimens in M. Spinola collection, Torino, are *Alysson spinosus*); Yildirim, Ljubomirov, and Lelej, 2014:5 (distribution in Turkey by biogeographic provinces); Koçak and Kemal, 2015:136 (in checklist of Hymenoptera of Turkey); P. Rosa and Pagliano, 2015:97 (Italy: Lombardia and Piemonte: Parco del Ticino); Samin and Bagriacik, 2015:71 (Iran: Northern Khorasan: Bojnord at 37°35'N 57°20'E); Simon, Lorthiois, and Macé, 2015:53 (France: alluvial terraces in Eure Department: Courcelles-sur-Seine); Cölln and Jakubzik, 2016:16 (Germany: Rheinland-Pfalz: Vulkaneifel District); Mokrousov and Popov, 2016:566 (Russia: Krasnodarskiy Kray); Saure, 1916c:204 (Germany: center of Berlin: Großer Tiergarten Park); Stolle and Saure, 2016:921 (Germany: in list of Sphecidae of Sachsen-Anhalt); Wiśniowski, 2016:127 (Poland: Ojców National Park); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:32 (Iran: known from North Khorasan

Province); Le Divelec, Chapelin-Viscardi, Larivière, and Adamski, 2017:62 (potential host of *Nysson dimidiatus* and *N. distinguendus*); Nemkov, 2017a:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae), 2017b:219 (in catalog of Bembicinae of Russia); Twerd, Krzyżyński, Waldon-Rudzionek, and Olszewski, 2017:6 (Poland: Inowrocław: nesting in postindustrial soda ash dumping sites); Nicolas and Vago, 2018:14 (France: Pas-de-Calais: Mont Saint-Frieux); Carminati, Mora, and Cretin, 2019:9 (France: in list of Sphecidae *sensu lato* of Franche-Comté); Olszewski and Kowalczyk, 2019:89 (Poland: Pomorze Gdańskie = Gdańsk Pomerania); Gayubo, Bitsch, Barbier, and Jacobs *in* Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and K. Schmidt, 2020:195 (in Sphecidae Fauna of Europe); Cross, Baldock, and Wood, 2021:22 (in catalog of Sphecidae *sensu lato* of Portugal); Kaplan and Yildirim, 2021:22 (in catalog of Crabronidae of Turkey); Olszewski, Wiśniowski, and Ljubomirov, 2021:83 (in commented list of Sphecidae *sensu lato* of Poland); Kaplan and Yildirim, 2023:1679 (in checklist of Sphecidae *sensu lato* of Turkey); Zou, Ma, Chen, and Li, 2025:586 (in key to *Alysson* of China).

Alyson katkovi Kokujev, 1906:214, ♀. Holotype: ♀, Ukraine: Ol'gopol' District: no specific locality (originally N. Kokujev coll., now ZIN). Synonymized with *Alysson spinosus* by Nemkov, 2014a:282. – R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Pulawski, 1978:212 (in key to Sphecidae of European part of USSR).

As *Spheg fuscatus* Fabricius (whose position within Sphecidae is unclear): Panzer, 1798, Heft 51:3 (in revision of German insects, illustration of habitus). – As *Alysson fuscatus*: Panzer, 1806:163 (new combination, in critical review of his previous work); Lepeletier de Saint Fargeau, 1845:88 (new combination, in revision of world Hymenoptera, as *Alyson*); Handlirsch, 1888a:246 (in revision of world Bembicinae); Kohl, 1893a:28 (Austria: Niederösterreich); Krieger, 1894:9 (Germany: Kingdom of Saxony); Handlirsch, 1895:8831 (additional bibliography); Dalla Torre, 1897:563 (in catalog of world Sphecidae); Mocsáry, 1897:81 (Kingdom of Hungary, some localities are in today's Croatia and Romania); Dominique, 1901a:506 (France: Loire-Inférieure, now Loire-Atlantique: Le Pouliguen); de Cobelli, 1903:114 (Italy: Province of Trentino); Antiga and Bofill, 1904:14 (Spain: Cataluña Province); W. Schulz, 1904b:97 (Czechoslovakia); Kokujev, 1906:216 (in revision of Russian Alyssontini); Schmiedeknecht, 1907:224, 225 (in key to Hymenoptera of Central Europe); de Gaulle, 1908:120 (in catalog of French Hymenoptera); Peets, 1909-1911:67 (*Spheg fuscatus* of Panzer = *Alyson fuscatus*); Dittrich, 1911b:24 (Germany: Niederschlesien Province, now Poland: Dolny Śląsk Region); Birula, 1912:540 (Ukraine: Zhytomyr Oblast': Radomyshl'); Schirmer, 1912:167 (Germany: Berlin area); Birula, 1913:540 (Ukraine: Zhytomyr Oblast': Radomyshl'), 1914:386 (Belarus: Luskopol' near Vitebsk); Szilády, 1914:89 (Hungary: Nagyenyed); Bouwman, 1915:2 (Netherlands: Bergen op Zoom, preying on cicadellids); Dubois, 1919:96 (Belgium: Bruxelles: Uccle); O. Meyer, 1919:154 (Germany: Posen Province, now Poland: Poznań Region); A. Wagner, 1920:36 (Germany: Lower Elbe region, not found but can be expected); Dubois, 1922:204, 205 (in key to Belgian Sphecidae); Fahringer, 1922:181 (Turkey); Hedicke, 1922:271 (Germany: Brandenburg: Gross Machnower Weinberg); Berland, 1925d:96 (in Sphecidae Fauna of France); Bischoff, 1925:300 (Latvia: Wezkukkul; Lithuania: Dolny near Vilnius; Poland: Grajewo area); Coulon, 1925:120 (France: Oise); Bouwman, 1926:280 (in list of Sphecidae of Netherlands); Friese, 1926:161 (in key to Sphecidae of Central Europe, short morphological and biological characteristics); Bouwman, 1927:94 (in revision of Sphecidae of Netherlands); F. Werner, 1927:66 (Austria: Niederösterreich: River Kamp valley); Crèvecoeur and Maréchal, 1928:172 (Belgium: Uccle); van der Vecht, 1928:93 (Netherlands: Meijendel sand dunes); Schmiedeknecht, 1930:686, 687 (in keys to Hymenoptera of North and Central Europe); Pulkkinen, 1931:66 (in revision of Sphecidae of Finland, not yet found in Finland); Bondroit, 1932:32 (Belgium: Solbosch); Giordani Soika, 1932a:22 (Italy: Lido di Venezia); Bondroit, 1934:59 (Belgium: Watermael); Bernard, 1935:64 (France: Var: Fréjus area); Crèvecoeur and Maréchal, 1936:253 (in catalog of Belgian Hymenoptera); Zavadil, Šustera, and Bat'a, 1937:195 (in catalog of Sphecidae of Czechoslovakia); Móczár, 1938c:81 (Hungary: Kőszeg Hills); A. Wagner, 1938a:121 (northwestern Germany); Móczár, 1939:5 (Hungary: Jászberény); Balthasar, 1941b:112 (Czech Republic: eastern Moravia); Paul, 1941:44 (Germany: Pommern: Mittel- and Ostpommern: no specific locality now Poland: Pomorze środkowe i wschodnie); Schneid, 1941:1042 (Germany: Bayern: Bamberg area); Giner Marí, 1943a:152 (in Sphecidae Fauna of Spain); Bat'a, 1944a:26 (Czech Republic), 1944b:92 (Czech Republic: Veselí); Guiglia, 1944a:147

(Cyprus: Limassol), 1944b:22 (Italy); Šnoflák, 1944a:144 (Czech Republic: serpentine steppe near Mohelno); Šnoflák in Zavadil and Šnoflák, 1948:78, 79 (in key to Sphecidae of Czechoslovakia); Ceballos, 1949:100 (Spain); de Andrade, 1949:17 (Portugal); Sala de Castellarnau, 1949:89 (Spain: Teruel: Bronchales, as *fuscatum*); Cavo, 1950:36 (France: Département du Nord); Pittioni, 1950:24 (Cyprus); Szulczewski, 1950:107 (Poland: Wielkopolski National Park 15 km south of Poznań, as *fasciatus*); Caruel, 1951:100 (France: Marne Department); Leininger, 1951:117 (Germany: Baden-Württemberg); Bat'a, 1952:167 (Czech Republic); Głowacki, 1953:520 (Poland: Podkova Leśna near Warszawa); Móczár, 1953:310 (Hungary); Bischoff and Stadler, 1954:126 (Germany: Bayern: Romberg); Puławski, 1954:106 (Poland: Błędów); Aerts, 1955:62 (Germany: Rheinland); Vergne, 1955:4 (France: Auvergne); Vogrin, 1955:36 (Yugoslavia); Ceballos, 1956:379 (in catalog of Hymenoptera of Spain); Bajári, 1957a:94, 95 (in key to Hungarian Sphecidae); Zimngiebl, 1957:182 (Germany: Rheinland-Pfalz: Birkenheide, Frankenthal, Leistadt); Benno, 1958b:10 (in review of Dutch Sphecidae); Diniz, 1959:29 (Portugal: Pao de Rei); Olberg, 1959:338-339 (photograph of female carrying cicadellid prey); Wolf, 1959b:29 (Germany: Nassau); Noskiewicz and Puławski, 1960:82, 83 (in key to Polish Sphecidae); Hoop, 1963a:73 (Germany: Schleswig-Holstein: Reher Kratt), 1963b:7 (Germany: Schleswig-Holstein: Lauenburg, Reher Kratt); Lehrer and Scutaru, 1963:283 (Romania: Moldova Pașcani and Mircești in Iași District, Baia in Suceava District); Babiș, 1964:116 (Austria: Land Salzburg); Benno, 1967:34 (in checklist of Sphecidae of Netherlands); Balthasar, M. Hrubant, and E. Hrubant, 1967:174 (Bulgaria: Slanchev Bryag near Nessebar); Niedl and Deneš, 1967:84 (Czech Republic: Třeboň area); Scobiola-Palade, 1967b:47 (Romania), 1968b:142 (Romania: Island of Letea in delta of Danube), 1968c:384 (Romania: Budești, Dăraști, Giurgiu, Tigănești); Dathe, 1969:439 (Germany: Berlin Zoological Garden); Lefebvre, 1969a:78 (Netherlands: Bergen op Zoom); Romanova, 1969:134 (Russia: North Caucasus); Tumšs, 1970:77 (Latvia); Balthasar, 1972:350 (in Sphecidae Fauna of Czechoslovakia); Niedl and Deneš, 1972:153 (Czechoslovakia); Scobiola-Palade, 1972a:150 (Romania: delta of Danube: Caraorman), 1972b:253 (Romania), 1974:144 (Romania: Sărăturile sandbank in Danube delta); Georgiou, 1977:188 (Cyprus); Radović and Krunic, 1979:unpaginated foldout (nesting in sand, foreleg structure); Casolari and Casolari Moreno, 1980:125 (specimens in M. Spinola collection, Torino); Józán, 1981:305 (Hungary: Hortobágy National Park); Skibińska, 1981:357 (Poland: Warszawa-Białoleka), 1982:123 (Poland: Warszawa area); Holtappels, 1987:26 (Germany: Nordrhein-Westfalen: Offender Waldsee); Kowalczyk, 1988b:64 (Poland: environs of Łódź); Pádr in Šedivý, 1989a:167 (in checklist of Czechoslovakian Sphecidae); Pádr, 1990:73 (Czech Republic: Praha area); Kowalczyk, 1990:291 (Poland: Góry Świętokrzyskie); Skibińska, 1995:424, 425 (Poland: pine forest in Białowieża Primeval Forest); Pawlikowski and Kruszyński, 1996:86 (Poland: Toruń Basin); Pagliano, 2014:118 (specimens in M. Spinola collection, Torino, are *Alysson costai*, *Alysson spinosus*, and *Alysson tricolor*).

As *Alysson pertheesi*: Nedelkov, 1914:197 (Bulgaria), corrected to *Alysson spinosus* by Ljubomirov, 1999a:22.

As *Alysson ratzeburgi*: Marquet, 1875:204 (France: Hérault: Montpellier), corrected to *Alysson bimaculatus* by Marquet, 1881:181; Dittrich, 1911b:24 (Germany: Niederschlesien: Mirkau, now Poland: Dolny Śląsk: Mirków near Wrocław), corrected to *Alysson fuscatus* by Noskiewicz and Chudoba, 1949:409; Nedelkov, 1914:197 (Bulgaria), corrected to *Alysson spinosus* by Ljubomirov, 1999a:22.

As *Alysson tricolor*: Dittrich, 1911b:23 (Germany: Niederschlesien: Mirkau, now Poland: Dolny Śląsk: Mirków near Wrocław), corrected to *Alysson fuscatus* by Noskiewicz and Chudoba, 1949:409.

33. *striatus* W. Fox

Alysson striatus W. Fox, 1894b:88, ♂. Lectotype: ♂, USA: District of Columbia (ANSP), designated by Cresson, 1928:48. – Handlirsch, 1895:834 (original description translated into German); Dalla Torre, 1897:565 (in catalog of world Sphecidae); Ashmead, 1899d:329 (in checklist of North American Crabronidae); Bradley, 1928:1015 (in catalog of New York Sphecidae); Krombein in Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1979b:1685 (in catalog of North American Hymenoptera).

34. *taiwanus* Sonan

Alysson taiwana Sonan, 1940:14, ♀, incorrect original termination. Holotype: ♀, Taiwan: Shinchiku (depository?) – Tsuneki, 1968h:5 (in key to *Alysson* of South and East Asia); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Wu and Zhou, 1987:215 (in key to *Alysson* of China); Porter, Stange, and Wang, 1999:13 (in checklist of Sphecidae of Taiwan); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Zou, Ma, Chen, and Li, 2025:586 (in key to *Alysson* of China).

35. *takasago* Tsuneki

Alysson takasago Tsuneki, 1977j:266, ♂. Holotype: ♂, Taiwan: Fuhosho (TMB). – Hua, 2006:277 (in list of Chinese insects, geographic distribution); Zou, Ma, Chen, and Li, 2025:586 (in key to *Alysson* of China).

36. *testaceitarsis* Cameron

Alysson testaceitarsis Cameron, 1902c:290, ♀. Holotype or syntypes: ♀, India: Himachal Pradesh: Simla (BMNH). – Tsuneki, 1968h:5 (in key to *Alysson* of South and East Asia); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Girish Kumar, Sheela, and Sharma, 2020:271 (in key to Indian *Alysson*).

37. *tomentosus* McLeay

Alysson tomentosus McLeay, 1828:457, sex not stated. Holotype or syntypes: Australia: no specific locality (depository unknown). – Not listed by Bohart and Menke, 1976; Cardale, 1985:303 (in catalog of Australian Sphecidae).

38. *triangularis* Krombein

Alysson triangularis Krombein, 1985a:13, ♀, ♂. Holotype: ♀, Sri Lanka: Central Province: Kandy District: Kandy - Udawattakele Sanctuary (USNM). – Evans and O'Neil, 2007:27 (summary of information on nesting habits); Nemkov, 2012b:119 (review of known prey); Girish Kumar, Sheela, and Sharma, 2020:271 (in key to Indian *Alysson*), 276 (color illustrations), 278 (first record from India: Tamil Nadu).

39. *triangulifer* Provancher

Alysson triangulifer Provancher, 1887:272, ... Lectotype: ♂, Canada: ... (Laval Univ.), designated by Gahan and Rohwer, 1917:305. – W. Fox, 1894b:88 (in revision of North American *Alysson*, description of ♀, as *trianguliferus*); Handlirsch, 1895:835 (original description translated into German); Dalla Torre, 1897:565 (in catalog of world Sphecidae); Ashmead, 1899d:329 (in checklist of North American Crabronidae); Harrington, 1902:223 (Canada: Ontario: Ottawa); H. Smith, 1908b:341 (in revision of Nebraskan Sphecidae); Mickel, 1918b:347 (in catalog of Nebraskan Sphecidae); Washburn, 1919:216 (in list of Hymenoptera of Minnesota, as *trianguliferum*); J.Ch. Bradley, 1920:131 (New York, as *trianguliferus*), 1928:1015 (in catalog of New York Sphecidae); Hendrickson, 1930:158 (Iowa); Brimley, 1938:446 (North Carolina: Hendersonville); Krombein in Muesebeck, Krombein, and Townes, 1951:981 (in catalog of North American Hymenoptera); Krombein, 1963f:276 (Maryland: Plummers Island near Washington, D.C., as *triangulifer triangulifer*); Steiner, 1973:27 (Canada: Northwest Territories); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1979b:1685 (in catalog of North American Hymenoptera); Finnamore, 1982:206 (in Sphecid Fauna of southern Quebec); Ahlstrom, 1995:112 (in checklist of insects of North Carolina); Finnamore, 1997:892 (Canada: Yukon, as *triangulifer triangulifer*); Buck, 2004:27 (Canada: in checklist of Crabronidae of Ontario); Ratzlaff, 2016:23 (Canada: British Columbia, after Bradley, 1920 and Krombein, 1979); Buck and Bennett, 2025:207 (in checklist of Sphecidae *sensu lato* of Canada, Alaska, and Greenland: USA: Alaska, Canada: British Columbia, Northwest Territories, Ontario, Quebec, Yukon Territory, first records from Alberta, Manitoba, New Brunswick, Newfoundland Island, and Saskatchewan: no specific locality).

ssp. *shawi* Bradley

Alysson shawi Bradley, 1920:130, ♀, ♂. Holotype: ♂, Canada: British Columbia: Carbonate on Columbia River (Cornell Univ.). – As *Alysson triangulifer shawi*: Krombein in Muesebeck, Krombein, and Townes, 1951:981 (new sta-

tus, in catalog of North American Hymenoptera); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Krombein, 1979b:1685 (in catalog of North American Hymenoptera).

40. *tricolor* Lepeletier de Saint Fargeau and Audinet-Serville

Alysson tricolor Lepeletier de Saint Fargeau and Audinet-Serville, 1825:50, ♀. Syntypes: ♀, France: Normandie: Luc, and Paris area (lost). – Vander Linden, 1829:92 (bibliographic references, Belgium: Bruxelles); Lepeletier de Saint Fargeau, 1845:89 (in revision of world Hymenoptera); Wesmael, 1852:261 (Belgium); F. Smith, 1856:373 (in catalog of Hymenoptera in British Museum); A. Costa, 1864b:98 (two specimens from France in Napoli Museum), 1867c:151 (in revision of Italian Sphecidae); Kirchner, 1867:19 (in catalog of European Hymenoptera); A. Costa, 1871b:82 (in revision of Italian Sphecidae); Dours, 1874:142 (in catalog of Hymenoptera of France); Wüstnei, 1886:9 (known from Germany, but not from Nassau); Handlirsch, 1888a:244 (in revision of world Bembicinae); Kohl, 1893a:28 (Austria: Niederösterreich: Wien); Handlirsch, 1895:831 (Switzerland: Grono); Acloque, 1897:85 (in Sphecid Fauna of France and Algeria); Dalla Torre, 1897:565 (in catalog of world Sphecidae); Mocsáry, 1897:81 (Hungary: Budapest); Adlerz, 1904:138 (known prey: cicadellids); W. Schulz, 1906:201 (correct authorship is Lepeletier de Saint Fargeau and Serville); Schmiedeknecht, 1907:224, 225 (in key to Hymenoptera of Central Europe); Ferton, 1908:558 (prey); de Gaulle, 1908:120 (in catalog of French Hymenoptera); nec Dittrich, 1911b:23 (= *Alysson spinosus*); Noskiewicz, 1918:137 (Galicia, now western Ukraine: Hołosko near Lwów); Grubod, 1919:857 (Italy: Piemonte: Susa); Steck, 1920:77 (Switzerland: Grono, Novaggio); Dubois, 1922:204, 205 (in key to Belgian Sphecidae); Berland, 1925d:96 (in Sphecid Fauna of France); Bouwman, 1927:93, 94 (in key to Sphecidae of Netherlands, not yet found in Netherlands); Schmiedeknecht, 1930:686, 687 (in keys to Hymenoptera of North and Central Europe); Bondroit, 1932:32 (Belgium: recorded from Bruxelles area by Wesmael, 1852); Dusmet y Alonso, 1935:53 (Spain: Madrid: El Pardo); Vergne, 1935:118 (France: Auvergne: Lezoux); Crèvecoeur and Maréchal, 1936:253 (in catalog of Belgian Hymenoptera); Móczár, 1939:5 (Hungary: Jászberény); Zavadil, 1939:120 (Czech Republic: Moravia: Kobyly); Giner Marí, 1943a:152 (in Sphecid Fauna of Spain); Šnoflák in Zavadil and Šnoflák, 1948:78, 79 (in key to Sphecidae of Czechoslovakia); Vogrin, 1955:36 (Yugoslavia); Ceballos, 1956:379 (in catalog of Hymenoptera of Spain); Bajári, 1957a:94, 95 (in key to Hungarian Sphecidae); Noskiewicz and Puławski, 1960:83, 84 (in key to Polish Sphecidae, not yet found in Poland); de Beaumont, 1964c:63 (in Sphecid Fauna of Switzerland), 1967a:326 (Turkey); Leclercq, 1968:map 1169 (Belgium: distribution); Scobiola-Palade, 1968c:384 (Romania: Pusnicu); Stritt, 1968:110 (Germany: Baden-Württemberg), 1969:131 (first record from Germany: Baden-Württemberg: Karlsruhe), 1971a:104 (first record from Germany: Baden-Württemberg: Karlsruhe), 1971b:155 (first record from Germany: Karlsruhe area); Balthasar, 1972:351 (in Sphecid Fauna of Czechoslovakia); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Kazenas, 1978b:94 (in key to Sphecidae of Kazakhstan and Central Asia); Leclercq, 1978b:map 1169 (distribution in Belgium); Puławski, 1978:212 (in key to Sphecidae of European part of USSR); Schmidt, 1979b:332 (Germany: Baden-Württemberg); Casolari and Casolari Moreno, 1980:125 (specimens in M. Spinola collection, Torino); Dollfuss, 1983a:75 (endangered in Niederösterreich, Austria), 1983b:20 (in catalog of Sphecidae of Austria: Wien; last found in 1866); Westrich and Schmidt, 1985b:110 (Germany: endangered in Baden-Württemberg); Pádr in Šedivý, 1989a:167 (in checklist of Czechoslovakian Sphecidae); Pagliano, 1990:67 (in catalog of Italian Sphecidae); Day, 1991:xii (in summary of European Endangered Hymenoptera Lists); Dollfuss, 1991:182 (in key to Sphecidae of North and Central Europe); Pagliano, 1992:31 (in key to Italian Nyssoninae, now Bembicinae); Dollfuss, 1994:97 (endangered in Austria); Negrisolo in Minelli, Ruffo, and La Posta, 1995b:3 (in catalog of Italian fauna); Gusenleitner, 1996a:8 (Austria: Burgenland); Schmid-Egger, Schmidt, and Doczkal, 1996:376 (Germany: Baden-Württemberg: endangered); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:128 (in Sphecid Fauna of Western Europe); Pagliano and Alma, 1997:179 (known prey listed); Schmidt and Schmid-Egger, 1997:5 (in checklist of German Sphecidae); Dollfuss, Gusenleitner, and Bregant, 1998:521 (Austria: Burgenland: Schützgraben east of Minihof-Liebau); Gorobchishin, 1998a:50 (Ukraine: Kiev and surroundings); Ljubomirov, 1999a:22 (Bulgaria),

1999b:56 (Bulgaria: Mount Vitosha near Sofia); Pauly, 1999:111 (in catalog of Belgian aculeates); Blösch, 2000:360 (in Sphecidae Fauna of Germany); Gayubo, González, and Torres, 2000:187 (Spain: Salamanca Province); Józán, 2001:280 (Hungary: Somogy County); Kazenas, 2001b:43 (tentatively in checklist of Sphecidae of Kazakhstan and Central Asia); Ohl et al., 2001:137 (distribution in Germany by provinces); Zettel, Gross, and Mazzucco, 2001:79 (Austria: Wien, not collected in XXth century); Gorobchishin, 2003:48 (Ukraine: Kiev: botanical garden); Mandery, Kraus, Voith, Wickl, Scheuchl, Schubert, and Warncke, 2003:63 (Germany: in catalog of wasps and bees of Bayern); Deván, 2004:146 (western Slovakia); Pagliano and Negrisolo, 2005:113 (in Sphecidae Fauna of Italy); Straka, 2005a:393 (extinct in Czech Republic); Gorobchishin, 2006:135 (Ukraine: Kiev); Jacobs, 2007:159, 160 (in key to Crabronidae of Germany); Józán, 2007:171, 180 (Hungary: Zselic hills); Vepřek and Straka, 2007:207 (in catalog of Sphecidae of Czech Republic and Slovakia, known from Moravia and Slovakia only); Ljubomirov and Yildirim, 2008:63 (in catalog of Crabronidae of Turkey); Baños-Picón, Asís, Gayubo, and Tormos, 2009:310 (Spain: frequency of specimens collected with hand nets and Malaise traps); González, Gayubo, Asís, and Tormos, 2009:625 (Spain: Salamanca and Zamora provinces: Arribes del Duero Natural Park); Pagliano, 2009:177 (Italy: Lazio: Morlupo); Bitsch, 2010:108 (in supplement to vol. II of Faune de France, 1997: review of recent records from Europe); Schmid-Egger, 2010a:13 (in red list of Aculeata of Germany: very rare); Shorenko and Kononov, 2010:13 (Ukraine: Khar'kov Oblast': Gaidary; Donetsk Oblast': Thorez); van der Smissen, 2010b:383 (France: Gard: Collias; Vaucluse: Le Beaucet-Venasque); Gogala, 2011:26 (Slovenia: Ljubljana); Józán, 2011:183 (in checklist of Sphecidae *sensu lato* of Hungary); Tischendorf, Frommer, Flügel et al., 2011:85 (Germany: Hessen: insufficient data to evaluate degree of endangerment); Strumia, Pagliano, and Gayubo, 2012:56 (Italy: Toscana: Riserva di San Rossore); Reder, 2013a:25 (Germany: Rheinland-Pfalz: three localities, review of German collecting localities with distribution map in Germany); Bitsch, 2014:398 (record from France by van der Smissen, 2010b reported); Pagliano, 2014:118 (specimens in M. Spinola collection, Torino, are *Alysson tricolor*); Yildirim, Ljubomirov, and Lelej, 2014:5 (distribution in Turkey by biogeographic provinces); Reder and Niehuis, 2014:6 (Germany: Birkenheide, Flörsheim-Dalsheim, and Worms); Koçak and Kemal, 2015:136 (in checklist of Hymenoptera of Turkey); Mokrousov, Shlyakhtenok, and Grebennikov, 2016:20 (Russia: Belgorod Oblast': Les na Vorskle Nature Reserve); Jansen, 2017:6 (Germany: Sachsen: Leipzig), 7 (color photographs); Nemkov, 2017a:Table p. 1 (in zoogeographic analysis of Palearctic Bembicinae), 2017b:219 (in catalog of Bembicinae of Russia); Gayubo, Bitsch, Barbier, and Jacobs in Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:196 (in Sphecidae Fauna of Europe); Bertsch, 2021:46 (Germany: Bayern: Bogen at 48.898°N 12.710°E and Straubing at 48.874°N 12.559°E; prey: four genera of Cicadellidae); Kaplan and Yildirim, 2021:23 (in catalog of Crabronidae of Turkey); Witt, 2021:50 (Germany: Niedersachsen: Hannover-Wülfel); Bertsch, 2023:36 (Germany: Bayern, Hessen, and Saarland); Kaplan and Yildirim, 2023:1679 (in checklist of Sphecidae *sensu lato* of Turkey).

41. *tridentatus* Wu and Zhou

Alysson tridentatus Wu and Zhou, 1987:218 and 221, ♀. Holotype: ♀, China: Jiangsi: Guli (Inst. Zool. Beijing). – Wu and Zhou, 1987:215 (in key to *Alysson* of China), 1996:133 (in revision in Economic Insect Fauna of China); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Zou, Ma, Chen, and Li, 2025:587 (in key to *Alysson* of China).

42. *verhoeffi* Tsuneki

Alysson verhoeffi Tsuneki, 1967e:8, ♀. Holotype: ♀, China: Heilongjiang: Harbin (RMNH). – Tsuneki, 1968h:6 (in key to *Alysson* of South and East Asia); R. Bohart and Menke, 1976:458 (in checklist of world Sphecidae); Wu and Zhou, 1987:215 (in key to *Alysson* of China); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Nemkov, 2014a:277 (in key to *Alysson* of Russia and adjacent countries); Mokrousov and Danilov, 2017:57 (China: Liaoning 50 km north of Shenyang = Mukden), Russia: Khabarovskiy Kray: Khabarovsk; Nemkov, 2017b:219 (in catalog of Bembicinae of Russia); Zou, Ma, Chen, and Li, 2025:586 (in key to *Alysson* of China).

43. *yunnanensis* Wu and Zhou

Alysson yunnanensis Wu and Zhou, 1987:215 and 221, ♀. Holotype: ♀, China: Yunnan: Xishuangbanna: Menglun (Inst. Zool. Beijing). – Wu and Zhou, 1987:214 (in key to *Alysson* of China), 1996:131 (in revision in Economic Insect Fauna of China); Hua, 2006:277 (in list of Chinese insects, geographic distribution); Zou, Ma, Chen, and Li, 2025:586 (in key to *Alysson* of China).

NOMEN NUDUM IN ALYSSON

Alysson triangularis Fox: Ashmead, 1899d:329 (in checklist of North American Crabronidae).

SPECIES DESCRIBED IN ALYSSON AND TRANSFERRED TO OTHER GENERA

aculeatus Cresson, 1865 = *Didineis*

ater Spinola, 1808 = *Pison*

barbieri de Beaumont, 1968 = *Didineis*

latro de Beaumont, 1967 = *Didineis*

nipponicus Tsuneki, 1968 = *Didineis*

texanus Cresson, 1872 = *Didineis*