

Last updated: 26 August 2024

BELOMICROIDES

Belomicroides Kohl, 1899:312. Type species: *Belomicroides schmiedeknechti* Kohl, 1899, by monotypy.

Key: Gayubo, 1983b:44.

Revision: Antropov, 1994 (Asian species), 2002 (world).

1. alessandroi Antropov

Belomicroides alessandroi Antropov, 2002a:565. Holotype: ♂, Egypt: Gebel Asfar near Cairo (Torino). – Antropov, 2002b:696 (in revision of world *Belomicroides*).

2. arenarius Arnold

Belomicroides arenarius Arnold, 1960b:476, ♂. Holotype: ♂, Zimbabwe: Sawmills (SAM). – Bohart and Menke, 1976:360 (in checklist of world Sphecidae). – Antropov, 2002b:696 (in revision of world *Belomicroides*, description of ♀).

3. bytinskii Antropov

Belomicroides bytinskii Antropov, 2002a:566, ♂. Holotype: ♂, Egypt: Sinai: Oasis Feiran (Tel Aviv. Univ.). – Antropov, 2002b:698 (in revision of world *Belomicroides*).

4. clypeatus Antropov

Belomicroides clypeatus Antropov, 2002a:568, ♂. Holotype: ♂, South Africa: Western Cape Province: northeast of Touwsrivier (SAM). – Antropov, 2002b:698 (in revision of world *Belomicroides*).

5. erythrogaster Antropov

Belomicroides erythrogaster Antropov, 2002a:569, ♀. Holotype: ♀, Namibia: Gobabis District: Gobabis (BMNH). – Antropov, 2002b:698 (in revision of world *Belomicroides*).

6. krombeini Antropov

Belomicroides krombeini Antropov, 2002a:571, ♂. Holotype: ♂, Egypt: Suez Province: Ain Sokhna (USNM). – Antropov, 2002b:699 (in revision of world *Belomicroides*).

7. marleyi Arnold

Belomicroides marleyi Arnold, 1927:58, ♀ (as *Marleyi*, incorrect original capitalization). Holotype: ♀, South Africa: KwaZulu-Natal: Durban (Durban Mus.). – Arnold, 1930:14 (in checklist of Afrotropical Sphecidae); Pate, 1931a:113 (in key to *Belomicroides*); R. Bohart and Menke, 1976:360 (in checklist of world Sphecidae); Antropov, 2002b:699 (in revision of world *Belomicroides*).

8. maurusius Pate

Belomicroides maurusius Pate, 1931a:111, ♂. Holotype: ♂, Algeria: Biskra (Cornell Univ.). – Pate, 1931a:113 (in key to *Belomicroides*); R. Bohart and Menke, 1976:360 (in checklist of world Sphecidae); Leclercq, 1993c:43 (in catalog of western Palearctic Crabronini and Oxybelini); Antropov, 2002b:700 (in revision of world *Belomicroides*), 2011a:626 (United Arab Emirates: Um al-Quwain at 25°32'N 55°32'E); Gadallah, 2020b:33 (in list of aculeate wasps of Arabian Peninsula).

9. maximiliani Antropov

Belomicroides maximiliani Antropov, 2002:572, ♀, ♂. Holotype: ♀, Namibia: Karibib District: 65 km southwest of Usakos (M. Schwarz coll., Ansfelden, Austria). – Antropov, 2002b:702 (in revision of world *Belomicroides*).

10. melas Antropov

Belomicroides melas Antropov, 1994:94, ♂. Holotype: ♂, Uzbekistan: 35 km southeast of Bukhara (ZMMU. – Medetov, Embergenov, Kholmatov, Elmurodova, Rakhimov, and Tajibaeva, 2024:424 (Uzbekistan: distribution).

11. pictus Arnold

Belomicroides pictus Arnold, 1927:56, ♀, ♂. Lectotype: ♂: South Africa: Eastern Cape Province: Willowmore (TMP), designated by Antropov, 200b:702. – Arnold, 1930:14 (in checklist of Afrotropical Sphecidae); Pate, 1931a:113 (in key to *Belomicroides*); R. Bohart and Menke, 1976:360 (in checklist of world Sphecidae); Antropov, 2002b:702 (in revision of world *Belomicroides*).

ssp. chrysoptilus Antropov

Belomicroides pictus chrysoptilus Antropov, 2002a:574, ♀. Holotype: ♀, Namibia: Rundu District: Rundu (M. Schwarz coll.). – Antropov, 2002b:705 (in revision of world *Belomicroides*).

12. politus Antropov

Belomicroides politus Antropov, 2002:576, ♀, ♂. Holotype: ♀, Namibia: Diamond Area No.1: Klinghardtberge: Spitzkuppe (PPRI). – Antropov, 2002b:705 (in revision of world *Belomicroides*).

13. rift Antropov

Belomicroides rift Antropov, 2002a:578, ♀, ♂. Holotype: ♀, Kenya: Rift Valley Province: Marich Pass at 1°32.2'N 35°27.4'E (CAS). – Antropov, 2002b:705 (in revision of world *Belomicroides*).

14. rufus Antropov

Belomicroides rufus Antropov, 2007c:634, ♀. Holotype: ♀, Morocco: Mhamid 80 km south of Zagora in Drâa Valley (ZMHU). Paratype: Egypt: Wadi Digla near Cairo.

15. schmiedeknehti Kohl

Belomicroides schmiedeknehti Kohl, 1899:313, ♀ (as *Schmiedeknehti*, incorrect original capitalization). Holotype: ♀, Tunisia: Tunis (NHMW). – Pate, 1931a:113 (in key to *Belomicroides*); R. Bohart and Menke, 1976:360 (in checklist of world Sphecidae); Dollfuss, 1989:7 (type material in NHMV); Leclercq, 1993c:43 (in catalog of western Palearctic Crabronini and Oxybelini); Antropov, 2002b:706 (in revision of world *Belomicroides*); Ben Khedher, Yildirim, and Braham, 2020a:104 (Tunisia: Mahdia Province: Sidi Alouane: Aouled Kloula at 35°22'09.4"N 10°51'22.6"E). – As *Belomicrus schmiedeknehti*: Kohl, 1923:117 (new combination, in key to world *Belomicrus*), 180 (in revision of world *Belomicrus*).

16. veronicae Antropov

Belomicroides veronicae Antropov, 2002a:580, ♀. Holotype: ♀, Ethiopia: Sidamo Province: 4 km east of Yabelo at 4°53'N 38°08'E (CAS). – Antropov, 2002b:707 (in revision of world *Belomicroides*).

17. woji Antropov

Belomicroides woji Antropov, 2002a:582, ♂. Holotype: Namibia: Okahandja District: 17 km west of Okahandja (CAS). – Antropov, 2002b:707 (in revision of world *Belomicroides*).

sp.

de Beaumont, 1958b:70 (Algeria: Tassili des Ajjer); S. Gess, 1996:301 (South Africa: floral records).

**SPECIES DESCRIBED IN *BELOMICROIDES* AND TRANSFERRED TO
*PSEUDOMICROIDES***

desertus Antropov, 1994 = *Pseudomicroides*

elvirae Gayubo, 1983 = *Pseudomicroides*

fergusoni de Beaumont, 1960 = *Pseudomicroides*

melas Antropov, 1994 = *Pseudomicroides*

olgae Kazenas, 1979= *Pseudomicroides*

rostratus Krombein, 1982 = *Pseudomicroides*