

Last updated: 18 November, 2020

HEMIDULA

Hemidula Burmeister, 1874:119. Type species: *Monedula singularis* Taschenberg, 1870, designated by J. Parker, 1917:2. – Pate, 1943c:2 (omitted from Pate's, 1937 Catalog of Generic Names).

1. *burmeisteri* Willink

Hemidula burmeisteri Willink, 1947:541, ♀. Holotype: ♀, Argentina: Santiago del Estero: no specific locality (MACN). – Willink, 1958:52 (description of ♂); R. Bohart and Menke, 1976:540 (in checklist of world Sphecidae); Genise, 1989b:275 (nest and prey); Amarante, 2002:10 (in catalog of Neotropical Crabronidae); Evans and O'Neil, 2007:131 (summary of information on nesting habits).

2. *singularis* (Taschenberg)

Monedula singularis Taschenberg, 1870:25, ♀. Lectotype: ♀, Argentina: Mendoza: no specific locality (Halle), designated by M.A. Mendoza, 2012:220. – Burmeister, 1874:121 (in revision of Bembicinae of Argentina); Handlirsch, 1890:125 (in revision of world Bembicinae); Dalla Torre, 1897:500 (in catalog of world Sphecidae); Schrottky, 1903b:177 (in checklist of Hymenoptera of Argentina, Paraguay, and Uruguay). – *As Stictia singularis*: Jörgensen, 1912:289 (new combination, Argentina: Mendoza Province); Schrottky, 1913a:232 (Argentina: Mendoza). – *As Hemidula singularis*: J. Parker, 1929:3 (new combination, in revision of Stizini and Bembicini); Willink, 1947:528 (in revision of Bembicini of Argentina); Lohrmann, 1948:462 (in review of *Hemidula*); Willink, 1952d:76 (Argentina: Chubut); R. Bohart and Menke, 1976:540 (in checklist of world Sphecidae); Genise, 1982e:77 (nesting behavior); Amarante, 2002:10 (in catalog of Neotropical Crabronidae); Evans and O'Neil, 2007:132 (summary of information on nesting habits); M.A. Mendoza, 2012:220 (redescription, locality records from Argentina, many color photographs).