

Last updated: 5 December 2020

KOHLIELLA

Kohliella Brauns, 1910a:668. Type species: *Kohliella alaris* Brauns, 1910, by monotypy.

Revision: Pulawski, 1991b.

1. *anula* Pulawski

Kohliella anula Pulawski, 1991b:295, ♀, ♂. Holotype: ♀, Sri Lanka: Mannar District: 0.8 km northeast of Kokmotte in Wilpattu National Park at about 8°32'N 80°02'E (USNM). – Mawadda, Girish Kummar, and Sureshan, 2020:148 (first record from India: Kerala: Kannur District: Kannapuram at 11°58'45.768"N, 75°18'25.884"E).

2. *alaris* Brauns

Kohliella alaris Brauns, 1910a:669, ♂. Holotype: ♂, South Africa: Eastern Cape Province: Willowmore (TMP). – Brauns, 1911a:239 (South Africa); Arnold, 1922:135 (in revision of South African Larrini), 1924:43 (description of ♀; South Africa: Western Cape Province: Hex River; Zimbabwe: Bulawayo), 1930:2 (in checklist of Afrotropical Sphecidae); de Beaumont, 1967b:510 (South Africa: Western Cape Province: Cape Peninsula and Table Mountain); R. Bohart and Menke, 1976:286 (in checklist of world Sphecidae); F. Gess and S. Gess, 1980a:45 (nesting habits); F. Gess, 1981:20 (South Africa: 18 km west-northwest of Grahamstown; nesting in friable soils), 40 (use of foretarsal rake), 47 (multiple prey, size of individual prey determines oviposition); Pulawski, 1991b:297 (in revision of *Kohliella*).

3. *stevensoni* Arnold

Kohliella stevensoni Arnold, 1924:42, ♂ (as *Stevensoni*, incorrect original capitalization). Lectotype: ♂, Zimbabwe: Sawmills (SAM), designated by Pulawski, 1991b:298. – Arnold, 1930:2 (in checklist of Afrotropical Sphecidae); R. Bohart and Menke, 1976:286 (in checklist of world Sphecidae); Pulawski, 1991b:298 (in revision of *Kohliella*).