

Last updated: 20 April 2025

PAE

Pae Pate, 1944b:364. Type species: *Pae paniquita* Pate, 1944, by original designation.

Lamocrabro Leclercq, 1951a:44 and 48. Type species: *Crabro nasicornis* F. Smith, 1873, by original designation.

Synonymized with *Pae* by Leclercq, 1995:83.

Revision: Leclercq, 1995a.

1. amaripa Pate

Pae amaripa Pate, 1944:369, ♀. Holotype: ♀, Guyana: Tukeit Trail along Potaro River at Kaietur Falls (BMNH). – Leclercq, 1951a:49 (Ecuador: Napo), 1954c:214 (in catalog of world Crabronini, Mexico); R. Bohart and Menke, 1976:413 (in checklist of world Sphecidae); Leclercq, 1995a:86 (faunal records, discussion of characters); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Starr and Hook, 2003:19 (in catalog of Aculeata of Trinidad, West Indies); Leclercq, 2005b:7 (locality records from Brazil, Ecuador, French Guyana, and Peru, those from Panama refer to *Pae paniquita*), 8 (comparison with *Pae manausae*), 2005e:19 (records from Panama in Leclercq, 2005b:7 refer to *Pae paniquita*); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

2. beniae Leclercq

Pae beniae Leclercq, 1995a:86, ♀. Holotype: ♀, Bolivia: Beni: Rurebanaque (BMNH). – Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

3. macasae Leclercq

Pae macasae Leclercq, 1995a:87, ♀. Holotype: ♀, Ecuador: Moronga-Santiago: 6 km of Macas (BMNH). – Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

4. manausae Leclercq

Pae manausae Leclercq, 2005b:7, ♂. Holotype: ♂, Brazil: Amazonas: 5 mi. southeast of Manaus (CNC).

5. napoensis Leclercq

Pae napoensis Leclercq, 1995a:88, ♀. Holotype: ♀, Ecuador: Napo: Panacocha (BMNH). Paratype: Colombia. – Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

6. nasicornis (F. Smith)

Crabro nasicornis F. Smith, 1873b:102, ♀. Syntypes: Brazil: Amazonas: Ega, now Tefé, and São Paulo de Olivença (BMNH). – Dalla Torre, 1897:611 (in catalog of world Sphecidae). – **As *Lamocrabro nasicornis***: Leclercq, 1951a:41 (new generic assignment), 1954c:214 (in catalog of world Crabronini); R. Bohart and Menke, 1976:413 (in checklist of world Sphecidae). – **As *Pae nasicornis***: Leclercq, 1995a:88 (new generic assignment, discussion of characters, locality records from Colombia, Guyana, Ecuador, Peru, Bolivia, Argentina, Brazil); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

Crabro dentatus F. Smith, 1873b:104, ♀. Holotype or syntypes: ♀, Brazil: Pará: Pará, now Belém (BMNH). Synonymized with *Lamocrabro nasicornis* by Leclercq in Bohart and Menke, 1976:413. – Dalla Torre, 1897:596 (in catalog of world Sphecidae, as *dentalis*). – **As *Lamocrabro dentatus***: Leclercq, 1951a:41 (new generic assignment), 1954c:214 (in catalog of world Crabronini).

7. paniquita Pate

Pae paniquita Pate, 1944:366. Holotype: ♀, Colombia: Boyaca Department: Muzo (MCZ). – Leclercq, 1954c:214 (in catalog of world Crabronini); R. Bohart and Menke, 1976:413 (in checklist of world Sphecidae); Leclercq, 1995a:89 (diagnostic characters; Costa Rica, Panama, Colombia); Hanson and Menke, 1995:644 (known from Costa Rica); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Ruíz Cancino, Coronado Blanco, Varela

Fuente, and Horta Vega, 2002:664 (in checklist of Mexican Sphecidae); Leclercq, 2005b:8 (Mexico: Vera Cruz: Lake Catemaco; difference with male of *Pae paniquita*), 2005e:19 (Panama: National Park Darién: Rancho Frío).

8. *surinamensis* Leclercq

Pae surinamensis Leclercq, 1995a:90, ♀. Holotype: ♀, Surinam: no specific locality (FSAG). – Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Leclercq, 2007g:194 (holotype in FSAG).