

Last updated: 10 May 2025

PRIONYX

Prionyx Vander Linden, 1827:362. Type species: *Ammophila kirbii* Vander Linden, 1827, by monotypy.

Priononyx Dahlbom, 1843:28. Type species: *Pepsis thomae* (Fabricius, 1804) [= *Sphex thomae* Fabricius, 1775], by monotypy.

Enodia Dahlbom, 1843:28, junior homonym of *Enodia* Hübner, 1819 (Lepidoptera). Type species: *Sphex albisectus* Lepeletier de Saint Fargeau and Serville, 1828 [= *Ammophila kirbii* Vander Linden, 1827], designated by Kohl, 1885b:164. Synonymized with *Prionyx* Vander Linden by Pate, 1935:250.

Harpactopus F. Smith, 1856:264. Type species: *Harpactopus crudelis* F. Smith, designated by Patton, 1880a:384.

Paraspheg F. Smith, 1856:267. Type species: *Sphex albisectus* Lepeletier de Saint Fargeau and Serville, 1828 [= *Ammophila kirbii* Vander Linden, 1827], designated by Kohl, 1885b:164. Synonymized with *Prionyx* Vander Linden by Pate, 1935:250.

Gastrosphaeria A. Costa, 1858b:10. Type species: *Gastrosphaeria anthracina* A. Costa, 1858 [= *Sphex subfuscatus* Dahlbom, 1845], by monotypy.

Pseudosphex Taschenberg, 1869:420, junior homonym of *Pseudosphex* Hübner, 1818 (Lepidoptera). Type species: *Pseudosphex pumilio* Taschenberg, 1869, by monotypy.

Calosphex Kohl, 1890b:113. Type species: *Sphex niveatus* Dufour, 1853, designated by Pate, 1937c:15. – As *Callosphex*: Rohwer, 1913:450 (misspelling).

Neosphex Reed, 1894:627. Type species: *Neosphex albospiniferus* Reed, 1894 [= *Pseudosphex pumilio* Taschenberg, 1869], by monotypy.

Foxiprionyx Danilov in Danilov and Odintsev, 2023:432. Type species: *R. Bohart* and Menke, 1963 by original designation and monotypy.

Key to species: F. Parker, 1960 (North America); Mingo and Gayubo, 1983:152 (Spain); Pagliano, 1984:366 (Italy); Guichard, 1988a:119 (Arabian Peninsula); Danilov, 2012a, b (Russia and adjacent countries); Can and Gülmez, 2019:348 (Turkey); Danilov, 2014b:514 (Siberia).

Classification: de Beaumont, 1968b:148 (Mediterranean species).

Review of biology: Kazenas: 2001b:81.

Recognition of subgenera: Danilov, 2019; Danilov and Odintsev, 2023 (key to subgenera).

1. afghaniensis (de Beaumont)

Sphex afghaniensis de Beaumont, 1970a:391, ♀. Holotype: ♀, Afghanistan: Kabul (Brno Mus.). – As *Prionyx afghaniensis*: R. Bohart and Menke, 1976:131 (new generic assignment, in checklist of world Sphecidae); Dunford, Turbyville, and Leavengood, 2014:11 (listed as medically important in Afghanistan); Danilov and Odintsev, 2023:432 (in subgenus *Calosphex*).

2. atratus (Lepeletier de Saint Fargeau)

Sphex atratus Lepeletier de Saint Fargeau, 1845:355, ♀. Holotype or syntypes: ♀, origin unknown (originally J. Serville coll., now M. Spinola collection, Torino). – Cresson, 1863:319 (in catalog of North American Hymenoptera), 1868:379 (New Mexico), 1876:208 (Colorado: Cannon City); Kohl, 1890b:357 (in revision of world Sphecini); Dalla Torre, 1897:415 (in catalog of world Hymenoptera); Strand, 1916:100 (diagnostic characters); Berland, 1926c:202 (specimens in MNHN); Murray in Muesebeck, Krombein, and Townes, 1951:973 (in catalog of North American Hymenoptera); Casolari and Casolari Moreno, 1980:102 (specimens in M. Spinola collection, Torino); Pagliano, 2008:527 (unpublished lectotype in Spinola collection, Torino). – As *Priononyx atratus*: F. Smith, 1856:266 (new generic assignment, in catalog of Hymenoptera in British Museum); Cresson, 1865b:464 (specimens in ANSP collection), 1868:379 (New Mexico), 1873:213 (Texas), 1875:715 (Colorado, New Mexico), 1876:208 (Colorado: Canon City); Riley, 1878:318 (Colorado, preying on

pupae of *Melanoplus spretus* (Walsh), an acridid); Patton, 1879d:354 (northwestern Kansas); Snow, 1881:96 (in checklist of Hymenoptera of Kansas: Douglas County: no specific locality); Cresson, 1887:276 (in catalog of North American Hymenoptera); Ashmead, 1890:33 (in checklist of Hymenoptera of Colorado, as *atrata*); C. Robertson, 1892:107 (visiting flowers of *Pycnanthemum linifolium* Ph., Lamiaceae, as *atrata*), 114 (visiting flowers of *Blephilia ciliata* Raf., Lamiaceae, as *atrata*), 1894:456 (visiting flowers of *Solidago canadensis* Linnaeus, Asteraceae, as *atrata*), 458 (visiting flowers of *Solidago lanceolata* Linnaeus, Asteraceae, as *atrata*), 469 (visiting flowers of *Lepachys pinnata* Torr. and Gray, Asteraceae), 470 (visiting flowers of *Helianthus grosse-serratus* Martens, Asteraceae), 475 (visiting flowers of *Helenium autumnale* Linnaeus, Asteraceae), 1896:72 (visiting flowers of *Polygonum pennsylvanicum* Linnaeus, Polygonaceae, as *atrata*), 73 (visiting flowers of *Polygonum hydropiperoides* Michx., Polygonaceae); G. Peckham and E. Peckham, 1898:171 (nesting habits, as *atrata*); Ashmead, 1899d:353 (in checklist of North American Sphecidae); Bridwell, 1899:209 (Kansas: Baldwin, as *atrata*); J. Smith, 1900:523 (in list of insects of New Jersey, as *atrata*); Adlerz, 1904:138 (known prey: tettigoniids); Hart, 1907:255 (Illinois); Rau and Rau, 1918:159 (nest digging and closing, prey: *Dissosteira carolina* (Linnaeus), Acrididae, nest structure, nest parasites, as *atratum*); Rau, 1922:23 (USA: Missouri: Saint Louis, prey); G. Carpenter, 1930b:294, 295 (nest closure); Rau, 1938b:541 (sleeping habits, as *atratum*); Strickland, 1947:128 (Canada: Alberta: Lethbridge, Medicine Hat); Evans and Lin, 1956a:142 (description of larva); R. Bohart, 1958b:92, 93 (in key to North American *Prionyx*); Evans, 1958a:178 (nesting behavior), 1959b:147 (additional larval characters); F. Parker, 1960:206, 207 (in key to North American *Prionyx*, as *atrata*). – **As *Chlorion atratum***: Fernald, 1906:338 (new generic assignment, in revision of Sphecini of North America and West Indies); H. Smith, 1908b:332 (in revision of Nebraskan Sphecidae); J. Smith, 1910:677 (in new list of insects of New Jersey); Rohwer, 1916b:679 (in catalog of Hymenoptera of Connecticut); Stevens, 1917:420 (North Dakota); Mickel, 1918b:397 (in catalog of Nebraskan Sphecidae); Washburn, 1919:222 (in list of Hymenoptera of Minnesota); Carter, 1925:132 (Canada: Alberta); Rohwer in Viereck, 1925:679 (in key to Sphecidae of Connecticut); J.Ch. Bradley, 1928:1011 (in catalog of New York Sphecidae); Fernald, 1931a:441 (Oregon, Washington); Krombein, 1936:98 (New York: Buffalo; floral records); Brimley, 1938:444 (North Carolina: Highlands, Raleigh, Wrightsville); Britton, 1938:149 (in addition to checklist of insects of Connecticut); Dreisbach, 1944:268 (in key to Sphecinae of Michigan), 272 (Michigan: locality records); Strandtmann, 1945a:308 (Texas, nest and prey); Spencer and Wellington, 1948:10 (British Columbia); Krombein, 1958f:191 (in supplement to catalog of North American Hymenoptera: description of larva by Evans and Lin, 1956a, reported); Piek and Spanjer, 1986:182 (in list of Sphecidae with known prey). – **As *Prionyx atratus***: R. Bohart and Menke, 1963:154 (new generic assignment, in revision of Nearctic Sphecini); Lavigne and Pfadt, 1966:31 (Wyoming; preying on *Melanoplus sanguinipes* (Fabricius), Acrididae); Pilon and Steiner, 1966:483 (locality records from Michigan and Quebec); Horning and Barr, 1970:104 (USA: Idaho: Craters of the Moon National Monument); R. Bohart and Menke, 1976:131 (in checklist of world Sphecidae); Kumar, Lavigne, Lloyd, and Pfadt, 1976:51 (Colorado: Pawnee National Grassland); Krombein, 1979b:1585 (in catalog of North American Hymenoptera); Finnamore, 1982:18 (Canada: in Sphecid Fauna of southern Quebec); Brockmann, 1985b:312 (nest closure summary); Radović, 1985:64 (sting apparatus analyzed); Piek and Spanjer, 1986:188 (in list of Sphecidae with known prey, as *astratus*); Steiner, 1986:96 (references to papers on nesting habits); O'Brien, 1989b:206 (distribution in Michigan); Spofford, Kurczewski, and Downes, 1989:256 (reference to publication on nest parasite *Metopia argyrocephala* (Meigen), a miltogrammine fly); Kurczewski and Acciavatti, 1990:59 (New York: Cayuga County); Betz, Struven, Wall, and Heitler, 1994:49 (pollinating *Asclepias verticillata* L., Apocynaceae); Ahlstrom, 1995:107 (in checklist of insects of North Carolina); O'Neil, 1995:248 (Montana, list of grasshopper prey); Blades and Maier, 1996:71 (Canada: British Columbia: Osoyoos - Mount Kobau area at 119°40'W 49°05'N); Kurczewski, 1998d:250 (pine barrens in upstate New York); Ruíz Cancino, Coronado Blanco, Varela Fuente, and Horta Vega, 2002:669 (in checklist of Mexican Sphecidae); Buck, 2004:24 (Canada: in checklist of Sphecidae of Ontario); Hua, 2006:276 (in list of Chinese insects, geographic distribution, obviously in error); Miller, Pearce, and O'Neill, 2009:3 (known to be parasitized by *Paraxenos duryi* (Pierce), Strepsiptera); Ratzlaff, 2016:29 (Canada: British Columbia: from Spencer and Wellington, 1948); R. Bohart and Menke, 1963, and

Buck, 2004); Danilov, 2019:78 (member of subgenus *Priononyx*); Danilov and Odintsev, 2023:433 (in subgenus *Priononyx*); Buck and Bennett, 2025:224 (in checklist of Sphecidae sensu lato of Canada, Alaska, and Greenland: Canada: Alberta, British Columbia, Ontario, Quebec, Saskatchewan, first record from Manitoba: no specific locality).

Sphex labrosus Harris, 1835:588. Nomen nudum. Synonymized with *Chlorion atratum* by Fernald, 1906:338.

Priononyx brunnipes Cresson, 1873:213, ♂. Holotype: ♂, USA: Texas: Bosque County: no specific locality (USNM). Synonymized with *Chlorion atratum* by Fernald, 1906:338. – Cresson, 1887:276 (in catalog of North American Hymenoptera); Ashmead, 1899d:353 (in checklist of North American Sphecidae, as *brunneipes*); Cresson, 1916:93 (holotype in USNM). – **As *Sphex brunnipes***: Kohl, 1890b:440 (new generic assignment, original description copied); Dalla Torre, 1897:417 (in catalog of world Hymenoptera); J. Smith, 1900:523 (in list of insects of New Jersey, as *brunneipes*).

3. ? *australis* (de Saussure)

Harpactopus australis de Saussure, 1867:42, ♀. Holotype or syntypes: ♀, Nova Hollandia, now Australia: no specific locality, in error (MHNG). – Froggatt, 1892:210 (in catalog of Australian Hymenoptera); R. Turner, 1910a:344 (as a synonym of *Prionyx globosus*); Danilov and Byvaltsev, 2020:402 (unrecognizable species). – **As *Sphex australis***: Kohl, 1890b:367 (new generic assignment, in revision of world Sphecini); Dalla Torre, 1897:416 (in catalog of world Hymenoptera); Montague, 1914:649 (Australia: Western Australia: Monte Bello Islands). – **As *Palmodes australis***: van der Vecht, 1973:352 (new generic assignment, type locality in error, type specimen in MHNG, not NHMW); Danilov and Odintsev, 2023:432 (in subgenus *Harpactopus*). – **As tentative subspecies of *Palmodes occitanicus***: R. Bohart and Menke, 1976:127 (new status, original type locality erroneous, in checklist of world Sphecidae). – **As *Palmodes occitanicus australis***: Dollfuss, 1989:11 (type material in NHMW); Danilov, 2019:77 (a subspecies of *Palmodes occitanicus*).

4. *bifoveolatus* (Taschenberg)

Priononyx bifoveolatus Taschenberg, 1869:408, ♂ (as *bifoveolata*, incorrect original termination). Syntypes: ♂, Brazil: Rio de Janeiro: Nova Friburgo (Halle). – F. Lynch Arribálzaga, 1878:329 (Argentina: Buenos Aires area); W. Fox, 1897b:378 (Brazil: Chapada and Corumbá); Harrington, 1902:224 (Canada: Ontario: Ottawa); Hart, 1907:255 (Illinois); Rau, 1922:23 (USA: Missouri: Saint Louis, prey carrying); K. Cooper, 1950:105 (Massachusetts: Cape Code: Woods Hole). – **As *Chlorion bifoveolatum***: Fernald, 1906:346 (new generic assignment, in revision of Sphecini of North America and West Indies); H. Smith, 1908b:333 (in revision of Nebraskan Sphecidae); J. Smith, 1910:677 (in new list of insects of New Jersey); Rohwer, 1916b:679 (in catalog of Hymenoptera of Connecticut); Strand, 1916b:99 (Kohl's 1890 reference to North America is in error); Mickel, 1918b:398 (in catalog of Nebraskan Sphecidae); Britton, 1920:340 (in checklist of insects of Connecticut); Carter, 1925:132 (Canada: Alberta; as *bifoveolatum* Fabricius); Rohwer in Viereck, 1925:679 (in key to Sphecidae of Connecticut; Milford); J.Ch. Bradley, 1928:1011 (in catalog of New York Sphecidae); Fernald, 1931a:441 (as synonym of *Chlorion pubidorsum*); Spencer and Wellington, 1948:10 (British Columbia). – **As *Sphex bifoveolatus***: Kohl, 1890b:360 (new generic assignment, in revision of world Sphecini); Dalla Torre, 1897:417 (in catalog of world Hymenoptera); J. Smith, 1900:522 (in list of insects of New Jersey); Berland, 1926c:204 (Mexico: Guadalajara). – **As *Prionyx bifoveolatus***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Genise, 1981b:19 (influence of meteorological factors on activity); Vardy, 1995a:12 (sleeping aggregation); Amarante, 2002:72 (in catalog of Neotropical Sphecidae); Dollfuss, 2008b:1408 (locality records from Chile); Buys, 2009e:277 (Brazil: Rio de Janeiro: Nova Friburgo, Petrópolis, Teresópolis); López García, Mazzitelli, Frutos, González, Marcucci, Giusti, Alemanno, del Barrio, Portela, and Debandi, 2019:314 (Argentina: Mendoza: departamento de Tupungato: vineyards in Gualtallary); Danilov, 2019:78 (member of subgenus *Priononyx*); Danilov and Odintsev, 2023:433 (in subgenus *Priononyx*).

Sphex striatulus Brèthes, 1908:147, ♀, ♂. Lectotype: ♂, Argentina: Buenos Aires (MACN), designated by Menke in R. Bohart and Menke, 1976:133. Synonymized with *Prionyx bifoveolatus* by Menke in R. Bohart and Menke, 1976:133. – Liebermann, 1931:23 (in revision of Argentinean Sphecini). – **As *Priononyx striatulus***: Jörgensen, 1912:286 (new generic assignment, Argentina: Mendoza Province); Schrottky, 1913a:225 (Argentina); Evans, 1958a:185 (observations by Lieber-

man, 1931). – **As *Chlorion striatulum***: Willink, 1948a:315 (new generic assignment, diagnostic characters), 318 (differences between *Chlorion thomae* and *Ch. striatulum*), 320 (in key to *Chlorion thomae* species group), 1951:190 (in revision of Argentinian Sphecini); Zapata, 1974:37 (Chile: Lampa near Santiago).

Sphex subexcisus Brèthes, 1908:148, ♀, ♂. Lectotype: ♀, Argentina?: no specific locality (MACN), designated by Menke in R. Bohart and Menke, 1976:133. Synonymized with *Chlorion striatulum* by Willink, 1948a:316, and with *Prionyx bifo-veolatus* by Menke in R. Bohart and Menke, 1976:133. – Liebermann, 1931:80 (in revision of Argentinean Sphecini). – **As *Priononyx subexcisus***: Schrottky, 1913a:225 (new generic assignment, Argentina).

Sphex wagneri Berland, 1926c:204, ♀, ♂. Lectotype: ♂, Argentina: Santiago de Estero: Icano (MNHN), designated by Menke in R. Bohart and Menke, 1976:133. Synonymized with *Chlorion striatulum* by Willink, 1948a:316 and with *Prionyx bifo-veolatus* by Menke in R. Bohart and Menke, 1976:133. – Liebermann, 1931:25 (in revision of Argentinian Sphecini).

Sphex caridei Liebermann, 1931:85, ♀, ♂. Holotype: ♀, Argentina: Buenos Aires: Partido de Guaminí: Distrito Casbas: estancia La Flora (MACN). Synonymized with *Chlorion striatulum* by Willink, 1948a:316 and with *Prionyx bifo-veolatus* by Menke in R. Bohart and Menke, 1976:133.

5. *binghami* Jha and Farooqi

Prionyx binghami Jha and Farooqi, 1996:15, ♀. Holotype: ♀, India: Bihar: Pusa (depository?). – Danilov and Odintsev, 2023:432 (in subgenus *Harpactopus*).

6. *canadensis* (Provancher)

Priononyx canadensis Provancher, 1887:258, ♀. Lectotype: ♂ [sic], Canada: Ontario: Ottawa (Laval Univ.), designated by Gahan and Rohwer, 1918:170. – F. Parker, 1960:206, 207 (resurrected from synonymy, in key to North American *Priononyx*). – **As *Sphex canadensis***: Kohl, 1890b:360, footnote (new generic assignment, original description copied, as tentative synonym of *Sphex bifo-veolatus*). – **As *Chlorion canadense***: Fernald, 1906:346 (new generic assignment, as synonym of *Chlorion bifo-veolatum*). – **As *Prionyx canadensis***: R. Bohart and Menke, 1963:157 (new generic assignment, in revision of Nearctic Sphecini); Lavigne and Pfadt, 1966:31 (Wyoming; preying on *Aulocara ellioti* (Thomas), Acrididae); Horning and Barr, 1970:104 (USA: Idaho: Craters of the Moon National Monument); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Krombein, 1979b:1585 (in catalog of North American Hymenoptera); Rust, Hanks and Bechtel, 1983:405 (Nevada: Churchill County: Sand Mountain); Rust, Menke, and Miller, 1985:46 (California: Channel Islands); Blades and Maier, 1996:71 (Canada: British Columbia: Osoyoos - Mount Kobau area at 119°40'W 49°05'N); Weissmann and Kondratieff, 1999:78 (Colorado: Great Sand Dunes National Monument); Ohl and Linde, 2003:149 (number of ovarioles, identification tentative); Buck, 2004:24 (Canada: in checklist of Sphecidae of Ontario), 74 (R. Bohart and Menke, 1963, map indicates two Ontario localities: London and Point Pelee); Horta Vega, Pinson Domínguez, Barrientos Lozano, and Correa Sandoval, 2007:48 (Mexico: Tamaulipas); Ratzlaff, 2016:29 (Canada: British Columbia: from Kohl, 1890b, Spencer and Wellington, 1948, and Buck, 2004); Danilov, 2019:78 (member of subgenus *Priononyx*); Danilov and Odintsev, 2023:433 (in subgenus *Priononyx*); Buck and Bennett, 2025:224 (in checklist of Sphecidae sensu lato of Canada, Alaska, and Greenland: Canada: Alberta, British Columbia, Manitoba, Saskatchewan: no specific locality).

Sphex excisus Kohl, 1890b:362, ♀, ♂. Syntypes: Canada: British Columbia: Island of Vancouver (NHMW). Synonymized with *Chlorion pubidorsum* by Fernald, 1931a:441 and with *Prionyx canadensis* by R. Bohart and Menke, 1963:157. – Dalla Torre, 1897:422 (in catalog of world Hymenoptera); Fernald, 1906:417 (unidentified species); Dollfuss, 1989:12 (type material in NHMW). – **As *Priononyx excisus***: Ashmead, 1899d:353 (new generic assignment, in checklist of North American Sphecidae).

7. *chobauti* (Roth)

Sphex chobauti Roth, 1925:388 (as *Chobauti*, incorrect original capitalization). Syntypes: ♀, ♂, Algeria: Ain Sefra and Sidi-bou-Rziguine; Morocco: Dar-Salem (MNHN). – Berland, 1926c:200 (Morocco: locality records); de Beaumont, 1951e:268 (Morocco); Leclercq, 1955h:26 (bibliographic references, faunal records); de Beaumont, 1956a:181 (Libya; as

chobauti subsp.). – **As *Priononyx chobauti***: de Beaumont, 1968b:150 (new generic assignment, member of *macula* species group). – **As *Prionyx chobauti***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Danilov and Odintsev, 2023:432 (in subgenus *Harpactopus*).

8. *crudelis* (F. Smith);

Sphex rufipennis Fabricius, 1793:200, sex not stated, junior primary homonym of *Sphex rufipennis* De Geer, 1778. Syntypes: ♀, India: Tamil Nadu: Tranquebar (ZMUC). – Fabricius, 1796:156 (in Index to his Entomologia Systematica, 1793); Turton, 1801:485 (redescription); Lepeletier de Saint Fargeau and Serville, 1828:462 (in list of *Sphex*); Guérin-Ménéville, 1829a:562 (is a member of *Sphex*); Dahlbom, 1845:XXI (specimens in collection Fabricius), 436 (in key to world *Sphex*); Lepeletier de Saint Fargeau, 1845:334 (in revision of world Hymenoptera); F. Smith, 1856:252 (in catalog of Hymenoptera in British Museum, probably a Brazilian species); A. Costa, 1864a:60 (three specimens from Mexico in Napoli Museum, clearly in error) A. Costa, 1864b:112 (two specimens from Guadalupe in Museo Zoologico di Napoli, clearly in error); Taschenberg, 1869:411 (redescription based on specimens from Brazil); nec Kohl, 1885b:198 (= *Sphex luteipennis*); Ed. André, 1888:150 (in revision of Sphecidae of Europe and Algeria), 9* (bibliographic references); Cameron, 1889c:108 (in list of Sphecidae of Oriental Region), 112 (records from South America refer to a different species; specimens described by Kohl, 1885b:198, are *luteipennis*); Kohl, 1889a:26 (specimens treated under this name by André, 1888, are several species); Kohl, 1890:408 (as a variety of *Sphex argentatus* which = *Sphex diabolicus*); Casolari and Casolari Moreno, 1980:102 (specimens in M. Spinola collection, Torino); Pagliano, 2008:524, 526 (specimens in M. Spinola collection, Torino). – **As *Sphex umbrosus* var. *rufipennis***: Ashmead, 1904d:150 (new status, Philippines).

? *Sphex hirtipes* Fabricius, 1793: 207, sex not stated. Holotype or syntypes: Guinea (type lost), junior primary homonym of *Sphex hirtipes* De Geer, 1778. Tentatively synonymized with *Harpactopus crudelis* by Kohl, 1890:351. – Fabricius, 1796:155 (in Index to his Entomologia Systematica, 1793); Kohl, 1890b:351 (as tentative synonym of *Sphex aegyptius* = *Prionyx crudelis*), 445 (original description copied); Dalla Torre, 1897:425 (in catalog of world Hymenoptera); van der Vecht, 1961a:32 (type material lost, may be a synonym of *Sphex obscurus*); R. Bohart and Menke, 1976:115 (as tentative synonym of *Sphex obscurus*); Krombein, 1984a:9 (cannot be synonym of *Sphex obscurus*); Casolari and Casolari Moreno, 1980:102 (specimens in M. Spinola collection, Torino); Pagliano, 2008:524 (specimens in M. Spinola collection, Torino). – **As *Pepsis hirtipes***: Fabricius, 1804:212 (new generic assignment, redescription).

Sphex aegyptius Lepeletier de Saint Fargeau, 1845:356, sex not stated (as *Aegyptia*, incorrect original capitalization and termination), junior primary homonym of *Sphex aegyptius* Linnaeus, 1758. Lectotype: ♀, Egypt: no specific locality (M. Spinola collection, Torino), designated by Menke in Bohart and Menke, 1976:133. – Taschenberg, 1869:412 (redescription, as *aegyptica*); Kohl, 1885b:181 (in revision of Palearctic *Sphex*); Ed. André, 1888:148 (in revision of Sphecidae of Europe and Algeria), 9* (bibliographic references); Cameron, 1889c:106 (in list of Sphecidae of Oriental Region); Kohl, 1889a:25 (comparison with *Sphex subfuscatus*), 1890b:351 (in revision of world Sphecini); Magretti, 1892 (Somalia: Herrer el Saghir); de Saussure, 1892:424 (Madagascar and Mauritius, redescription); Kohl, 1893e:183 (Tanzania: Bagamoyo); Bingham, 1897:245 (in revision of wasps and bees of British India); Dalla Torre, 1897:413 (in catalog of world Hymenoptera); Kohl, 1906a:197 (Yemen: Aden); Pérez, 1907:495 (Oman: Dibba); Kohl, 1909:370 (Comoros); R. Turner, 1911b:370 (Seychelles Islands); Maidl, 1913:560 (Egypt: Helwan); Storey, 1916:107 (Egypt: everywhere); Strand, 1916b:102 (German East Africa, now Tanzania); Fahringer and Friese, 1921:160 (Turkey: Erzurum: Djihan valley in Amanus Mountains = Gavur Dağları); Maidl, 1924:246 (Sudan: Bara, Gullfan); Berland, 1926c:200 (miscellaneous locality records); von Schulthess, 1927:299 (Iran: Bushehr); Guiglia, 1928:500 (Somalia); Schoutedden, 1930:95 (Zaire); Guiglia, 1932:124 (Somalia: Brava; Equatorial Africa: Lado); Gussakovskij, 1933b:372 (Iran); C. Williams, 1930:56 and 1933:474 (as *aegyptiacus*, following swarm of locust *Schistocerca gregaria* Forsk. in Amani, Tanzania); Giordani Soika, 1939c:105 (Eritrea: Keren, as *aegyptium*); Pittioni, 1950:21 (Cyprus); Atanassov, 1955:205 (first record from Bulgaria: Pirin); Haskell, 1955:284 (accompanying swarms of *Schistocerca gregaria* Forsk. in Kenya and Tanzania); Berland, 1956:1169 (in revision of African Sphecini); Bradley, 1957:40 (Lepeletier de Saint Fargeau's specimens in M. Spinola

collection, Torino); Myartseva, 1963b:59 (Turkmenistan: lower Murgab River); Guiglia, 1968:164 (Yemen: Taiz, as *aegyptius* Kohl); Kazenas, 1969a:22 (Kazakhstan: Mangyshlak Peninsula, Golodnaya Step'); I. Robertson, 1969:480 (Tanzania: Ukiriguru, as *aegyptiacus*); Chhotani and Ray, 1975:27 (India: Rajasthan: Sambhar Lake); Georghiou, 1977:191 (Cyprus); Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino; as *aegyptia*); Piek and Spanjer, 1986:189 (in list of Sphecidae with known prey); Pagliano, 2008:535 (specimens in M. Spinola collection, Torino, as *aegyptia*). – **As *Priononyx aegyptia***: F. Smith, 1856:266 (new generic assignment, in catalog of Hymenoptera in British Museum). – **As *Chlorion aegyptium***: Arnold, 1928c:359 (new generic assignment, description by Kohl, 1890b, translated into English), 1930:17 (in checklist of Afrotropical Sphecidae); Scott in Arnold, 1933a:370 (Ethiopia); Arnold, 1935b:1 and 8 (Mauritania: Nema); Guiglia, 1940e:293 (Italian Somalia: no specific locality).

Harpactopus crudelis F. Smith, 1856:264, ♀. Holotype or syntypes: ♀, India: Madras (BMNH). Synonymized with *Sphex aegyptius* Lepeletier de Saint Fargeau by Kohl, 1885b:181, and with *Sphex rufipennis* Fabricius by van der Vecht, 1961a:34. – F. Smith, 1871a:362 (in catalog of Oriental Aculeata); Walker, 1871:20 (Sinai Peninsula: Wadi Hebran; Africa: Red Sea coast: Island of Akeek, Harkeko); Magretti, 1884a:249 (Ethiopia), 1884c:582 (Ethiopia: Metemma); Innes Bey, 1912:110 (specimens recorded by Walker, 1871, now destroyed by dermestids, came from unknown locality). – **As *Sphex crudelis***: de Beaumont, 1949a:127 (new generic assignment, *Sphex crudelis*, and not *soror*, is the correct name for *aegyptius* Lepeletier de Saint Fargeau, nec Linnaeus), 1950d:7 (Egypt: Siwa oasis); de Beaumont and Bytinski-Salz, 1955:41 (Israel); Leclercq, 1955h:25 (bibliographic references, faunal records from Africa); de Beaumont, 1960a:5 (Greece: Island of Rhodes), 1961e:2 (Iraq); Leclercq, 1961d:108 (Madagascar); de Beaumont, 1962c:221 (Arabia: El Riyadh); Diniz, 1964b:237 (Guinea Bissau, redescription); Myartseva, 1964:73 (nesting habits in Turkmenistan); Iwata, 1965:106 (number of oocytes); Myartseva, 1965:82 (Turkmenistan: Akibay; Sakar-Chaginsk district; Murgab district); de Beaumont, 1966:211 (Egypt: Abukir), 1967a:273 (Turkey), 1968b:149 (member of *subfuscatus* species group); Guiglia, 1968:164 (Yemen); de Beaumont, 1970c:4 (Iran: Khorassan); Myartseva, 1971b:180 (Myartseva, 1971b:80 (Turanian species, ranging to Kashmir and tropical Africa); Erlandsson, 1974:58 (Greece); Kazenas, 1978b:41 (in key to Sphecidae of Kazakhstan and Central Asia). – **As *Chlorion crudele***: Derwesh, 1965:70 (new generic assignment, Iraq: no specific locality). – **As *Prionyx crudelis***: Myartseva, 1966:48 (new generic assignment, preying on orthopterans), 49 (Turkmenistan: lower course of Murgab and Tedjen rivers; preying on adult acridids *Anacridium aegyptium aegyptium* (L.) and *Calliptamus italicus italicus* (L.)), 1972a:84 (Turkmenistan); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Guichard, 1988a:121 (Arabian Peninsula); Al-Houty, 1989:162 (Kuwait: Kathma); Jha and Farooqi, 1994:11 (description and illustration of male genitalia); Roche and Zalut, 1994:113 (Egypt: Sinai Peninsula); van Vondel, 1995:29 (specimens from India in Natuurmuseum Rotterdam); Al-Houty, 1997:161 (Kuwait: no specific locality); Nazarova and Shomirsaidov, 1997:23 (Tajikistan: fruit tree orchards in Vakhsh River valley); Nazarova, 1998:40 (Tajikistan: Tigrovaya Balka Nature Reserve); Ivanov and Ljubomirov, 2001:210 (Bulgaria: Kresna Gorge at 41°48'N 23°10'E); Kazenas, 2001b:14 (in checklist of Sphecidae of Kazakhstan and Central Asia), 84 (nesting habits); Madl, 2001:1109 (Al-dabra Island group); Seyoum and Pulawski, 2001:321 (potential control agent of acridid pests in Ethiopia); Kazenas, 2002a:28 (geographic distribution, collecting localities in Kazakhstan); Ohl and Linde, 2003:149 (number of ovarioles); Pulawski, 2003b:795 (in checklist of Malagasy Sphecidae); Gadallah and Assery, 2004a:221 (in catalog of Sphecidae of Saudi Arabia); Nazarova, 2005:93 (alfalfa fields in southwestern Tajikistan); Tezcan, Yildirim, Anlaş, and Beyaz, 2006:58 (Turkey: Manisa: Turgutlu: Çıkrıkçı, on flowers of *Coridothymus capitatus*); Roche, 2007a:39 (in checklist of Egyptian Sphecidae, redescription), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Dollfuss, 2008b:1409 (Tanzania: 35 km north of Dodoma); Ljubomirov and Yildirim, 2008:26 (in catalog of Sphecidae of Turkey); Bitsch, 2010:105 (in supplement to vol. II of Faune de France, 1997: taxonomic history, morphological characteristics, geographic distribution, recent record from Bulgaria); Danilov, 2010b:44 (distribution of Palearctic-Ethiopian type); Sakenin, Samin, and Bagriacik, 2010:17 (Iran: Khuzestan: Khorramshahr); Murai and Amr, 2011:120 (recorded from Syria by de Beaumont and Bytinski-Salz, 1955); Schmid-Egger, 2011b:603 (recorded from United Arab Emirates by Guichard, 1988a); Danilov, 2012a:163, 164 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:62 (bibliographic references, geographic distribution); Gadallah, Al Dhafer,

Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia); Ebrahimi, 2014:21 (Iran: Kermān and Kermānshāh provinces); Madl, 2014a:1018 (in catalog of Ampulicidae, Crabronidae, and Sphecidae of Madagascar, with synonymy and locality records), 2014b:23 (Madagascar: Fort Dauphin); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces); Augul, Abdul-Rassoul, and Kaddou, 2015:116 (in key to Sphecini of Iraq; Iraq: locality records), 117, 118 (illustrations); Koçak and Kemal, 2015:278 (in checklist of Hymenoptera of Turkey); Arens, 2017a:630 (first record from continental Greece: Peloponnesus); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:25 (Iran: known from Kerman, Kermanshah, Khuzestan, Sistan-o Baluchestan, and South Khorasan provinces); Augul, 2019:496 (Iraq: Baghdad Province: Bab Al Muadham); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Can and Gülmez, 2019:349 (in key to *Prionyx* of Turkey); Bitsch, Barbier, and Jacobs *in* Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:138 (in Sphecid Fauna of Europe); Gadallah, 2020d:85 (in list of aculeate wasps of Arabian Peninsula); Örgel, Anlaş, and Tezcan, 2020:638 (Turkey: Manisa Province); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

Sphex grandis Radoszkowski, 1876b:132, ♂. Holotype or syntypes: ♂, Ethiopia: no specific locality (Kraków). Synonymized with *Sphex aegyptius* by Ed. André, 1888:*9.

? *Sphex aegyptius* var. *turcomanicus* Radoszkowski, 1893a:58, sex not stated. Syntypes: Turkmenistan: Serax (Kraków). Synonymized with ..

As *Sphex soror*: Honoré, 1944a:69 (in revision of Egyptian Sphecini), present correction.

9. *damascenus* (de Beaumont)

Sphex damascenus de Beaumont, 1968b:154, ♀, ♂. Holotype: ♀, Syria: Mezze near Damascus (Lausanne or A. Mochi coll., now Torino). – As *Prionyx damascenus*: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Murai and Amr, 2011:120 (recorded from Syria by de Beaumont, 1968b); Danilov and Odintsev, 2023:432 (in subgenus *Calosphex*).

10. *elegantulus* (R. Turner)

Sphex elegantulus R. Turner, 1912g:369, ♀. Holotype or syntypes: ♀, China: Lo-Fou Mountains: no specific locality (BMNH). – As *Prionyx elegantulus*: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Hua, 2006:276 (in list of Chinese insects, geographic distribution); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*).

11. *erythrogaster* (Rohwer)

Callosphex [sic] *erythrogaster* Rohwer, 1913:450, ♀ (as *erythrogastra*, incorrect original termination). Holotype: ♀, Peru: Cuzco (USNM). – As *Prionyx erythrogaster*: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Amarante, 2002:72 (in catalog of Neotropical Sphecidae); Rasmussen and Asenjo, 2009:16 (in checklist of Crabronidae of Peru, as *erythrogastra*); Danilov and Odintsev, 2023:433 (in subgenus *Neosphex*).

12. *fervens* (Linnaeus)

Sphex fervens Linnaeus, 1758:569, sex not stated. Holotype: ♀, India, actually West Indies: no specific locality (Museum Ludovicae Ulricae, Uppsala). – Linnaeus, 1764:406 (in museum of Queen Ludovica Ulrica, redescription); Fabricius, 1775:347 (redescription); Christ, 1781:294 (redescription); Fabricius, 1781:444 (redescription), 1787:275 (redescription); Gmelin, 1790:2726 (redescription); Fabricius, 1793:200 (redescription), 1796:155 (in Index to his Entomologia Systematica, 1793); Turton, 1801:484 (redescription); Jurine, 1807:129 (in list of *Sphex*); Erichson, 1849:589 (British Guyana); Kohl, 1890b:334 (in revision of world Sphecini, as unrecognizable species, Conil's description of *Enodia fervens* copied); Dalla Torre, 1897:422 (in catalog of world Hymenoptera, *Enodia fervens* of Conil listed as separate species); Schrottky, 1903b:123 (in checklist of Hymenoptera of Argentina, Paraguay, and Uruguay, referring to *Enodia fervens* of Conil, 1880); W. Schulz, 1912:56 (study of type); van der Vecht, 1959b:130 (identification of species by W. Schulz, 1912, was correct); Day and Fitton, 1978:193 (curation of Linnean type material); Day, 1979:62 (taxonomic

- history); Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino); Pagliano, 2008:533 (specimens in M. Spinola collection, Torino). – **As *Pepsis fervens***: Dahlbom, 1845:XXI (specimens in collection Fabricius are *Enodia fervens*). – **As *Enodia fervens***: Dahlbom, 1845:XXI (new generic assignment), 439 (in key to world *Enodia*); Fairmaire, 1858:264 (Gabon, clearly in error); A. Costa, 1864b:112 (15 specimens from Senegal in Museo Zoologico di Napoli, clearly in error); Conil, 1880:241 (nesting habits and redescription), 1881:454 (same information). – **As *Parasphepex fervens***: F. Smith, 1856:267 (new generic assignment, in catalog of Hymenoptera in British Museum, 1871a:362 (in catalog of Oriental Aculeata); Walker, 1871:20 (Sinai Peninsula: Tor; Djibouti: Tajura; Africa: Red Sea coast: Harkeko, clearly in error); Magretti, 1884a:249 (Sudan, clearly in error), 1884c:582 (Sudan: Kassala, clearly in error, as *fervens* Fabricius); Innes Bey, 1912:111 (specimens recorded by Walker, 1871, now destroyed by dermestids, were collected at Tor, Sinai Peninsula, obviously a misidentification); Roche, 2007a:139 (recorded from Egypt, but not occurring there). – **As *Priononyx fervens***: F. Parker, 1960:206, 207 (new generic assignment, in key to North American *Priononyx*). – **As *Prionyx fervens***: R. Bohart and Menke, 1963:158 (new generic assignment, in revision of Nearctic Sphecini), 1976:133 (in checklist of world Sphecidae); Krombein, 1979b:1585 (in catalog of North American Hymenoptera); Nascimento and Overal, 1980:8 (Argentina, Brazil); Yústiz, 1987:13 (Venezuela: Central Lara Depression); F. Fernández, 1990:24 (Colombia: Meta: Parque Nacional Natural La Macarena); Hanson and Menke, 1995:637 (known from Costa Rica); Amarante, 2002:72 (in catalog of Neotropical Sphecidae); Ruíz Cancino, Coronado Blanco, Varela Fuente, and Horta Vega, 2002:669 (in checklist of Mexican Sphecidae); Buys, 2006b:311 (nesting habits, behavior of larva); Dollfuss, 2008b:1409 (Argentina: La Rioja province; Mexico: Hidalgo: Metztitlán 70 km of Pachuca); Buys, 2009e:277 (Brazil: Rio de Janeiro: Macaé, Rio de Janeiro), 2011b:2 (Brazil: Rio de Janeiro: Arraial do Cabo, Cabo Frio); Rodrigues and Buys, 2013:214 (Brazil; Espírito Santo: Serra); Buys and Rodrigues, 2014:40 (Brazil: State of Espírito Santo: Linhares, Serra); Danilov, 2019:78 (member of subgenus *Priononyx*); Buys, 2020b:80 (in cladistic analysis of larvae of Sphecidae s.s.); Danilov and Odintsev, 2023:433 (in subgenus *Priononyx*).
- Pepsis johannis* Fabricius, 1804:208, sex not stated (as *Johannis*, incorrect original capitalization). Holotype: ♀, South American Islands: no specific locality (ZMUC). Synonymized with *Sphex striatus* by F. Smith, 1756:260, 266 (tentatively), synonymy confirmed by Burmeister, 1872:239, and with *Priononyx fervens* by van der Vecht, 1961a:34. – **As *Sphex johannis***: Dalla Torre, 1897:427 (new generic assignment, in catalog of world Hymenoptera); Schrottky, 1902a:315 (Brazil), 1903b:123 (in checklist of Hymenoptera of Argentina, Paraguay, and Uruguay); Autran, 1907:207 (corrected *Sphex fervens* of Conil, 1881, to *Sphex johannis*).
- Sphex doumerci* Lepeletier de Saint Fargeau, 1845:357, ♀ (as *Doumerci*, incorrect original capitalization). Holotype: ♀, Syntypes: Brazil and Cayenne (originally Audinet-Serville coll., now M. Spinola collection, Torino). Synonymized with ... – Kohl, 1890:356, footnote (original description copied, as tentative synonym of *Sphex striatus*); Dalla Torre, 1897:421 (in catalog of world Hymenoptera); Bradley, 1957:40 (Lepeletier de Saint Fargeau's specimens in M. Spinola collection, Torino); Casolari and Casolari Moreno, 1980:103 (specimen in M. Spinola collection, Torino); Pagliano, 2008:533 (holotype in M. Spinola collection, Torino). – **As *Priononyx doumerci***: F. Smith, 1856:266 (new generic assignment, in catalog of Hymenoptera in British Museum).
- Priononyx striatus* F. Smith, 1856:266, ♀ (as *striata*, incorrect original termination). Syntypes: ♀, Brazil: Pará: Pará, now Belém; and Amazonas: Villa Nova, now Parintins (BMNH). Synonymized with *Sphex fervens* by W. Schultz, 1912:56. – Taschenberg, 1869:408 (redescription); Burmeister, 1872:239 (Argentina: Paraná, Mendoza); Schrottky, 1909b:244 (Argentina: Catamarca), 1913a:225 (Argentina, Paraguay); R. Bohart, 1958b:92, 93 (in key to North American *Prionyx*, as *striata*); Evans, 1958a:184 (nesting behavior). – **As *Sphex striatus***: Kohl, 1890b:356 (new generic assignment, in revision of world Sphecini); Dalla Torre, 1897:442 (in catalog of world Hymenoptera); Ducke, 1901:241 (Brazil: Pará: Belém); W. Schulz, 1906:193 (Argentina: Tucumán: Tapia; variation); Brèthes, 1908:144 (in revision of *Sphex thomae* group); Ducke, 1908b:82 (Brazil: Ceará State); Lüderwaldt, 1910:177 (nesting habits); Strand, 1910a:133 (Paraguay); Jörgensen, 1912:285 (Argentina: Mendoza Province); Poulton, 1918:xxxvii (Brazil, prey); Berland, 1926c:202 (miscellaneous locality records); Liebermann, 1931:24 (in revision of Argentinean Sphecini); Murray in Muesebeck, Krombein, and Townes, 1951:973 (in catalog of North American Hymenoptera). – **As *Prionyx striatus***: Snow, 1906: new generic assignment, :335

(new generic assignment, in revision of Sphecini of North America and West Indies, first record from USA), 1907:264 (Argentina), 1931a:440 (synonymy); Fernald, 1942:30 (Guyana: Kartabo); Willink, 1948a:319 (in key to *Chlorion thomae* species group), 1951:194 (in revision of Argentinean Sphecini); Krombein, 1958f:191 (in supplement to catalog of North American Hymenoptera: California).

Sphex laerma Cameron, 1897b:370, sex not stated. Holotype or syntypes: Mexico: Guerrero: Río Papagaio (BMNH). Synonymized with *Chlorion striatum* by Fernald, 1906:335 (tentatively) and 1931a:440 (definitely).

13. *foxi* R. Bohart and Menke

Sphex ferrugineus W. Fox, 1892f:170, ♀, junior primary homonym of *Sphex ferrugineus* Lepeletier de Saint Fargeau, 1845. Holotype: ♀, USA: southern California: no specific locality (USNM). – Kohl, 1895:48 (original description copied); Dalla Torre, 1897:422 (in catalog of world Hymenoptera); Murray in Muesebeck, Krombein, and Townes, 1951:973 (in catalog of North American Hymenoptera). – **As *Chlorion ferrugineum***: Fernald, 1906:331 (new generic assignment, in revision of Sphecini of North America and West Indies, description of ♂). – **As *Priononyx ferrugineus***: Ashmead, 1899d:353 (new generic assignment, in checklist of North American Sphecidae, as *ferruginosus*); F. Parker, 1960:206 (in key to North American *Prionyx*, as *ferruginea*). – **As *Parasphex ferrugineus***: Snow, 1906:133 (new generic assignment, Arizona).

Prionyx foxi R. Bohart and Menke, 1963:52. Substitute name for *Sphex ferrugineus* W. Fox. – R. Bohart and Menke, 1963:152 (in revision of Nearctic Sphecini), 1976:133 (in checklist of world Sphecidae); Krombein, 1979b:1586 (in catalog of North American Hymenoptera); Ruíz Cancino, Coronado Blanco, Varela Fuente, and Horta Vega, 2002:669 (in checklist of Mexican Sphecidae); Danilov and Odintsev, 2023:432 (in subgenus *Foxiprionyx*).

14. *fragilis* (Nurse)

Sphex fragilis Nurse, 1903b:10, ♀, ♂. Syntypes: India: Gujarat: Deesa; Pakistan: Quetta (BMNH). – Ramakrishna Aiyar, 1916:554 (in catalog of Indian aculeates described after Bingham, 1897). – **As *Prionyx fragilis***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Danilov and Odintsev, 2023:432 (in subgenus *Calosphex*).

15. *funebis* (Berland)

Sphex funebis Berland, 1926c:202, ♀. Holotype: ♀, Kenya: Bura (MNHN); paratypes: Ethiopia, South Africa. – Leclercq, 1955h:26 (bibliographic references, locality records); de Beaumont, 1967b:502 (Namibia). – **As *Chlorion funebis***: Arnold, 1928c:358 (new generic assignment, in revision of Sphecini of southern Africa), 1930:17 (in checklist of Afrotropical Sphecidae); Scott in Arnold, 1933a:370 (described from Ethiopia). – **As *Prionyx funebis***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Seyoum and Pulawski, 2001:321 (potential control agent of acridid pests in Ethiopia); S. Gess and F. Gess, 2003:91 (Namibia and South Africa: visiting flowers of unnamed Acanthaceae, *Asclepias buchenaviana* Schinz, Apocynaceae, *Sarcostemma viminalis*, Apocynaceae, *Leucas pechuelii* (Kuntze) Guerke, Lamiaceae, *Acacia karroo* Hayne, Fabaceae, and *Medicago sativa* L., Fabaceae); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

16. *globosus* (F. Smith)

Sphex globosus F. Smith, 1856:251, ♀, ♂ (as *globosa*, incorrect original termination). Syntypes: Australia: Van Diemen's Land, now Tasmania: no specific locality (BMNH). – Kohl, 1890b:368 (in revision of world Sphecini); Froggatt, 1892:210 (in catalog of Australian Hymenoptera); Dalla Torre, 1897:424 (in catalog of world Hymenoptera); R. Turner, 1910a:343 (in key to Australian Sphecini); Berland, 1926c:206 (Tasmania). – **As *Chlorion globosus***: R. Turner, 1915b:551 (new generic assignment, Tasmania, continental Australia); Chandler, 1928:177 (Australia: Victoria: Red Cliffs; nesting behavior). – **As *Prionyx globosus***: R. Bohart and Menke, 1963:152 (new generic assignment, single member of *Prionyx globosus* species group); Riek, 1970:940 (in Insect Fauna of Australia); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Evans, Hook, and Matthews, 1982:223 (nesting behavior); Callan, 1984:38 (Australia: sleeping aggregation); Cardale, 1985:227 (in catalog of Australian Sphecidae); McCorquodale and Thomson, 1989:94 (prey: Acrididae); K.

Walker, Naumann, Austin, Taylor, and Cardale, 1992:49 (in catalog of insects of Tasmania); Naumann, 1993:181 (Australia: Queensland: Heathlands area in Cape York), 1998:182 (Australia: northwestern Queensland: Musselbrook area, approximately 18°40'S 138°23'E); Pagliano, 2003a:504 (Australia); Dollfuss, 2008b:1409 (Australia: Coopers Creek); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

17. *guichardi* (de Beaumont)

Sphex guichardi de Beaumont, 1967a:273, ♀. Holotype: ♀, Turkey: Kayseri: Sultahani (BMNH). – **As *Priononyx guichardi*:** de Beaumont, 1968b:150 (new generic assignment, member of *Sphex macula* species group). – **As *Prionyx guichardi*:** R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Ljubomirov and Yildirim, 2008:26 (in catalog of Sphecidae of Turkey); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey), Can and Gülmez, 2019:348 (in key to *Prionyx* of Turkey); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

18. *haberhaueri* (Radoszkowski)

Sphex haberhaueri Radoszkowski, 1871:199, ♀ (as *Haberhaueri*, incorrect original capitalization). Holotype or syntypes: ♀, Iran: Golestan: Astrabad, now Gorgan (Kraków or ZMHU). – Kohl, 1885b:183 (in revision of Palearctic *Sphex*); Ed. André, 1888:127 (in revision of Sphecidae of Europe and Algeria); Kohl, 1889a:24 (critique of André's characteristic), 1890b:331 (in revision of world Sphecini); F. Morawitz, 1894:339 (Turkmenistan: Atrek); Dalla Torre, 1897:424 (in catalog of world Hymenoptera); Berland, 1926b:168 (one ♀ with no locality data in coll. Pérez, MNHN); Gussakovskij, 1933b:273 (Iran); de Beaumont and Bytinski-Salz, 1955:41 (Israel); Kazenas, 1969a:21 (Kazakhstan: Charyn River); de Beaumont, 1970a:391 (Afghanistan), 1970c:4 (Iran: Baluchistan); Kazenas, 1972b:11 (Kazakhstan: Charyn River in Alma Ata Oblast'); Esmaili and Rastegar, 1974:45 (Iran); Kazenas, 1978b:41 (in key to Sphecidae of Kazakhstan and Central Asia); Pulawski, 1978:184 (in key to Sphecidae of European part of USSR). – **As *Prionyx haberhaueri*:** R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Islamov, 1986:516 (Uzbekistan: Tashkent Oblast'); Kazenas, 1998b:109 (in Sphecid Fauna of Kazakhstan); Nazarova, 1998:40 (Tajikistan: Tigrovaya Balka Nature Reserve); Esenbekova and Kazenas, 2000:9 (southeast Kazakhstan: 6 km south of Podgornyi); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:28 (geographic distribution, collecting localities in Kazakhstan), 2004b:98 (Kazakhstan: western Tien Shan Mountains), 2008a:98 (Southeast Kazakhstan: River Ili valley, lower course of River Charyn, and foothills of Dzunggar Alatau: Kyzyl Agach and Ush-Tube), 2008c:255 (Kazakhstan: village Koktum south of Lake Alakol'); Danilov, 2009:54 (Russia: Western Siberia: Kulundinskaya Steppe), Danilov, 2010b:44 (distribution of Tethyan type); Kazenas, 2010a:168 (Kazakhstan: Sogety Range near village Kokpek); Danilov, 2012a:163 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:63 (bibliographic references, geographic distribution); Kazenas, 2013a:26 (color photograph of adults, short information on geographic distribution and nesting habits); Danilov, 2014b:514 (in key to Sphecidae s.s. of Siberia, not yet found in Siberia); Dunford, Turbyville, and Leavengood, 2014:11 (listed as medically important in Afghanistan); Ebrahimi, 2014:21 (Iran: Hamadān: Asadābad); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:25 (Iran: known from Golestan, Hamadan, and Sistan-o Baluchestan provinces); Kemal and Koçak, 2018:43 (Turkey: in list of pterygot insects of Van Province); Can and Gülmez, 2019:348 (in key to *Prionyx* of Turkey); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Calosphex*); Daminova, 2019:84 (endangered in Uzbekistan); Danilov, 2020:319 (specimen from Kyrgyzstan in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Danilov and Odintsev, 2023:432 (in subgenus *Calosphex*); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

Enodia vittata Kohl, 1884a:385, ♂. Syntypes: on Caspian Sea: no specific locality (NHMW). Synonymized with *Prionyx haberhaueri* by Danilov, 2012b:63. – **As *Sphex vittatus*:** Kohl, 1885b:184 (new generic assignment, in revision of Palearctic *Sphex*); Ed. André, 1888:142 (in revision of Sphecidae of Europe and Algeria), 9* (bibliographic references); Kohl, 1890b:331 (in revision of world Sphecini); Dalla Torre, 1897:447 (in catalog of world Hymenoptera); de Beaumont and

Bytinski-Salz, 1955:41 (Israel); de Beaumont, 1961b:272 (Afghanistan), 1967a:273 (Turkey); Kazenas, 1969a:21 (Kazakhstan: Sharyn River, foothills of Dzhungarian Alatau), 1972b:110 (Kazakhstan), 1978b:41 (in key to Sphecidae of Kazakhstan and Central Asia); Pulawski, 1978:184 (in key to Sphecidae of European part of USSR). – **As *Prionyx vittatus***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Dollfuss, 1989:12 (type material in NHMW); Ebrahimi, 2008:95 (first record from Iran: Hamedan: Psadabad); Ljubomirov and Yildirim, 2008:32 (in catalog of Sphecidae of Turkey); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

19. *herrerai* (Brèthes)

Sphex herrerai Brèthes, 1926:46, sex not stated (as *Herrerai*, incorrect original capitalization). Holotype or syntypes: Peru: Cuzco (Buenos Aires). – **As *Prionyx herrerai***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Genise, 1990:27 (type material in MACN); Amarante, 2002:72 (in catalog of Neotropical Sphecidae); Rasmussen and Asenjo, 2009:16 (in checklist of Crabronidae of Peru); Danilov and Odintsev, 2023:433 (in subgenus *Neosphex*).

? *Sphex villarubiai* Giner Marí, 1944:349, ♀, ♂. Syntypes: Peru: no specific locality (Mus. Barcelona). Tentatively synonymized with *Prionyx herrerai* by R. Bohart and Menke, 1976:133.

20. *indus* (Linnaeus)

Sphex indus Linnaeus, 1758:569 (as *inda*, incorrect original termination). Holotype: ♀, India: no specific locality (Museum Ludovicae Ulricae, Uppsala). – Day, 1979:65 (notes on holotype). – **As *Prionyx indus***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); S. Gess and F. Gess, 2003:91 (Namibia and South Africa: visiting flowers of *Psilocaulon salicornioides* (Pax) Schwant., Aizoaceae, *Hermbsaedia* sp., Amaranthaceae, *Asclepias buchenaviana* Schinz, Apocynaceae, and *Acacia karroo* Hayne, Fabaceae); Dollfuss, 2008b:1409 (South Africa: Vioolsdrift, Zimbabwe: Kwekwe; correction to Arnold's 1928 key); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

Sphex indostanus Linnaeus, 1764:942 (as *indostana*, incorrect original termination). Unjustified emendation of *Sphex indus*. – Linnaeus, 1764:407 (in museum of Queen Ludovicae Ulricae, redescription); Gmelin, 1790:2726 (redescription); Christ, 1791:295 and 307 (redescription); Turton, 1801:488 (redescription); Dalla Torre, 1897:427 (in catalog of world Hymenoptera); Day and Fitton, 1978:193 (recuration of Linnean type material: no specimens).

Harpactopus tyrannus F. Smith, 1856:264, ♀. Holotype: ♀, South Africa: KwaZulu-Natal: Port Natal, now Durban (BMNH). Synonymized with *Sphex indus* by .. – Cameron, 1910b:139 (South Africa: Transvaal). – **As *Sphex tyrannus***: Radoszkowski, 1881:210 (new generic assignment, Angola); Kohl, 1890b:349 (in revision of world Sphecini); Dalla Torre, 1897:445 (in catalog of world Hymenoptera); Bingham, 1902:216 (Malawi); Cameron, 1910b:139 (South Africa: Mpumalanga: Pretoria, Westfalia; Limpopo: Zoutpansberg District, now Soutpansberg); Brauns, 1911a:117 (South Africa); Leclercq, 1955h:27 (bibliographic references, faunal records from Africa, variation in wing venation), 1961b:47 (Zaire); Diniz, 1964c:100 (in key to Angolan *Sphex*), 102 (Angola: Lunda: Andrada); Casolari and Casolari Moreno, 1980:102 (specimens in M. Spinola collection; as *tyrannus*); van Vondel, 1995:24 (specimens from Namibia in Natuurmuseum Rotterdam); Pagliano, 2008:525 (specimens in M. Spinola collection, Torino, are a *Sphex* sp.). – **As *Chlorion tyrannus***: R. Turner, 1918b:361 (new generic assignment, synonymy); Arnold, 1928c:357 (in revision of Sphecini of southern Africa), 1930:17 (in checklist of Afrotropical Sphecidae), 1935a:503 (South Africa: Kalahari). – **As *Chlorion aegyptum* [sic] *tyrannum***: Arnold, 1947:145 (new status).

Sphex vagus Radoszkowski, 1881:209, ♂, junior primary homonym of *Sphex vagus* Linnaeus, 1768. Holotype or syntypes: ♂, Angola: no specific locality (Kraków). Synonymized with *Chlorion tyrannus* by R. Turner, 1918b:361. – Dalla Torre, 1897:446 (in catalog of world Hymenoptera).

Sphex englebegi Brauns, 1899:392, ♀, ♂ (as *Englebegi*, incorrect original capitalization). Syntypes: South Africa: Gauteng: Tamatsetse, Zwartkop; Free State: Bloemfontein (NHMW, AMG, TMP). Synonymized with *Chlorion tyrannum* by Arnold, 1928c:357. – Brauns, 1911a:118 (prey, following acridid swarms); Strand, 1916b:102 (German East Africa, now Tanzania: Kigonsera); Berland, 1926c:200 (South Africa and Tanzania: locality records).

21. *insignis* (Kohl)

Sphex insignis Kohl, 1885b:189, ♀. Holotype or syntypes: ♀, Syria: no specific locality (NHMW). – Ed. André, 1888:132 (in revision of Sphecidae of Europe and Algeria), 9* (bibliographic references); Kohl, 1890b:343 (in revision of world Sphecini); Dalla Torre, 1897:427 (in catalog of world Hymenoptera); Dollfuss, 1989:12 (type material in NHMW). – **As *Prionyx insignis***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Murai and Amr, 2011:120 (recorded from Syria by Kohl, 1885); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*).

22. *judaesus* (de Beaumont)

Sphex judaeus de Beaumont, 1968b:150, ♀, ♂. Holotype: ♀, Jordan: Jericho (Lausanne). – de Beaumont, 1968b:149 (member of *subfuscatus* species group). – **As *Prionyx judaeus***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Roche, 2007a:40 (in checklist of Egyptian Sphecidae, redescription), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

23. *kirbii* (Vander Linden)

Ammophila? kirbii Vander Linden, 1827:360, ♀, ♂. Syntypes: France and Spain: no specific localities (lost). – **As *Sphex kirbii***: Scobiola-Palade, 1987:65 (new generic assignment, Romania: Dobrogea), 1989:87 (Romania: delta of Danube); Kazenas, 1978b:44 (in key to Sphecidae of Kazakhstan and Central Asia, as *kirbyi*); Pulawski, 1978:183 (in key to Sphecidae of European part of USSR). – **As *Prionyx kirbii***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae, as *kirbyi*), 627 (*kirbii* is correct spelling); Guichard, 1978:270 (Greece); Valetta, 1979:215 (Malta; as *kirbi*); Pagliano, 1980:110 (Italy: Liguria, Valle d'Aosta); Gayubo, 1981a:136 (northern Spain: Sierra de Béjar); Gess, 1981:17 (South Africa: Hilton Farm 18 km west-northwest of Grahamstown; in non-friable soils nesting in old or abandoned nests of *Parachilus insignis*), 19 (digging own nests in friable soils), 54 (facultative nesting in preexisting cavities); Gayubo, 1982f:245 (Spain: Cádiz Province); Dollfuss, 1983b:2 (occurrence in Austria doubtful); Gayubo, 1983c:231 (Spain: Salamanca Province); Mingo and Gayubo, 1983:154 (Spain); Schmidt and Westrich, 1983:121 (Greece); Gayubo, 1984c:356 (Portugal: El Algarve Province); Gayubo and Tormos, 1984:9 (Spain: Valencia); Pagliano, 1984:367 (Italy); Chevin and Chevin, 1985:38 (France: Aude, as *kirbyi*); Brockmann, 1985b:312 (nest closure summary); Eiroa and Novoa, 1985:23 (Spain: Pontevedra: Barra beach near Cangas); Gayubo, 1985c:166 (Spain: Avila: Guisando; Valladolid: Traspinedo); Józán, 1985b:55 (Hungary south of Lake Balaton; as *kirbyi*), 76 (floral records), 83 (ecological and zoogeographic characteristics); Pagliano, 1985:8 (Italy); Gayubo, 1986b:36 (Spain: Andalucía), 1986c:30 (Spain: Zamora Province), 1986f:997 (prey, strepsipteran parasite); Gayubo and Heras, 1986:28 (Spain: Segovia and Valladolid Provinces; floral records); Gayubo and Sanza, 1986:27 (Spain: Burgos, Soria); Gayubo and Tormos, 1986a:8 (Spain: Castellón de la Plana), 1986b:4 (Spain: Valencia); Islamov, 1986:516 (Uzbekistan: Tashkent Oblast'); Józán, 1986:367 (Hungary: Kiskunság National Park, as *kirbyi*); Steiner, 1986:97 (references to papers on nesting habits, as *kirbyi*); Asís and Jiménez, 1987:24 (Spain: Provincia de Castellón); Gayubo, 1987:107 (Spain: Provincia de Ciudad Real); Tormos and Jiménez, 1987a:122 (Spain: Valencia; as *kirbyi*), 1987b:316 (Spain: Valencia Province: Dehesa de El Saler); Karsai, 1988:99 (Hungary: Kiskunság National Park); Gayubo, Asís, and Tormos, 1990a:10 (Spain); Dollfuss, 1990:122 (Central African Republic); Pagliano, 1990:58 (in catalogue of Italian Sphecidae); Dollfuss, 1991:29 (in key to Sphecidae of North and Central Europe); Gayubo, Borsato, and Osella, 1991:394 (Italy); Gayubo and Torres, 1991:Table I (Spain: Salamanca; effects of urban pressure); Hamon, Fonfria, and Tussac, 1991:128 (in key to French Sphecini), 133 (in France extending north to Alsace and Loire-Atlantique); Leclercq, 1991a:274 (correction to Leclercq's, 1979 catalog of French and Benelux Sphecidae: *kirbii* is correct spelling); Kazenas and Nasyrova, 1991:38 (Kazakhstan: preying on *Calliptamus barbarus*).

cephalotes F.W. and *Notostaurus albicornis* (Ev.)); Negrisolo, 1991:316 (Italy: Gorizia and Udine Provinces); Schembri, 1991:176 (recorded from Malta by Valetta, 1979); Gayubo, Borsato, and Osella, 1992:276 (France: Corse; Spain, Greece); Józán, 1992b:171 (Hungary: Boronka-melléki Protected Area, as *kirbyi*); Gayubo, Tormos, and Asís, 1993b:307 (parasitized by *Paraxenos sphecidarum* Dufour); Luchetti, 1993:104 (Italy: Sardegna: Maddalena archipelago); Mochi and Luchetti, 1993:104 (Italy; France: Corse); Torregrosa, Gayubo, Tormos, and Asís, 1993:11 (Spain: Alicante Province); Gayubo and Borsato, 1994:200 (Italy: Toscana); Roche and Zalat, 1994:113 (Egypt: Sinai Peninsula, as *kirbyi*); Tormos, Asís, and Gayubo, 1994:187, 194 (Spain: Albacete Province, nest and prey); Józán, 1995:104 (Hungary: projected Duna-Dráva National Park, as *kirbyi*); Negrisolo, 1995a:19 (visiting flowers of *Limonium bellidifolium* (Gouan) Dumort), 22 (Italy: Veneto); Negrisolo in Minelli, Ruffo, and La Posta, 1995b:2 (in catalog of Italian fauna); Pagliano and Pesarini, 1995:83 (Italy: Ferrara Province); Pagliano and Scaramozzino, 1995:730 (Italy: Island of Lampedusa); Scharrer, 1995:22 (Morocco: Ifrane and Tanger); Vernier, 1995:176 (in key to Sphecini of Switzerland); S. Gess, 1996:283 (floral records); Gusenleitner, 1996a:5 (first unquestionable record from Austria: Burgenland); Kuhlmann, 1996:220 (Portugal: Serra de Estrela); Minoranskiy and Shkuratov, 1996:81 (Russia: Rostov Oblast'); Wu and Zhou, 1996a:45 (in revision in Economic Insect Fauna of China, as *kirbyi*); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:59 (in Sphecidae Fauna of Western Europe); Stoyanov and Ljubomirov, 1997:25 (Bulgaria: Rila Mountains); Dollfuss, Gusenleitner, and Bregant, 1998:509 (Austria: summary of collecting records from Burgenland); Gusenleitner, 1998:498 (Austria: Burgenland: Nickelsdorf); González, Gayubo, and Torres, 1998:72, 73 (Spain: Valladolid Province); Józán, 1998:310 (Hungary: Duna-Dráva National Park); Kazenas, 1998b:111 (in Sphecidae Fauna of Kazakhstan, as *kirbyi*); Nazarova, 1998:40 (Tajikistan: Tigrovaya Balka Nature Reserve, as *kirbyi*); Gayubo, García, Torres, and González, 1999:89 (Spain: Soria Province); González, Gayubo, and Torres, 1999:354 (Spain: Valladolid: Viana de Cega); Zehnder and Zettel, 1999:131 (Switzerland: recolonization of flooded area in Valais Canton); Esenbekova and Kazenas, 2000:9 (southeast Kazakhstan: 10 km northwest of Chemolgan, near Kapchagay, 35-45 km northwest of Kapchagay, 45 km northwest of Suzak); Gayubo, González, and Torres, 2000:184 (Spain: Salamanca Province); Giachino, Grosso, Marchetti, Pagliano, Scaramozzino, and Vailati, 2000:104 (Greece); Józán, 2000:104 (Hungary: Bakony Mountains, as *kirbyi*); Ljubomirov, 2000:7 (Bulgaria, specimens in N. Nedelkov collection); Shkuratov, 2000:55 (Russia: Rostov Oblast': Vëshenskaya village area at 49°37'N 41°45'E); Vicidomini, 2000b:27 (Italy: previously recorded from Campania); Basset, 2001:79 (France: Département de Gironde); Ivanov and Ljubomirov, 2001:210 (Bulgaria: Kresna Gorge at 41°48'N 23°10'E); Józán, 2001:277 (Hungary: Somogy County, as *kirbyi*); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 84 (review of nesting habits); Seyoum and Pulawski, 2001:321 (potential control agent of acridid pests in Ethiopia; Ethiopia: Harerge: 37 km southeast of Jijiga at 9°11'N 43°05'E, 44 km east-northeast of Jijiga at 9°29'N 43°10'E, Sidamo: Moyale at 3°33'N 39°03'E, 4 km east of Yabelo at 4°53'N 38°08'E); Shkuratov, 2001:16 (prey: *Calliptamus italicus* L.); Kazenas, 2002a:28 (geographic distribution, collecting localities in Kazakhstan, as *kirbyi*); Shkuratov, 2002a:383 (Russia: common in Rostov Oblast'), 2002b:138 (Russia: Rostov Oblast': Rostovskiy Nature Reserve at 46°27'N 42°41'E); Drewes, 2003:141 (Spain: Barcelona, Tarragona); Generani, Pagliano, Scaramozzino, and Strumia, 2003:64 (Italy: Arcipelago Toscano); S. Gess and F. Gess, 2003:93 (South Africa: visiting flowers of unnamed Mesembryanthema, *Hermbsaedia odorata* (Burch.) T. Cooke, Amaranthaceae, *Asclepias buchenaviana* Schinz, Apocynaceae, *Lasiospermum bipinnatum* (Thunb.) Druce, Asteraceae, *Geigeria pectidea* (DC.) Harv., Asteraceae, *Bergia glomerata* L. f., Elatinaceae, *Acacia karroo* Hayne, Fabaceae, and *Zygophyllum simplex* L., Zygophyllaceae); González, Gayubo, Asís, Tormos, and García, 2003:61 (Spain: Soria: Chavaler); Józán, 2003: 226 (Hungary: Látrányi Pusztá Nature Conservation Area, as *kirbyi*); Nieves-Aldrey et al., 2003:42 (Spain: Madrid: Estación Biogeológica de El Ventorrillo in Sierra de Guadarrama); Schmid-Egger, 2003:757 (Italy: Sicilia: Ragusa); Wu, Zhou, Q. Li, and Yang, 2003:807 (China: Fujian Province); Gayubo et al. 2004:130 (Spain: Madrid: Estación Biogeológica de El Ventorrillo in Sierra de Guadarrama); Gayubo, Nieves-Aldrey, González, Tormos, Rey del Castillo, and Asís, 2004:108 (Spain: Madrid: Monte de El Pardo); Kazenas, 2004b:98 (Kazakhstan: western Tien Shan Mountains, as *kirbyi*), 2004d:26 (Kazakhstan: northern Caspian region, as *kirbyi*); Shkuratov, 2004a:73 (Russia: Rostov Oblast'), 2004b:164 (Russia: Rostov Oblast': Gosudarstvennyi Muzei-Zapovednik M.A. Sholokhova); Straka, Bo-

gusch, Tyrner, and Vepřek, 2004:146 (Czech Republic: Vojenské Cvičiště Nature Reserve); Cruz-Sánchez, Gayubo, González, and Torres, 2005:219 (Spain: Salamanca: San Martín del Castañar); Gayubo and Özbek, 2005:8 (Turkey: Antalya: Arapsuyu, Manavgat; Erzurum: University campus); Gülmez and Tüzün, 2005:47 (Turkey: Ankara Province); Jacobs, 2005a:437 (Bulgaria); Pagliano and Negrisola, 2005:53 (in Sphecidae Fauna of Italy); Shorenko, 2005a:162 (Ukraine: Crimea); Straka, 2005a:399 (critically endangered in Czech Republic); Blösch, 2006:63 (specimens spend nights attached to plant stems with their heads down); Gadoum and Barbier, 2006:42 (France: départements of Val-d'Oise and Yvelines: Parc Naturel Régional du Vexin Français); Hua, 2006:276 (in list of Chinese insects, geographic distribution); Ljubomirov, 2006:536 (Bulgaria: previous records from Rhodope Mountains summarized); Magdalou, 2006a:6, 9 (France: Pyrénées-Orientales: Réserve Naturelle de la Massane), 2006b:109 (France: Pyrénées-Orientales: Mas-Larrieu Nature Reserve near Argelès-sur-Mer), 2006c:7 (France: Pyrénées-Orientales: Réserve Naturelle du Mas-Larrieu); Polidori, Tormos, Asís, Mendiola, and Andrietti, 2006:405 (kleptoparasite on *Stizus continuus* in mixed colonies, description of mature larva); Standfuss and Standfuss, 2006c:307 (Greece: Thessalia: Magnisia Peninsula at 39°N 23°E); Burguet, 2007:unnumbered p. 5 (France: Puy-de-Dôme: sands Girauds-Faures in Orléat commune); Jacobs, 2007:42 (in key to Sphecidae of Germany, not yet found in Germany); Baños-Picón, Gayubo, Asís, and González, 2007:255, 258 (Spain: Zamora: Cabañas de Aliste); Roche, 2007a:41 (in checklist of Egyptian Sphecidae, redescription, as *kirbii kirbii*), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Sipos and Móczár, 2007:204 (Hungary: Bács-Kiskun megye: vicinity of Foktő); Vepřek and Straka, 2007:199 (in catalog of Sphecidae of Czech Republic and Slovakia: known from Moravia and Slovakia only), 211 (recently recorded from Moravia and Slovakia by Straka et al., 2004); Yildirim and Ljubomirov, 2007:116 (Turkey: Erzurum: Oltu; Tortu; İçel: Aydıncık); Danilov, 2008:348 (Russia: Altayskiy Kray: Barnaul area, as *kirbyi*); Dollfuss, 2008b:1409 (locality records from 30 African and European countries); Gayubo, González, Tormos, and Asís, 2008:136 (Spain: Salamanca: Parque Natural de Las Batuecas – Sierra de Francia); Kazenas, 2008c:255 (Kazakhstan: village Koktum south of Lake Alakol', as *kirbyi*); Ljubomirov and Yildirim, 2008:27 (in catalog of Sphecidae of Turkey); Zettel, Wiesbauer, and Zimmermann, 2008:134 (Austria: Burgenland: several localities; Niederösterreich: several localities; Wien,); Baños-Picón, Asís, Gayubo, and Tormos, 2009:310 (Spain: frequency of specimens collected with hand nets and Malaise traps); Danilov, 2009:54 (Russia: Western Siberia: Kulundinskaya Steppe); Gayubo, González, Tormos, and Asís, 2009:362 (Spain: Valladolid: Reserva Natural Riberas de Castronuño – Vega del Duero); González, Gayubo, Asís, and Tormos, 2009:622 (Spain: Salamanca and Zamora provinces: Arribes del Duero Natural Park); Józán, 2009:165 (Croatia: Bibinje, as *kirbyi*); Shorenko, 2009:366 (in list of Sphecidae *sensu lato* of Crimea); Bitsch, 2010:105 (in supplement to vol. II of Faune de France, 1997: recent biological and distributional records reported); Burguet *in* Durand and Burguet, 2010:22 (France Département d'Ardèche: Ardèche Gorges); Danilov, 2010b:44 (distribution of Palearctic-Ethiopian type); Shorenko and Konovalov, 2010:12 (Ukraine); Smetana, Roller, Beneš, Bogusch, Dvořák, Holý, Karas, Macek, Straka, Šima, Tyrner, Vepřek, and Zeman, 2010:87 (Western Slovakia: Borská Nížina), 100 (Western Slovakia: recorded from Borská Nížina); van der Smissen, 2010b:386 (France: Ardèche, Gard, and Vaucluse: locality records); Baghirov, 2011b:140 (Russia: Altayskiy Kray: Barnaul); Cruz-Sánchez, Asís, Gayubo, Tormos, and González, 2011:497 (Spain: Salamanca and Zamora provinces: Arribes del Duero Natural Park: effects of wildfire); Field, Ohl, and Kennedy, 2011:734 (in molecular cladistic analysis of Ammophilini); Gogala, 2011:6 (Slovenia); Józán, 2011:179 (in checklist of Sphecidae *sensu lato* of Hungary); Murai and Amr, 2011:120 (recorded from Syria by Coulon, 1925 and Dollfuss, 2008b); Schmid-Egger, 2011b:603 (United Arab Emirates: Wadi Wurayah), 604 (color photograph of male); Wiesbauer and Zettel, 2011:269 (Austria: Niederösterreich: March = Morava lowland); Burguet *in* Durand and Burguet, 2012:2 (France: Puy-de-Dôme: sands Girauds-Faures in Orléat commune); Danilov, 2012a:164 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:63; Japoshvili and Ljubomirov, 2012:96: Turkey: Isparta: Gölcük Nature Park 8 km SW city of Isparta, as *kiirbii kirbii*); Prisniy, 2012:46 (Russia: Belgorod Oblast'); Protsenko, Fateryga, Ivanov, and Puzanov, 2012:58 (Ukraine: Crimea); Yildirim, 2012:74 (Turkey: Erzurum: Oltu); Strumia, Pagliano, and Gayubo, 2012:55 (Italy: Toscana: Riserva di San Rossore); Bayındır, Gürbüz, Ljubomirov, and Pohl, 2013:146 (Turkey: Isparta: Kasnak Oak Forest Nature Reserve); Kazenas, 2013a:18, 19 (color photographs of

adult wasps, short information on geographic distribution and nesting habits, as *kirbyi*); Shlyakhtenok, 2013:130 (in annotated catalog of aculeate wasps of Belarus'); Vieira, Oliveira, Brewster, and Gayubo, 2013:15 (Portugal: Douro International Natural Park); Baldock, 2014:354 (Spain: Island of Mallorca: four localities); Bitsch, 2014:396 (France: Aisne: Grisolles); Danilov, 2014a:424 (Russia: Siberia: Altayskiy Kray), 2014b:516 (in key to Sphecidae s.s. of Siberia); Kazenas, 2014a:131 (Kazakhstan: Karatau Mountain Range); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces, as *kirbii kirbii*); Zettel, Ockermüller, and Wiesbauer, 2014:162 (Austria: Wien-Lobau); Augul, Abdul-Rassoul, and Kaddou, 2015:115, 116 (in key to Sphecini of Iraq), 116 (Iraq: locality records), 117, 118 (illustrations); Gülmez and Can, 2015:12 (usability of ITS2 and 28S-D2 gene regions for species identification, as *kirbyi*); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey); Lett, 2015:5 (France: Loir-et-Cher: Gièvre); P. Rosa and Pagliano, 2015:96 (Italy: Lombardia and Piemonte: Parco del Ticino); Shorenko, 2015:317 (in list of Sphecidae *sensu lato* of Crimea); Simon, Lorthiois, and Macé, 2015:54 (France: alluvial terraces in Eure Department: Courcelles-sur-Seine and Seine-Maritime Department: Anneville-Ambourville); Gülmez and Dizer, 2016:58 (Turkey: Tokat Province); Levárdá and Matache, 2016:42 (in catalog of Sphecidae s.s. of Romania); Mokrousov and Popov, 2016:564 (Russia: Abkhazia, Krasnodarskiy Kray); Arens, 2017a:631 (Greece: Peloponnesus); Burguet and Durand, 2017:9 (France: Puy-de-Dôme: basin of river Crouël); Danilov, 2017b:215 (in catalog of Sphecidae s.s. of Russia); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:25 (Iran: known from Ardabil and Golestan provinces); Danilov and Mokrousov, 2017a:109 (Russia: Dagestan, Kalmykia); Magdalou, 2017:16 (France: Pyrénées-Orientales: Réserves Naturelles Catalanes: Forêt de la Massane, Mas Larrieu); Shorenko, 2017:76 (in Crimea collected in June through August); Lett, 2018:202 (France: Loir-et-Cher: Gièvres at 47°16'39"N 1°41'52"E); Shorenko, 2018:127 (Crimea, including localities, habitats, and number of specimens); Augul, 2019:497 (Iraq: Baghdad, Karbala, and Wasit provinces); Can and Gülmez, 2019:349 (in key to *Prionyx* of Turkey); Carminati, Mora, and Cretin, 2019:29 (France: in list of Sphecidae *sensu lato* of Franche-Comté); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Prionyx* s.s.); Gülmez, 2019:3 (Turkey: Amasya, Ankara, Sivas, and Tokat provinces: no specific localities), 4 (abnormal forewing venation); Ben Khedher, Yildirim, Braham, and Ljubomirov, 2020a:314 (in list of Tunisian Sphecidae *sensu stricto*; additional records: Tunisia: Beja, Jendouba, Kairouan, Kasserine, Kebili, Le Kef, Mahdia, Nabeul, Sidi Bouzid, Sousse, Tataouine, Tozeur, and Zaghouan provinces); Bitsch, Barbier, and Jacobs *in* Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:138 (in Sphecid Fauna of Europe); Cassar and Mifsud, 2020:164 (in checklist of Sphecidae s.s. of Malta); Danilov, 2020:320 (specimens from Kazakhstan, Kyrgyzstan, and Russia: Altay Kray, Astrakhan' Oblast', Dagestan in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Field, Gonzalez-Voyer, and Boulton, 2020:7 (evolution of parental care); Gadallah, 2020d:85 (in list of aculeate wasps of Arabian Peninsula); Maharramov, Mokrousov, and Proshchalykin, 2020:46 (Azarbaijan: Nakichivan Autonomous Republic); Örgel, Anlaş, and Tezcan, 2020:638 (Turkey: Manisa Province); Verheyde, Waanders, and Theite, 2020:165 (first record from Belgium: West-Vlaanderen Province: Ieper, and Netherlands: Limburg Province: Brunssum); Can and Gülmez, 2021a:3012 (Turkey: Giresun and Sivas provinces); Cross, Baldock, and Wood, 2021:18 (in catalog of Sphecidae *sensu lato* of Portugal); Nix, Staudt, and Trifonov, 2021:42 (first record from Germany: six localities in southwestern Germany); Can, 2022b:1875 (parasite: mite *Leptus* sp., Trombidiformes, Erythraeoidea); N. Schneider and Cungs, 2022:19 (first record from Luxemburg); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

Sphex albisectus Lepeletier de Saint Fargeau and Serville, 1828:462, ♀, ♂. Lectotype: ♀, Italy: Piemonte: no specific locality (MNHN), designated by Menke *in* R. Bohart and Menke, 1976:133. Synonymized with *Parasphex kirbii* by F. Smith, 1856:267 (but used as valid name). – Ghiliani, 1842:24 (listed from Italy: Sicilia, as *albiset*); Lepeletier de Saint Fargeau, 1845:358 (in revision of world Hymenoptera); Lucas, 1849:272 (Algeria: Oran); Fabre, 1856a:149 (nesting habits); Girard, 1879:964 (color, distribution, and habits); Becker, 1880:153 (Russia: Sarepta, now Krasnoarmeysk south of Volgograd); Kohl 1885b:185 (in revision of Palearctic *Sphex*); Gasperini, 1887:18 (recorded from Dalmatia, now Croatia, by Kohl, 1885b); Ed. André, 1888:130 (in revision of Sphecidae of Europe and Algeria), 10* (bibliographic references); Kohl, 1888b:730 (Austria: Tirol, now Italy: Alto Adige); Radoszkowski, 1888a:329 (genitalia, as *albisecta*); Gasperini, 1889:70 (Dalmatia: Lesina, now Croatia: Hvar); Cameron, 1889c:106 (in list of Sphecidae of Oriental Region); Kohl and

Handlirsch, 1889:275 (Turkmenistan: Chuli); Kohl, 1890b:335 (in revision of world Sphecini); F. Morawitz, 1891a:202 (Russia: Astrakhan Government); Kohl, 1894:342 (Congo: Gabun, Mozambique: Delagoa Bay, now Maputo Bay, Tanzania: Island of Zanzibar, East Africa (country unknown): Chama); Sickmann, 1894:216 (China: Hopei Province: Tientsin); De Stefani Perez, 1895:226 (in catalog of Sicilian Hymenoptera); Laboulbène, 1875:179 (reference to Fabre's observation on prey); Medina, 1894a:260 (Spain: Sevilla and Pozuelo de Calatrava); Schletterer, 1894:34 (Istria Peninsula, now part of Croatia, Slovenia, and Italy); Dalla Torre, 1897:414 (in catalog of world Hymenoptera); Mocsáry, 1897:79 (Kingdom of Hungary, some localities are in today's Croatia and Romania); Ferton, 1901a:144 (fly parasite), 1902:512 (nesting habits); Adlerz, 1904:139 (known prey: acridids); Antiga and Bofill, 1904:5 (Spain: Cataluña Province); E. Saunders, 1904c:636 (Spain); W. Schulz, 1904b:93 (Spain: Granada; Lebanon: Beirut); Mantero, 1905:69 (Italy: Toscana: Isola del Giglio); W. Schulz, 1905b:9 (West Africa: Muculla), 34 (Algeria); Vángel, 1905:166 (Hungary); Dusmet and Mercet, 1906:507, 515 (in key to Spanish Sphecini); Graeffe, 1906:456 (Tunisia: Hammam el Lif); Magretti, 1906:12 (Eritrea, as *albisecta*); W. Schulz, 1906:193 (correct authorship); Móczár and Henter, 1907:205 (Hungary: Tiszaalpar); Schmiedeknecht, 1907:244 (in key to Hymenoptera of Central Europe); Cameron, 1908a:263 (Tanzania: Kilimanjaro and Meru); de Gaulle, 1908:104 (in catalog of French Hymenoptera); Cameron, 1910b:138 (South Africa: Transvaal: Kranspoort); Brauns, 1911a:118 (South Africa); Graeffe, 1911:49 (Italy: Trieste area); Morice, 1911:75 (Algeria: Biskra, Bône, now Annaba); Mantero, 1911:72 (Italy: Sardegna: Isola dell'Asinara); W. Schulz, 1911b:164 (Gribodo's 1884 determination confirmed); Maidl, 1913:560 (Egypt: Birket Karoun); Smits van Burgst, 1913a:319 and 1913b:6 (Tunisia: no specific locality); Dusmet y Alonso, 1915:86 (Spain: Aragón); Mantero, 1915:325 (Libya); Maidl, 1922:67 (Croatia); Ferton, 1923:86 (preying on acridids, like *Tachysphex panzeri*); Berland, 1924:89 (France: Var: Callian, stridulation during nest excavation), 1925b:45 (nesting habits), 1925d:37 (in Sphecidae Fauna of France); Coulon, 1925:116 (France, Syria, Morocco: Tanger); Roth, 1925:383 (in revision of North African Sphecini); Zanon, 1926:90 (Libya: Fueihat 15 km south of Benghazi); Berland, 1926a:45 (prey, homing), 1926b:168 (miscellaneous locality records); von Schulthess, 1926b:209 (Tunisia, Libya); Dusmet y Alonso, 1927:25 (Spain: Cataluña: Tarragona); Berland, 1928b:175 (add Roth, 1925 to bibliography in Berland, 1925d); Grandi, 1928a:12 (nesting habits); Guiglia, 1929:395 (Libya: Cirenaica: Cirene); Kruger, 1929a:21 and 1929b:56 (Libya: Cyrenaica: Giarabub); Bischoff, 1930a:216 (Tajikistan: Pamir); Schmiedeknecht, 1930:705 (in keys to Hymenoptera of North and Central Europe); Schouteden, 1930:95 (Zaire); Giordani Soika, 1931b:23 (Italy: lido di Venezia, unusually small specimen); Berland, 1932a:22 (France: Var Department; utilizing old charcoal preparation sites for nesting); Giordani Soika, 1932a:20 (Italy: Lido di Venezia); Bischoff, 1933:5 (Morocco); Nadig, 1933:103 (Morocco); Bernard, 1934b:249 (prey: *Stauroderus vagans* Eversmann); Giner Marí, 1934:130 (Spain); Grandi, 1934:63 (nesting habits), 130 (Italy: Emilia-Romagna: Cervia, and Lazio: Acilia); Guiglia, 1934b:294 (Libya: bibliography and summary of locality records); Maidl, 1934:64 (Greece: Aegean Islands: Milos and Seriphos); Bernard, 1935:61 (France: Var: Fréjus area; prey: *Stauroderus vagans* Eversmann); Vergne, 1935:117 (France: Auvergne: Les Martres-d'Artières); Zavadil, 1937a:73 (eastern Slovakia); Zavadil, Šusterka, and Bat'a, 1937:212 (in catalog of Sphecidae of Czechoslovakia); Móczár, 1938d:80 (Hungary: Pótharasz puszta); Yasumatsu, 1938c:83 (China: Manchuria; in revision of East Asian Sphecini); Móczár, 1939:5 (Hungary: Jászberény); Deleurance, 1941:278 (France: Camargue; prey: *Dociostaurus genei* (Ocskay)); Yasumatsu, 1942c:106 (China: Beijing; Nei Mongol: Apaka at 44°N 114.96°E); Giner Marí, 1943a:81 (in Sphecidae Fauna of Spain); Guiglia, 1943c:76 (Ethiopia: Gamo Gofa: Sagan-Omo region, as *albisectum*); Timon-David, 1943:29 (France: Bouches-du-Rhône: plage de Fos); Guiglia, 1944b:7 (Italy); Honoré, 1944a:63 (in revision of Egyptian Sphecini); Giner Marí, 1945b:359 (eastern Morocco: Muley Rechid); Deleurance, 1946b:62 (list of prey), 67 (France: Bouche-du-Rhône: Camargue: Bois des Rièges); Chaudoir, 1947:142 (France: Gard: Roquemaure); Dulac, 1947:53 (France: Saône-et-Loire Department: Creusot area); Soyer, 1947:117 (details of nest closure); Zavadil in Zavadil and Šnoflák, 1948:167 (in key to Sphecidae of Czechoslovakia); de Andrade, 1949:9 (Portugal); Berland and Bernard, 1949:2 (in revision of French *Sphex* s. l.), 7 (review of biological data); de Beaumont, 1950f:396 (Algeria); Guiglia, 1950:248 (Ethiopia: Gamo Gofa: Caschei, as *albisectum*); Scobiola, 1950:21 (Romania); de Beaumont, 1951e:267 (Morocco); Cleu, 1953:50 (France: Ardèche River basin); Móczár, 1953:309 (Hungary); Nouvel and Ribaut, 1953:177 (France: Haute-Garonne: Saint-Béat); Grandi, 1954:157 (nest and prey: *Chortippus bicolor* Charp.), 236 (Italy); Hertzog, 1954:99 (France: Bouches-du-Rhône: Camargue); de Beaumont and Bytinski-Salz, 1955:41 (Israel); Leclercq, 1955h:31 (bibliographic references, faunal records from Africa, nomenclatural history, variation); Steiner, 1955:133 (France: Dordogne); Vergne,

1955:4 (France: Auvergne); Vogrin, 1955:31 (Yugoslavia); Berland, 1956:1168 (in revision of African Sphecini); Bytinski-Salz, 1956:224 (Turkey: Bursa, Elmalı, İzmir, Karapınar, Kemer); Ceballos, 1956:362 (in catalog of Hymenoptera of Spain); de Beaumont, 1956a:181 (Libya); Dulac, 1956:9 (France: Isère: Grenoble, Saône-et-Loire: St. Laurent-d'Andenay E Creusot); Hertzog, 1956:157 (prey: acridids *Omocestus ventralis* (Zetterstedt), *Euchortippus pulvinatus* (Fischer Waldheim), *Aiolopus thalassinus* (Fabricius), *Oedipoda coerulescens* (Linnaeus), prey transport, nest closure, nest structure); Morel, Nouvel, and Ribaut, 1956:337 (France: Département des Pyrénées-Orientales); Bajári, 1957a:8, 10 (in key to Hungarian Sphecidae); de Beaumont, 1957b:130 (northern Iran); Guiglia, 1957:144 (Italy: Isole Pelagie: Lampedusa); Balthasar, 1958:339 (Slovakia: Čenkov); Nouvel and Ribaut, 1958:8 (France: Pyrénées-Orientales: Banyuls-sur-Mer area); Pulawski, 1958a:164 (Bulgaria: Aitos, Sozopol, Varna); Benz, 1959 (nesting habits); de Beaumont, 1959a:10 (Italy); Diniz, 1959:27 (Portugal: nine localities); Scobiola-Palade, 1959:497 (Romania: Constanța Region: Agigea); Suárez, 1959:53 (Spain: Almería Province); de Beaumont, 1960a:5 (Greece: Island of Rhodes), 1960b:227 (Libya); Guiglia, 1960:360 (Italy: Isole Pelagie: Lampedusa); Scobiola-Palade, 1960b:232 (Romania: several localities, male genitalia illustrated); Wenger, 1960:421 (nesting habits); Grandi, 1961:146 (nesting habits); Leclercq, 1961b:47 (Zaire); Atanassov, 1962:125 (Bulgaria: Petrich area); de Beaumont, 1962b:19 (Spain); Lehrer and Scutaru, 1963:287 (Romania: Iași); Scobiola, 1963:825 (Romania: Periprava); Tsuneki, 1963b:48 (nesting habits); Ceballos, 1964:87 (in supplement to catalog of Spanish Sphecidae); de Beaumont, 1964c:28 (in Sphecid Fauna of Switzerland); Diniz, 1964c:100 (in key to Angolan *Sphe*x); de Beaumont, 1965a:13 (Greece); Isensee, Lesemann, and Röseler, 1965:612 (Spain: Gerona: no specific locality); Balthasar, Hrubant, and Hrubant, 1967:175 (Bulgaria: Slanchev Bryag near Nessebar); Carayon, 1967:744, 748 (France: Vaucluse Department: nocturnal rest site); de Beaumont, 1967a:273 (Turkey); Scobiola-Palade, 1968b:141 (Romania: Island of Letea in delta of Danube), 1968c:382 (Romania: Budești, Copăceni); Benedek, 1969a:83 (Hungary: marshy meadow); Kazenas, 1969a:21 (Kazakhstan: Ili River, foothills of Dzhungarian Alatau, Arkharly Range, Golodnaya Step, Uil River, Gur'yev, now Atyraū); Tsuneki, 1971m:2 (China: Peking: Tiendang); Balthasar, 1972:424 (in Sphecid Fauna of Czechoslovakia); Kazenas, 1972b:113 (Kazakhstan); Myartseva, 1972a:83 (Turkmenistan); Scobiola-Palade, 1972a:148 (Romania: delta of Danube: Caraorman); Simon Thomas, 1972:175 (France: Gironde, Landes, Lot-et-Garonne, Périgord Noir; Haute Garonne: Saint-Béat); Erlandsson, 1974:58 (France, Italy, Spain); Kazenas, 1974b:109 (feeding on flowers of *Tamarix* sp. and *Statice gmelini* Willd., Plumbaginaceae, in Kazakhstan), 112 (feeding on flowers of *Apocynum lancifolium* Russ., Apocynaceae, in Kazakhstan); Simon Thomas, 1976:3 (France: Lot-et-Garonne: Forêt de Campet); Marion, 1978:86 (France); Radović and Krunic, 1979:unpaginated foldout (nesting in sand, foreleg structure); Abdu and Shaumar, 1985:229 (Qatar: Doha, Karaanah); Benz, 1985:1228 (nesting habits summary); Meyer-Holzappel, 1986:100 and 103 (nest parasite: *Hilarella stictica* Meigen, Sarcophagidae); Piek and Spanjer, 1986:189 (in list of Sphecidae with known prey, as *albisecta*); Bonelli, 1988:87 (prey and nest); Pádr in Šedivý, 1989a:166 (in checklist of Czechoslovakian Sphecidae); Delarze, 1992:68 (Switzerland: Valais: Les Follatères); Alieva and Humbatov, 2007:77 (nesting and prey, from literature); Pagliano, 2008:532 (specimens in M. Spinola collection, Torino). – **As *Sphe*x *albicinctus*** [sic]: Guérin-Méneville, 1829a:562 (in Dictionnaire classique d'histoire naturelle); Scobiola-Palade, 1985:95 (new generic assignment, Romania: delta of Danube). – **As *Enodia albisecta***: Dahlbom, 1843:28 (new generic assignment, in revision of Sphecidae and Pompilidae), 1845:438 (in key to world *Enodia*); A. Costa, 1858b:12 (in revision of Sphecidae of Kingdom of Naples), 1867b:71 and 1867c:15 (in revision of Italian Sphecidae); Palma, 1869:38 (Italy: Sicilia settentrionale); von Aichinger, 1870:322 (Austria: Tirol); Martinez y Saez, 1874:31 (specimens from Spain: Madrid area, donated to Leon Dufour, as *albisectus* Bonelli); Radoszkowski, 1881:210 (Angola); A. Costa, 1882b:22 (Italy: Sardegna), 1883:57 (Italy: Sardegna: island of Asinara and Terranova), 1884b:323 (Italy: Sardegna: Cagliari); Radoszkowski, 1887b:91 (in list of Transcaspiian Hymenoptera), 1892:586 (description of male genitalia); Roth, 1924:123 (Algeria: Nemours, now Ghazaouet); Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino). – **As *Parasphe*x *albisecta***: F. Smith, 1856:267 (new generic assignment, in catalog of Hymenoptera in British Museum); Kirchner, 1867:217 (in catalog of European Hymenoptera); Dours, 1874:146 (in catalog of Hymenoptera of France); Mocsáry, 1874:120 (Siebenbürgen, now Romania: Transylvania: no specific locality); Marquet, 1875:207 (France: Haute-Garonne: Toulouse; Hérault: Cette, now Sète); Frivaldszky, 1876:354 (Hungary: Temes Komitat: Grebenács, now in Timiș District in Romania); Becker, 1880:153 (Russia: Sarepta, now Krasnoarmeysk south of Volgograd); Kohl, 1880:182 (Italy: Trentino–Alto Adige); Marquet, 1881:178 (southern France); Sajó, 1882:5 (Hungary); Kohl, 1883e:674 (Switzerland); Gribodo, 1884c:302 (Ethiopia:

Let Marefia 16 km north of Ankober); Cuní y Martorell, 1897:331 (Spain: Cataluña: villa de Calella). – **As Chlorion albisectum**: Arnold, 1928c:350 (new generic assignment, in revision of Sphecini of southern Africa), 1930:17 (in checklist of Afrotropical Sphecidae); Bischoff, 1931:9 (Spain). – **As Tachysphex [sic] albisectus**: Ferton, 1905:70 (position of egg on prey; generic name a lapsus). – **As Prionyx albisectus**: Diniz, 1965:3 (new generic assignment, Portugal: twenty four localities); Leclercq, 1969:1050 (Congo Brazzaville); Radović, 1985:64 (sting apparatus analyzed).

As *Sphex vetusta*: Pagliano, 2008:532 (specimens in M. Spinola collection, Torino).

ssp. marginatus (F. Smith)

Parasphex marginatus F. Smith, 1856:267, ♀, ♂ (as *marginata*, incorrect original termination). Syntypes: Gambia: no specific locality (BMNH). – **As Sphex marginatus**: W. Fox, 1896e:552 (new generic assignment, Ethiopia: Bale Province: Sheikh Husein, now Shek Husen); Kohl, 1890b:337 (in revision of world Sphecini); Dalla Torre, 1897:430 (in catalog of world Hymenoptera); Kohl, 1909:370 (Tanzania: Pemba Island); Strand, 1916b:104 (German East Africa, now Tanzania); G. Carpenter, 1930b:293 (nest closure). – **As Chlorion albisectum race marginatum**: Arnold, 1928c:351 (new status, in revision of Sphecini of southern Africa, West Africa, Africa from Zimbabwe to Cape Province), 1930:17 (in checklist of Afrotropical Sphecidae); Guiglia, 1939d:74 (Ethiopia: Sidamo: Neghelli, now Negele, as var. *marginatum*), 1940e:293 (Ethiopia: Harer: Gotqa, as var. *marginatum*). – **As Sphex albisectus marginatus**: Leclercq, 1955h:33 (new status, in key to subspecies of *Sphex albisectus*; locality records from various African countries), 1955i:406 (Burundi), 1962a:393 (Tanzania: Uluguru Mountains); Diniz, 1964c:102 (Angola: Lunda: Dundo); de Beaumont, 1976b:502 (South Africa: Cape Province). – **As Prionyx kirbii marginatus**: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Rodgers and Homewood, 1982:233 (Tanzania: Usambara Mountains); Guichard, 1988a:121 (Arabian Peninsula); Gadallah and Assery, 2004a:221 (in catalog of Sphecidae of Saudi Arabia); Schmid-Egger, 2011b:603 (is just a desert color form); Gadallah, Al Dhafer, Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia).

Sphex sjöstedti Cameron, 1908a:263, ♂ (as *Sjöstedti*, incorrect original capitalization and diacritic mark). Holotype or syntypes: ♂ Tanzania: Mount Meru lowlands (NRS). Synonymized with *Chlorion albisectum marginatum* by Arnold, 1928c:351. – **As Sphex albisectus var. sjöstedti**: Leclercq, 1961b:47 (new status, Zaire). – **As Sphex marginatus sjöstedti**: Diniz, 1964c:103 (new status, Angola: Lunda: Dundo, Matala). – **As Sphex albisectus marginatus sjöstedti**: Leclercq, 1955h:33 (new status, in key to subspecies of *Sphex albisectus*), 35 (locality records from Zaire and South Africa).

Sphex curvilineatus Cameron, 1912b:397, ♂. Holotype or syntypes: ♂, Zaire: Lukombe (MRAC). Synonymized with *Chlorion albisectum marginatum* by Arnold, 1928c:351. – Schouteden, 1930:95 (Zaire).

Sphex albisectus var. *alluaudi* Berland, 1926b:168, ♀, ♂. Lectotype: ♀, Ivory Coast: Assinie (MNHN), designated by Menke in R. Bohart and Menke, 1976:133. Synonymized with ... by ... – Diniz, 1964c:103 (Angola: Lunda: Dundo). – **As Chlorion albisectum var. Alluaudi**: Arnold, 1928c:352 (new generic assignment, new status, original description translated into English), 1930:17 (in checklist of Afrotropical Sphecidae); Leclercq, 1961b:47 (Zaire). – **As Sphex albisectus marginatus alluaudi**: Leclercq, 1955h:34 (new status, in key to subspecies of *Sphex albisectus*), 35 (Zaire: locality records).

Sphex albisectus var. *chudeaui* Berland, 1926b:168, ♀. Holotype: ♀, Senegal: Balé (MNHN). – **As Chlorion albisectum var. Chudeaui**: Arnold, 1928c:352 (new generic assignment, new status, original description translated into English), 1930:17 (in checklist of Afrotropical Sphecidae, as *chudeauxi*). – **As Sphex albisectus marginatus chudeaui**: Leclercq, 1955h:33 (new status, in key to subspecies of *Sphex albisectus*).

Sphex albisectus var. *congoensis* Berland, 1926b:169, ♀. Lectotype: ♀, Gabon: Libreville (MNHN), designated by Menke in R. Bohart and Menke, 1976:133. Synonymized with *Sphex albisectus marginatus sjöstedti* by Leclercq, 1955h:35. – **As Chlorion albisectum var. congoensis**: Arnold, 1928c:352 (new generic assignment, original description translated into English), 1930:17 (in checklist of Afrotropical Sphecidae).

24. kurdistanicus (Balthasar)

Sphex kurdistanicus Balthasar, 1954b:281, ♀, ♂. Holotype: ♀, Iraq: Kurdistan: Erbil (originally V. Balthasar coll., now NMPC). – **As Prionyx kurdistanicus**: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world

Sphecidae); Augul, 2019:497 (recorded from Iraq by Balthasar, 1954b); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*).

25. *lividocinctus* (A. Costa)

Enodia lividocincta A. Costa, 1861:39, ♀, ♂. Syntypes: Italy: Reggio di Calabria: Brancaleone; and Terra d'Otranto, now Apulia (Napoli) – A. Costa, 1863:65 (Italy: Calabria Ulteriore: Aspromonte), 1867b:71 and 1867c:15 (in revision of Italian Sphecidae); Palma, 1869:38 (Italy: Sicilia settentrionale); A. Costa, 1886b:21 (Italy: Sardegna); Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino); Pagliano, 2008:532 (specimens in M. Spinola collection, Torino). – *As Parasphex lividocincta*: Kirchner, 1867:217 (new generic assignment, in catalog of European Hymenoptera). – *As Sphex lividocinctus*: Kohl, 1885b:190 (new generic assignment, in revision of Palearctic *Sphex*); Kohl and Handlirsch, 1889:275 (Turkmenistan); Kohl, 1890b:339 (in revision of world Sphecini); A. Costa, 1893b:3 (Tunisia); Dalla Torre, 1897:429 (in catalog of world Hymenoptera); Antiga and Bofill, 1904:5 (Spain: Cataluña Province); Dusmet and Mercet, 1906:507, 514 (in key to Spanish Sphecini); Ferton, 1908:560 (France: Corse: Bonifacio; nest); Dusmet y Alonso, 1915:87 (Spain: Aragón); Berland, 1925d:38 (in Sphecid Fauna of France); Roth, 1925:384 (in revision of North African Sphecini); Nadig, 1933:103 (Morocco); Guiglia, 1934b:294 (Libya: bibliography and summary of locality records); Maidl, 1934:64 (Greece: Aegean Islands: Mytilene, determination tentative); Nadig, 1934:33 (Italy: Sardegna: Aritzo); Gussakovskij, 1935:412 (Tajikistan); Guiglia, 1938b:9 (Italy: Sardegna); Bernard, 1939:167 (France: Var: Fréjus); Giner Marí, 1943a:82 (in Sphecid Fauna of Spain); Honoré, 1944a:67 (in revision of Egyptian Sphecini); Chaudoir, 1947:142 (France: Gard: Roquemaure); de Beaumont, 1947b:382 (Cyprus); Guiglia, 1948c:200 (Italy: Sardegna: Villasalto); de Andrade, 1949:9 (Portugal); Berland and Bernard, 1949:2 (in revision of French *Sphex sensu lato*); Pittioni, 1950:20 (Cyprus); de Beaumont, 1951e:268 (Morocco); Cleu, 1953:50 (France: Ardèche River basin); de Beaumont, 1954e:86 (Italy); Harant and Leclercq, 1955:250 (France: Hérault: Maguelonne); Leclercq, 1955h:37 (bibliographic references, faunal records from Africa); Ceballos, 1956:363 (in catalog of Hymenoptera of Spain); de Beaumont, 1956a:181 (Libya); Morel, Nouvel, and Ribaut, 1956:337 (France: Département des Pyrénées-Orientales); de Beaumont, 1957b:130 (northern Iran); Nouvel and Ribaut, 1958:8 (France: Pyrénées-Orientales: Banyuls-sur-Mer area); Pulawski, 1958a:164 (Bulgaria: Sandanski); de Beaumont, 1959a:10 (Italy); Diniz, 1959:27 (Portugal: S. João do Estoril, Soure); Suárez, 1959:53 (Spain: Almería Province); de Beaumont, 1960a:5 (Greece: Island of Rhodes), 1960b:227 (Libya); Ceballos, 1964:87 (in supplement to catalog of Spanish Sphecidae); de Beaumont, 1965a:13 (Greece); Carayon, 1967:744, 748 (France: Vaucluse Department: nocturnal rest site); de Beaumont, 1967a:272 (Turkey), 1969:81 (Turkey); Kazenas, 1969a:21 (Kazakhstan: Ili River, foothills of Dzhungarian Alatau, Kyzylagash, Golodnaya Step'); Islamov, 1970:63 (Uzbekistan: Chirchik Basin); Balthasar, 1972:425 (in Sphecid Fauna of Czechoslovakia: may be expected in the country); Kazenas, 1972b:113 (Kazakhstan); Erlandsson, 1974:58 (Greece); Georgiou, 1977:192 (Cyprus); Kazenas, 1978b:44 (in key to Sphecidae of Kazakhstan and Central Asia); Pulawski, 1978:183 (in key to Sphecidae of European part of USSR). – *As Prionyx lividocinctus*: Diniz, 1965:4 (new generic assignment, Portugal: Estoril); Myartseva, 1972a:84 (Turkmenistan); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Pagliano, 1980:110 (Italy: Lazio); Gayubo, 1982f:245 (Spain: Cádiz Province: Puerto de Santa Maria); Radović Krunić, and Brajković, 1982:28 (Yugoslavia); Gayubo, 1983c:231 (Spain: Salamanca Province: Cerralbo); Mingo and Gayubo, 1983:154 (Spain); Schmidt and Westrich, 1983:121 (Greece); Gayubo, 1984c:357 (Portugal: El Algarve Province); Gayubo and Tormos, 1984:9 (Spain: Valencia); Pagliano, 1984:367 (Italy); Chevin and Chevin, 1985:38 (France: Aude); Pagliano, 1985:8 (Italy); Radović, 1985:64 (sting apparatus analyzed); Gayubo, 1986b:36 (Spain: Andalucía); Islamov, 1986:516 (Uzbekistan: Surkhandarya and Tashkent Oblasts); Gayubo, 1987:107 (Spain: Ciudad Real Province); Pagliano, 1990:58 (in catalog of Italian Sphecidae); Hamon, Fonfria, and Tussac, 1991:128 (in key to French Sphecini), 133 (in France along Mediterranean shore); Kazenas and Nasyrova, 1991:38 (Kazakhstan: preying on *Dociostaurus tartarus* (Stshelk.)); Schembri, 1991:176 (first record from Malta); Gayubo, Borsato, and Osella, 1992:277 (Greece); Ebrahimi, 1993:96 (Iran); Gayubo and Borsato, 1994:200 (Italy: Toscana, Sardegna), 201 (map of distribution in Italy); Negrisolo in Minelli, Ruffo, and La Posta, 1995b:2 (in catalog of Italian fauna); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:61 (in Sphecid Fauna of Western Europe); Kazenas,

1998b:113 (in Sphecidae Fauna of Kazakhstan); Nazarova, 1998:40 (Tajikistan: Tigrovaya Balka Nature Reserve); Esenbekova and Kazenas, 2000:9 (southeast Kazakhstan: near lake Biylikul', 45 km NW Suzak); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:28 (geographic distribution, collecting localities in Kazakhstan); Gayubo, Nieves-Aldrey, González, Tormos, Rey del Castillo, and Asís, 2004:108 (Spain: Madrid: Monte de El Pardo); Kazenas, 2004b:99 (Kazakhstan: western Tien Shan Mountains), 2004d:26 (Kazakhstan: northern Caspian region); Gayubo and Özbek, 2005:8 (Turkey: Antalya: Belkız; Ýcel: Silifke); Gülmez and Tüzün, 2005:47 (Turkey: Ankara Province); Pagliano and Negrisol, 2005:54 (in Sphecidae Fauna of Italy); Blösch, 2006:63 (specimens spend nights attached to plant stems with their heads down); Standfuss and Standfuss, 2006c:307 (Greece: Thessalia: Magnisia Peninsula at 39°N 23°E); Roche, 2007a:42 (in checklist of Egyptian Sphecidae, redescription, as *lividocinctus lividocinctus*), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Dollfuss, 2008b:1412 (locality records from Bulgaria, Greece, Italy, Kazakhstan, Kyrgyzstan, Morocco, Turkey, and Turkmenistan); Kazenas, 2008c:255 (Kazakhstan: village Koktum south of Lake Alakol'); Ljubomirov and Yildirim, 2008:27 (in catalog of Sphecidae of Turkey); Baños-Picón, Asís, Gayubo, and Tormos, 2009:310 (Spain: frequency of specimens collected with hand nets and Malaise traps); Danilov, 2009:54 (Russia: Western Siberia: Kulundinskaya Steppe); Pagliano, 2009:175 (Italy: Molise: Mafalda); Bitsch, 2010:105 (in supplement to vol. II of Faune de France, 1997: France: first records from the Départements of Alpes-de-Haute-Provence: Ganagobie monastère, Sainte-Croix-de-Verdon, and Landes: Saint Geours-de-Marenne); Danilov, 2010b:44 (distribution of Tethyan type), 2012a:164 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:63 (bibliographic references, geographic distribution); Kazenas, 2013a:19, 20 (color photograph of female, short information on geographic distribution and nesting habits); Baldock, 2014:354 (Spain: Island of Mallorca: Campos); Danilov, 2014b:515 (in key to Sphecidae s.s. of Siberia, not yet found in Siberia); Kazenas, 2014a:131 (Kazakhstan: Karatau Mountain Range); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces, as *lividocinctus lividocinctus*); Augul, Abdul-Rassoul, and Kaddou, 2015:116 (in key to Sphecini of Iraq), 117 (locality records); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey); Mokrousov and Popov, 2016:564 (Russia: Krasnodarskiy Krai); Arens, 2017a:631 (Greece: Peloponnesus); Danilov, 2017b:215 (in catalog of Sphecidae s.s. of Russia); Danilov and Mokrousov, 2017a:109 (Russia: Crimea, Dagestan, Kalmykia); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:25 (Iran: known from Ardabil, Golestan, Khuzestan, and Qazvin provinces); Augul, 2019:497 (recorded from Iraq by Kaddou, 1967); Can and Gülmez, 2019:349 (in key to *Prionyx* of Turkey); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Prionyx* s.s.); Gülmez, 2019:3 (Turkey: Ankara Province: no specific locality); Ben Khedher, Yildirim, Braham, and Ljubomirov, 2020a:315 (in list of Tunisian Sphecidae *sensu stricto*); Bitsch, Barbier, and Jacobs in Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:139 (in Sphecidae Fauna of Europe); Cassar and Mifsud, 2020:164 (in checklist of Sphecidae s.s. of Malta); Danilov, 2020:320 (specimens from Kyrgyzstan and Russia: Astrakhan' Oblast', Crimea, Dagestan, and Kalmykia in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Tursi, Altadonna, Lo Cascio, Nobile, and Selis, 2020:728 (Italy: Aeolian Archipelago: island of Lipari); Can and Gülmez, 2021a:3012 (Turkey: Gümüşhane: Şiran: Güreşköy); Cross, Baldock, and Wood, 2021:18 (in catalog of Sphecidae *sensu lato* of Portugal); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*); Embergenov, Akhmedov, and Medetov, 2023:24 (Uzbekistan: southern Lake Aral region); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

Priononyx isselii Gribodo, 1880b:401, ♀ (as *Isselii*, incorrect original capitalization). Holotype: ♀, Tunisia: Galite Island (MSNG). Synonymized with *Sphex lividocinctus* by Kohl, 1885b:190 and 1885c:165, synonymy confirmed by de Beaumont, 1950e:262. – Penati and Mariotti, 2015: 70 (in list of Hymenoptera described by G. Gribodo).

Enodia obliquestriata Mocsáry, 1883:37, ♀ (as *oblique-striata*, incorrect original hyphenation). Holotype or syntypes: ♀, Lebanon: Beirut (TMB). Synonymized with *Sphex lividocinctus* by Kohl, 1885b:190.

Enodia graeca Mocsáry, 1883:35, ♂ (as *Graeca*, incorrect original capitalization). Holotype or syntypes: ♂, Greece: Corfu Island (TMB). Synonymized with *Sphex lividocinctus* by Kohl, 1890b:339. – As *Sphex graecus*: Kohl, 1885b:189 (new generic assignment, in revision of Palearctic *Sphex*); Ed. André, 1888:134 (in revision of Sphecidae of Europe and Algeria), 9* (bibliographic references).

As *Sphex micans*: Ed. André, 1888:133 (in revision of Sphecidae of Europe and Algeria), corrected to *Sphex lividocinctus* by Kohl, 1889a:24.

ssp. apakensis (Tsuneki)

Sphex lividocinctus apakensis Tsuneki, 1971m:2, ♀. Holotype: ♀, China: Nei Mongol (= Inner Mongolia): Apaka at 44°N 114.96°E (originally K. Tsuneki coll., now USNM). – Nuhn and Menke, 1994:25 (holotype transferred to USNM). – As *Prionyx lividocinctus apakensis*: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Hua, 2006:276 (in list of Chinese insects, geographic distribution).

ssp. oasis (Tsuneki)

Sphex lividocinctus oasis Tsuneki, 1971k:143, ♀, ♂. Holotype: ♂, Mongolia: Bayanhongor Aymag: Tsagan Bogd ul (TMB). – As *Prionyx lividocinctus oasis*: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae).

26. macula (Fabricius)

Pepsis macula Fabricius, 1804:210, sex not stated. Holotype: ♂, Saudi Arabia: no specific locality (MNHN, coll. Bosc, now possibly lost: van der Vecht, 1961a:34). – As *Sphex macula*: Kohl in Dalla Torre, 1897:430 (new generic assignment, in catalog of world Hymenoptera); W. Schulz, 1911b:164 (probable syntype material in Jurine collection, MHNG); Berland, 1926c:201 (redescription of type); de Beaumont, 1962c:221 (Arabia: El Riyadh), 1970c:4 (Iran: Baluchistan). – As *Prionyx macula*: de Beaumont, 1961e:2 (new generic assignment, may be a senior synonym of *Prionyx eatoni* and *lugens*), 1968b:150 (member of *macula* species group). – As *Prionyx macula*: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Guichard, 1988a:120 (Arabian Peninsula); Roche and Zalut, 1994:113 (Egypt: Sinai Peninsula); Kazenas, 1998b:114 (in Sphecid Fauna of Kazakhstan), 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:29 (geographic distribution, collecting localities in Kazakhstan); Gadallah and Assery, 2004a:221 (in catalog of Sphecidae of Saudi Arabia); Roche, 2007a:44 (in checklist of Egyptian Sphecidae, redescription, as *macula macula*), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae, as *macula macula*); Danilov, 2010b:44 (distribution of Tethyan type); Al-Houty, 2011:89 (Kuwait: no specific locality); Danilov, 2012a:163 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:63 (bibliographic references, geographic distribution); Gadallah, Al Dhafer, Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia); Dunford, Turbyville, and Leavengood, 2014:11 (listed as medically important in Afghanistan); Ebrahimi, 2014:21 (Iran: Bushehr, Ilām, and Sīstān-Baluchestān provinces); Augul, Abdul-Rassoul, and Kaddou, 2015:116 (in key to Sphecini of Iraq; Iraq: locality records), 117, 118 (illustration); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:26 (Iran: known from Alborz, Bushehr, Ilam, and Sistan-o Baluchestan provinces); Augul, 2019:497 (recorded from Iraq by de Beaumont, 1961e); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Kazenas, 2019:85 (endangered in Uzbekistan); Ben Khedher, Yildirim, Braham, and Ljubomirov, 2020a:315 (in list of Tunisian Sphecidae *sensu stricto*; additional record: Tunisia: Kasserine: Sbeitla); Gadallah, 2020d:85 (in list of aculeate wasps of Arabian Peninsula); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).
Sphex eatoni E. Saunders, 1910:518, ♀, ♂. Syntypes: Algeria: Biskra (OXUM). Synonymized with *Sphex macula* by de Beaumont, 1962c:221. – R. Turner, 1912c:413 (British East Africa, now Kenya: Makindu); Roth, 1925:387 (in revision of North African Sphecini); Honoré, 1944a:70 (in revision of Egyptian Sphecini); de Beaumont and Bytinski-Salz, 1955:41 (Israel); Leclercq, 1955h:26 (bibliographic references); Grandi, 1959b:287 (Libya: Tripolitania: Jefren); de Beaumont, 1961e:2 (Iraq); Ebrahimi, 1993:94 (Iran).

As *Sphex lugens*: Grandi, 1935a:111 (Libya), corrected to *Sphex eatoni* by Grandi, 1959b:287.

ssp. lugens (Kohl)

Sphex lugens Kohl, 1889a:25, ♀, ♂. Syntypes: Armenia: Arax River Valley: no specific locality (NHMW). – Kohl, 1890b:348 (in revision of world Sphecini); Dalla Torre, 1897:430 (in catalog of world Hymenoptera); W. Schulz, 1911b:165 (as new synonym of *Sphex macula*); Gussakovskij, 1933b:278 (Iran); nec Grandi, 1935a:111 (= *Prionyx macula*); de Beaumont,

1960c:170 (Afghanistan; diagnostic characters, taxonomic problems). – **As *Sphex macula lugens***: de Beaumont, 1962c:221 (new status); Dollfuss, 1989:12 (type material in NHMW). – **As *Priononyx lugens***: de Beaumont, 1968b:150 (new generic assignment, member of *macula* species group). – **As *Prionyx macula lugens***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia).

27. *melanotus* (F. Morawitz)

Sphex melanotus F. Morawitz, 1890:575, ♀, ♂. Lectotype: Transcaspia: no specific locality (ZIN), designated by Danilov, 2012b:64. – Kohl, 1890b:346 (in revision of world Sphecini); Dalla Torre, 1897:432 (in catalog of world Hymenoptera); Berland, 1926c:200 (Turkestan, Altai). – **As *Priononyx melanotus***: de Beaumont, 1968b:149 (new generic assignment, member of *macula* species group). – **As *Prionyx melanotus***: Myartseva, 1972a:84 (new generic assignment, Turkmenistan); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Dollfuss, 1989:12 (type material in NHMW); Danilov, 2012a:163, 164 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:64 (bibliographic references, geographic distribution), 2016:341 (lectotype preserved in Zoological Institute, Sankt Petersburg, Russia); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

28. *neoxenus* (Kohl)

? *Sphex melaenus* Spinola, 1851a:398, ♀, ♂ (as *melaena*, incorrect original termination). Syntypes: Chile: central provinces: no specific locality (Torino). – F. Smith, 1856:260 (in catalog of Hymenoptera in British Museum); Kohl, 1890b:448 (original description copied); Dalla Torre, 1897:432 (in catalog of world Hymenoptera); R. Bohart and Menke, 1976:133 (as tentative senior synonym of *Prionyx neoxenus*). – **As *Priononyx melaenus***: Reed, 1894:626 (new generic assignment, revision). – **As *Chlorion melaenum***: Willink, 1951:203 (new generic assignment, original description translated into Spanish).

Sphex neoxenus Kohl, 1890b:363, ♀. Holotype: ♀, Canada: British Columbia: Vancouver Island, but actually South America (NHMW) – Fernald, 1906:418 (probably a South American species); Brèthes, 1908:146 (in revision of *Sphex thomae* group; Argentina); Jörgensen, 1912:286 (Argentina: Mendoza Province). – **As *Priononyx neoxenus***: Schrottky, 1913a:225 (new generic assignment, Argentina). – **As *Chlorion neoxenum***: Willink, 1948a:319 (new generic assignment, in key to *Chlorion thomae* species group), 1951:177 (in revision of Argentinean Sphecini). – **As *Prionyx neoxenus***: Pérez d'Angello, 1974:145 (new generic assignment, Chile; color variation); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Sielfeld, 1980b:72 (in checklist of Chilean Sphecidae); Pérez d'Angello, 1989:263 (Chile: Magellan Region); Amarante, 2002:72 (in catalog of Neotropical Sphecidae); Dollfuss, 2008b:1413 (locality records from Chile); Rasmussen and Asenjo, 2009:16 (in checklist of Crabronidae of Peru); Chiappa, 2012:8 (Chile: Región de Valparaíso: no specific locality); Danilov, 2019:78 (member of subgenus *Priononyx*); Danilov and Odintsev, 2023:433 (in subgenus *Priononyx*).

Sphex omissus Kohl, 1890b:364, ♂. Syntypes: ♂, Chile: Valparaíso (NHMW). Synonymized with *Chlorion neoxenum* by Fernald, 1931a:441. – Dalla Torre, 1897:435 (in catalog of world Hymenoptera, as *omissus*); Strand, 1910b:15 (Peru, actually Bolivia: Guaqui), 1916b:100 (description of ♀); Brèthes, 1918a:124 (Peru: Arequipa); Ruiz Pereira, 1924:102 (Chile: Cerro San Cristóbal); C. Reed, 1928:317 (resembling in coloration *Pompilus diaphanicus* Spinola and the Diptera *Laphria rufiventris* Blanch. and *Acylaticus pictus* Phil.); Gazulla and Ruiz Pereira, 1929:299 (Chile: Hacienda de "Las Mercedes", as *ommissus*); Ruiz Pereira, 1934:167 (Chile: Pahuano; as *ommissus*), 1937:164 (Chile: Coquimbo Province, as *ommissus*); Dollfuss, 1989:12 (type material in NHMW).

Sphex neoxenus var. *melanogaster* Brèthes, 1910a:261, ♀. Holotype: Argentina: Mendoza: Blanco Encalada (depository unknown). Synonymized with *Chlorion neoxenum* by Willink, 1951:177 and 179. – Jörgensen, 1912:286 (Argentina: Mendoza Province); Schrottky, 1913a:225 (Argentina: Mendoza); Liebermann, 1931:22 (in revision of Argentinian Sphecini); Genise, 1990:27 (depository of type material unknown).

Sphex gayi Berland, 1926c:203, ♀, ♂. Lectotype: ♂, Chile: no specific locality (MNHN), designated by Menke in R. Bohart and Menke, 1976:133. Synonymized with *Chlorion neoxenum* by Willink, 1951:177 and 180.

Sphex nigricapillus Berland, 1926c:205, ♀, ♂. Lectotype: ♂, Peru: Arequipa (MNHN), designated by Menke in R. Bohart and Menke, 1976:133. Synonymized with ..

29. *nigropectinatus* (Taschenberg)

Enodia nigropectinata Taschenberg, 1869:409, ♀ (as *nigro-pectinata*, incorrect original hyphenation). Syntypes: ♀, Sudan: Khartoum (Halle). – **As *Sphex nigropectinatus***: Kohl, 1885b:183 (new generic assignment, in revision of Palearctic *Sphex*); Ed. André, 1888:129 (in revision of Sphecidae of Europe and Algeria), 10* (bibliographic references); Kohl, 1890b:329 (in revision of world Sphecini.); Dalla Torre, 1897:434 (in catalog of world Hymenoptera); Bingham, 1898a:105 (Yemen: Aden); Morice, 1911:76 (Algeria: Biskra); Roth, 1925:381 (in revision of North African Sphecini.); Berland, 1926b:168 (miscellaneous locality records from Africa; prey: migratory locust); Kruger, 1929a:21 and 1929b:56 (Libya: Cyrenaica: Giarabub, as *nigripectinatus*); Guiglia, 1932d:472 (Libya: 85 km south of Gialo); Gussakovskij, 1933b:273 (Iran); Guiglia, 1934b:295 (Libya: bibliography and summary of locality records); Roth, 1934a:394 (Algeria: In-Salah); Giordani Soika, 1935:234 (Libya: Fezzan: Gat); Gussakovskij, 1935:413 (Tajikistan); Guiglia, 1936:4 (Libya: Gialo and Tazerbo in Cufra oasis), 1942b:230 (Libya); Honoré, 1944a:62 (in revision of Egyptian Sphecini.); Giner Marí, 1947:19 (Morocco: Western Sahara); Berland, 1950a:126 (Mauritania: Ford Gouraud, now Fdérík); de Beaumont and Bytinski-Salz, 1955:41 (Israel); Leclercq, 1955h:30 (bibliographic references, faunal records from Africa); Berland, 1956:1166 (in revision of African Sphecini.); Myartseva, 1963b:59 (Turkmenistan: lower Murgab River), 1965:83 (Turkmenistan: Akibay); Kazenas, 1969a:21 (Kazakhstan: southeast Kyzylkum Desert, Golodnaya Step', Ili River); Kazenas, 1978b:41 (in key to Sphecidae of Kazakhstan and Central Asia). – **As *Prionyx nigropectinatus***: Myartseva, 1971b:80 (new generic assignment, occurring only in sand deserts of southern Palearctic Region), 1972a:84 (Turkmenistan); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Guichard, 1980:224 (Oman), 1988a:121 (South Yemen); Ebrahimi, 1993:98 (Iran); Roche and Zalat, 1994:113 (Egypt: Sinai Peninsula); Kazenas, 1998b:116 (in Sphecid Fauna of Kazakhstan), 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 85 (list of prey), 2002a:29 (geographic distribution, collecting localities in Kazakhstan); Roche, 2007a:45 (in checklist of Egyptian Sphecidae), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Danilov, 2010b:44 (distribution of Palearctic-Ethiopian type), 2012a:163 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:64 (bibliographic references, geographic distribution), 2014b:515 (in key to Sphecidae s.s. of Siberia, not yet found in Siberia); Ebrahimi, 2014:21 (Iran: Sistān-Baluchestān: Zābol: Jalālābād); S. Gess and Roosenschoon, 2016:104 (Dubai Desert Conservation Reserve, visiting flowers of *Prosopis cineraria* (L.) Druce, Fabaceae); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:26 (Iran: known from Alborz, Sistan-o Baluchestan, and Tehran provinces); Madl, 2018:944 (in catalog of Hymenoptera of Djibouti); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Calosphex*); Daminova, 2019:86 (endangered in Uzbekistan); Danilov, 2020:319 (specimens from Kazakhstan in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Gadallah, 2020d:85 (in list of aculeate wasps of Arabian Peninsula); Danilov and Odintsev, 2023:432 (in subgenus *Calosphex*).

? *Sphex dives* Lepeletier de Saint Fargeau, 1845:359, sex not stated. Holotype: ♀, origin unknown (originally J. Serville coll., now ?). Tentatively synonymized with *Sphex nigropectinatus* by Kohl, 1890b:329. – F. Smith, 1856:243 (in catalog of Hymenoptera in British Museum); Dalla Torre, 1897:421 (in catalog of world Hymenoptera).

? *Harpactopus nivosus* F. Smith, 1856:265, ♀. Holotype or syntypes: ♀, northern India: no specific locality (BMNH). Tentatively synonymized with *Sphex nigropectinatus* by Kohl, 1890b:329. – F. Smith, 1871a:362 (in catalog of Oriental Aculeata). – **As *Sphex nivosus***: Cameron, 1889c:106 (new generic assignment, in list of Sphecidae of Oriental Region, as *nivosus*); Bingham, 1897:244 (in revision of wasps and bees of British India); Dalla Torre, 1897:434 (in catalog of world Hymenoptera). – **As *Prionyx nivosus***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae, as questionable synonym of *Prionyx nigropectinatus*).

30. *niveatus* (Dufour)

Sphex niveatus Dufour, 1854a:377, ♀, ♂ (as *niveata*, incorrect original termination). Lectotype: ♀, Algeria: Pontéba, now Oumm ed Drou (MNHN), designated by Menke in R. Bohart and Menke, 1976:133. – Kohl, 1885b:182 (in revision of Palearctic *Sphex*); Ed. André, 1888:128 (in revision of Sphecidae of Europe and Algeria), 10* (bibliographic references); Kohl and Handlirsch, 1889:275 (Turkmenistan); Kohl, 1890b:328 (in revision of world Sphecini); Dalla Torre, 1897:434 (in catalog of world Hymenoptera); Bingham, 1902:216 (South Africa, Malawi); Kohl, 1906a: (Yemen: Aden); Morice, 1911:75 (Algeria: Biskra; prey: *Sphingonotus* sp., a grasshopper); Storey, 1916:107 (Egypt: Maadi, Massara, Moqattam Hills, Suez); Maidl, 1924:246 (Sudan: Tuti Island near Khartum); Roth, 1925:380 (in revision of North African Sphecini); Berland, 1926b:167 (Algeria, Djibouti: Obock; color variation); Guiglia, 1932d:473 (Libya: 85 km south of Gialo); Bischoff, 1933:5 (Morocco); Gussakovskij, 1933b:273 (Iran); Nadig, 1933:103 (Morocco); Guiglia, 1934b:295 (Libya: bibliography and summary of locality records), 1937:184 (Libya: Cyrenaica: Zauiet Msus), 1940a:288 (Libya: Mizda), 1942b:230 (Libya); Honoré, 1944a:60 (in revision of Egyptian Sphecini); Giner Marí, 1945b:359 eastern Morocco: Ixmoart), 1945e:220 (Morocco: Western Sahara), 1947:19 (Morocco: Western Sahara); Berland, 1950a:126 (Mauritania: Ford Gouraud, now Fdérík), 1950b:294 (Niger: Aïr area); de Beaumont, 1950f:396 (Algeria), 1951e:267 (Morocco), 1952c:188 (Algeria: Hoggar); de Beaumont and Bytinski-Salz, 1955:41 (Israel); Leclercq, 1955h:29 (bibliographic references, faunal records from Africa); de Beaumont, 1956a:181 (Libya), 1960b:227 (Libya); Myartseva, 1963b:59 (Turkmenistan: lower Murgab River); Pulawski, 1964:65 (Egypt: Kom Osheim); Myartseva, 1965:83 (Turkmenistan: Akibay; Bayram-Ali district); de Beaumont, 1966:211 (Egypt: Abukir), 1967a:273 (Turkey); Kazenas, 1969a:21 (Kazakhstan: Ili River between Ili and Ayak-Kalkan), 1972b:111 (Kazakhstan), 1978b:41 (in key to Sphecidae of Kazakhstan and Central Asia); Abdu and Shaumar, 1985:229 (Qatar: Rodat Rashed). – **As *Calosphex niveatus***: Grandi, 1959b:287 (new generic assignment, Algeria: Biskra). – **As *Prionyx niveatus***: Myartseva, 1966:48 (new generic assignment, preying on orthopterans), 1971b:80 (occurring only in sand deserts of southern Palearctic Region), 1972a:84 (Turkmenistan); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Guichard, 1980:224 (Oman); Roche, 1981:1 (in checklist of Sphecidae of United Arab Emirates); Islamov, 1986:516 (Uzbekistan: Tashkent Oblast); Steiner, 1986:97 (references to papers on nesting habits); Guichard, 1988a:121 (Arabian Peninsula); Al-Houty, 1989:162 (Kuwait: Sulabiya, Wadi Al-Batin); Guichard, 1991a:338 (Jordan); Kazenas and Nasyrova, 1991:38 (Kazakhstan: preying on *Calliptamus barbarus cephalotes* F.W. and *Dociostaurus tartarus* (Stshelk.)); Kazenas, 1992c:25 (Turkmenistan: Repetek Nature Reserve); Gayubo, Tormos, and Asís, 1993a:201 (first record from Spain: Almería: Cabo de Gata); Al-Houty, 1997:161 (Kuwait: no specific locality); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:62 (in Sphecid Fauna of Western Europe); Kazenas, 1998b:117 (in Sphecid Fauna of Kazakhstan); Nazarova, 1998:40 (Tajikistan: Tigrovaya Balka Nature Reserve); Esenbekova and Kazenas, 2000:9 (southeast Kazakhstan: five localities); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 85 (review of known biology), 2002a:29 (geographic distribution, collecting localities in Kazakhstan); Gadallah and Assery, 2004a:222 (in catalog of Sphecidae of Saudi Arabia); Kazenas, 2004b:99 (Kazakhstan: western Tien Shan Mountains), 2004d:26 (Kazakhstan: northern Caspian region); Nazarova, 2004:104 (Tajikistan: Badakhshan Region: Visav village in Bartash River valley in Rushan District); Gülmez and Tüzün, 2005:47 (Turkey: Ankara Province); Roche, 2007a:45 (in checklist of Egyptian Sphecidae, redescription, as *niveatus niveatus*), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Dollfuss, 2008b:1413 (locality records from Egypt, Jordan, Kazakhstan, Mongolia, Morocco, Tunisia, and Turkmenistan); Ljubomirov and Yildirim, 2008:29 (in catalog of Sphecidae of Turkey); Danilov, 2010b:44 (distribution of Palearctic-Ethiopian type); Schmid-Egger, 2011b:603 (United Arab Emirates: Um al-Quwain), 604 (color photograph of male); Danilov, 2012a:163 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English); Gadallah, Al Dhafer, Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia); Kazenas, 2013a:20, 21 (color photograph of female, short information on geographic distribution and nesting habits); Danilov, 2014b:515 (in key to Sphecidae s.s. of Siberia, not yet found in Siberia); Ebrahimi, 2014:22 (Iran: Māzandarān: Behshahr: Miānkāleh); Kazenas, 2014a:132 (Kazakhstan: Karatau Mountain Range); Schmid-Egger, 2014:623 (United Arab Emirates); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces, as *niveatus niveatus*); Augul, Abdul-Rassoul, and Kaddou, 2015:116 (in key to Sphecini of Iraq),

117 (Iraq: locality records), 119 (illustrations); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey); Mamadou, Mazih, Ghaout, and Hormatallah, 2015:59 (Niger: Agadir Region: Tafidet Valley at 18°09'16"N 9°30'52"E; use of anti-locust chemicals chlorpyrifos ethyl and fenitrothion greatly reduced this species population); Mokrousov, 2015:537 (southeastern European Russia); Danilov, 2017b:215 (in catalog of Sphecidae s.s. of Russia); Danilov and Mokrousov, 2017a:109 (Russia: Astrakhan Oblast', Volgograd Oblast'); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:26 (Iran: known from Mazandaran and North Khorasan provinces); Madl, 2018:944 (in catalog of Hymenoptera of Djibouti); Augul, 2019:497 (recorded from Iraq by Augul, Abdul-Rassoul, and Kaddou, 2015); Danilov, 2019:77 (in review of Palearctic Prionychini, member of subgenus *Calosphex*); Gülmez, 2019:3 (Turkey: Ankara Province: no specific locality); Can and Gülmez, 2019:348 (in key to *Prionyx* of Turkey); Ben Khedher, Yildirim, Braham, and Ljubomirov, 2020a:315 (in list of Tunisian Sphecidae *sensu stricto*; additional records: Tunisia: Kasserin, Sidi Bouzid, Tataouine, and Tozeur Provinces); Bitsch, Barbier, and Jacobs in Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:140 (in Sphecidae Fauna of Europe); Danilov, 2020:319 (specimens from Kazakhstan, Tajikistan, and Russia: Astrakhan' Oblast' in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Gadallah, 2020d:85 (in list of aculeate wasps of Arabian Peninsula); Embergenov, Élmurodova, Amirov and Kimyonazarov, 2022:47 (Uzbekistan: Khiva area); Danilov and Odintsev, 2023:432 (in subgenus *Calosphex*); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

Enodia albopictinata [sic] Taschenberg, 1869:410, ♀ (as *albo-pictinata*, incorrect original hyphenation). Holotype or syntypes: ♀, Sudan: Khartoum (Halle). Synonymized with *Sphex niveatus* by Kohl, 1885b:182.

Podium maracandicum Radoszkowski, 1877:7, ♂. Lectotype: ♂, Uzbekistan: Samarkand (ZMMU), designated by Danilov, 2012b:64. Synonymized with *Sphex nigropectinatus* by Kohl, 1885b:183, and 1885c:165, and with *Prionyx niveatus* by Danilov, 2012b:64. – **As *Sphex maracandica***: Radoszkowski, 1888a:328 (new generic assignment, genitalia).

Sphex suavis F. Morawitz, 1893b:405, ♀, ♂. Lectotype: ♀, Tajikistan: Iskander-kul Lake in Zeravshan River valley at 39°04'N 68°21'E (ZIN), designated by Danilov, 2012b:64. Tentatively synonymized with ..., synonymy confirmed by Danilov, 2012b: – Kohl, 1895:46 (original description copied); Dalla Torre, 1897:442 (in catalog of world Hymenoptera); Danilov, 2016:344 (lectotype preserved in Zoological Institute, Sankt Petersburg, Russia).

Sphex afghaniensis de Beaumont, 1970a:391, ♀. Holotype: ♀, Afghanistan: Kabul (Brno Mus.). Synonymized with *Prionyx niveatus* by Danilov, 2012b:64. – **As *Prionyx afghaniensis***: R. Bohart and Menke, 1976:131 (new generic assignment, in checklist of world Sphecidae). Dunford, Turbyville, and Leavengood, 2014:11 (listed as medically important in Afghanistan).

ssp. *ettingol* (Tsuneki)

Sphex niveatus ettingol Tsuneki, 1971k:143, ♀, ♂. Holotype: ♂, Mongolia: Bayanhongor Aymag: Tsagan Bogd ul (TMB). – **As *Prionyx niveatus ettingol***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae).

31. *notinitidus* (Willink)

Chlorion notinitidum Willink, 1951:184, ♀, ♂. Holotype: ♀, Argentina: no specific locality (MACN). – **As *Prionyx notinitidus***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Sielfeld, 1980b:72 (in checklist of Chilean Sphecidae, as *notonitidus*); Pérez d'Angello, 1989:264 (Chile: Magellan Region); Amarante, 2002:72 (in catalog of Neotropical Sphecidae); Chiappa, 2012:8 (Chile: Región de Valparaíso: no specific locality, as *notonitidus*); Danilov and Odintsev, 2023:434 (tentatively in subgenus *Priononyx*)

32. *nudatus* (Kohl)

Sphex nudatus Kohl, 1885:187, ♀, ♂. Syntypes: Ukraine: Yekaterinoslav, now Dnepropetrovsk, Russia: Sarepta, now, Krasnoarmeysk; Caucasus: no specific locality; Turkey: Brussa, now Bursa; Dalmatia, now coastal Croatia and Montenegro; Egypt: no specific locality (NHMW). – Gasperini, 1887:18 (recorded from Dalmatia, now Croatia, by Kohl, 1885b); F. Morawitz, 1889a:129 (China: Ordos Region); Kohl, 1890:342 (incorrectly synonymized *Sphex nudatus* with *Sphex mocsaryi*); E. Saunders, 1904c:636 (Spain); Kohl, 1913b:15 (Russia: Voronezh Oblast': Valuyki at 50°14'N 38°08'E);

Fahringer, 1922:177 (Turkey); Kuznetsov-Ugamskij, 1927:249 (Kazakhstan: Aulie-Ata, now Djambul); Honoré, 1944a:68 (in revision of Egyptian Sphecini); de Beaumont, 1957b:130 (northern Iran; valid name; diagnostic characters); Suárez, 1959:53 (Spain: Provincia de Almería); de Beaumont, 1957b:130 (Iran: Aliabad, Kamalabad, Rayne, Sirah; taxonomic history, diagnostic characters), 1962b:19 (Spain); Ceballos, 1964:87 (in supplement to catalog of Spanish Sphecidae); de Beaumont, 1967a:272 (Turkey); Kazenas, 1969a:21 (Kazakhstan: Alma Ata, Ili River, Kegen' River, Sharyn River); de Beaumont, 1970a:393 (Afghanistan); Kazenas, 1972b:112 (Kazakhstan), 1974b:109 (feeding on flowers of *Tamarix* sp., Tamaricaceae, *Statice gmelini* Willd., Plumbaginaceae, and *Euphorbia* spp., Euphorbiaceae, in Kazakhstan), 112 (feeding on flowers of *Apocynum lancifolium* Russ., Apocynaceae, in Kazakhstan), 1978b:44 (in key to Sphecidae of Kazakhstan and Central Asia); Pulawski, 1978:184 (in key to Sphecidae of European part of USSR); Kuznetsova, 1990:17 (Russia: Voronezh Oblast': Galich'ya Gora Nature Reserve). – **As *Spheg mocsaryi* var. *nudatus***: F. Morawitz, 1891a:202 (new status, Russia: Astrakhan Government); Dalla Torre, 1897:432 (in catalog of world Hymenoptera); Ceballos, 1956:364 (in catalog of Hymenoptera of Spain). – **As *Enodia nudata***: Radoszkowski, 1892:586 (new generic assignment, male genitalia). – **As *Prionyx nudatus***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae); Mingo and Gayubo, 1983:155 (Spain; redescription); Gayubo and Tormos, 1984:9 (Spain: Valencia); Gayubo, 1986b:36 (Spain: Andalucía); Gayubo and Sanza, 1986:28 (Spain: Burgos, Soria); Islamov, 1986:516 (Uzbekistan: Tashkent Oblast'); Ebrahimi, 1993:97 (Iran); Torregrosa, Gayubo, Tormos, and Asís, 1993:11 (Spain: Alicante Province); Tormos, Asís, and Gayubo, 1994:187, 195 (Spain: Albacete Province); Gorobchishin, 1995:17 (Ukraine: Kanev Nature Reserve), 1996:53 (Ukraine: Kanev Nature Reserve); Voblenko, Gorobchishin, and Nesterov, 1996:14 (Ukraine: Polesye Region); Gorobchishin, 1998a:48 (Ukraine: Kiev and surroundings); Kazenas, 1998b:119 (in Sphecid Fauna of Kazakhstan); Anan'eva and Kochetkov, 1999:6 (Russia: Ryazan Oblast': no specific locality); Esenbekova and Kazenas, 2000:10 (southeastern Kazakhstan: near Alma Ata, 10 km northwest of Chemolgan, 4 km southeast and 22 km west of Furmanovka, 45 km northwest of Suzak); Gayubo, González, and Torres, 2000:184 (Spain: Salamanca Province); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia); Kazenas and Esenbekova, 2001:133 (Kazakhstan: Almatinskiy Nature Reserve); Kazenas, 2002a:29 (geographic distribution, collecting localities in Kazakhstan); Shkuratov, 2002a:383 (Russia: common in Rostov Oblast'), 2004b:164 (Russia: Rostov Oblast': Gosudarstvennyi Muzey-Zapovednik M.A. Sholokhova); Protsenko, 2003:68, 69 (Ukraine: Odessa Oblast': Malyi Tataru island in Danube delta at 45.21°N 29.00°E); Gorobchishin, 2004:34 (Ukraine: Zaporiz'ka Oblast': Obitichna Kosa Nature Reserve); Kazenas, 2004b:99 (Kazakhstan: western Tien Shan Mountains), 2004d:26 (Kazakhstan: northern Caspian region); Shkuratov, 2004a:73 (Russia: Rostov Oblast'); Gayubo and Özbek, 2005:8 (Turkey: many localities); Gülmez and Tüzün, 2005:47 (Turkey: Ankara Province); Shoreiko, 2005a:162 (Ukraine: Crimea), 2005b:97 (Ukraine: Crimea: Karadagh Nature Reserve); Yildirim and Ljubomirov, 2005:1787 (Turkey: Erzurum: Pasinler; Kars: Sarikamis); Baghirov, 2007:93 (Russia: southwestern Siberia); Kazenas, 2007a:89 (Kazakhstan: Akkala Oblast': Kurgandzhin Nature Reserve and vicinity); Roche, 2007a:42 (in checklist of Egyptian Sphecidae), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Danilov, 2008:348 (Russia: Altayskiy Kray: Barnaul area); Dollfuss, 2008b:1413 (locality records from Kazakhstan, Kyrgyzstan, Morocco, Russia, Turkey, and Ukraine); Gayubo, González, Tormos, and Asís, 2008:136 (Spain: Salamanca: Parque Natural de Las Batuecas – Sierra de Francia); Kazenas, 2008b:111 (Kazakhstan: foothills of Zailiskiy Alatau: frequently found on loess cliffs), 2008c:255 (Kazakhstan: village Koktum south of Lake Alakol'); Ljubomirov and Yildirim, 2008:29 (in catalog of Sphecidae of Turkey); Nemkov, 2008b:17 (in catalog of Sphecidae of Asiatic Russia); Baños-Picón, Asís, Gayubo, and Tormos, 2009:310 (Spain: frequency of specimens collected with hand nets and Malaise traps); Danilov, 2009:54 (Russia: Western Siberia: Kulundinskaya Steppe); González, Gayubo, Asís, and Tormos, 2009:622 (Spain: Salamanca and Zamora provinces: Arribes del Duero Natural Park); Nemkov, 2009b:46 (in new catalog of Sphecidae and Crabronidae of Asiatic Russia); Shoreiko, 2009:366 (in list of Sphecidae *sensu lato* of Crimea); Danilov, 2010b:45 (distribution of Tethyan type); Mokrousov, 2010a:60 (Russia: Nizhniy Novgorod Oblast': no specific localities), 2010b:100 (Russia: Nizhniy Novgorod Oblast' no specific locality); Mokrousov and Zryanin, 2010:98 (needs to be protected in Russia: Novgorod Oblast'); Rudoiskatel', 2010:147 (Russia: southern Ural Mountains); Shoreiko and Kononov, 2010:12 (Ukraine); Mokrousov, Berezin, and Egorov, 2011:65 (Russia: Chuvash Republic); Danilov, 2012a:164 and 2012b:61 (in keys to *Prionyx* of

Russia and adjacent countries, respectively in Russian and English), 2012b:65 (bibliographic references, geographic distribution); Kazenas, 2012b:168 (Kazakhstan: Korgalzhin State Nature Reserve); Prisniy, 2012:46 (Russia: Belgorod Oblast'); Protsenko, Fateryga, Ivanov, and Puzanov, 2012:58 (Ukraine: Crimea); Yildirim, 2012:74 (Turkey: Iğdır: Melekli, Tuzluca; Kars: Sarıkamış: Karakurt); Kazenas, 2013a:21, 22 (color photographs of females, short information on geographic distribution and nesting habits); Baldock, 2014:354 (Spain: Island of Mallorca); Danilov, 2014a:424 (Russia: Siberia: Novosibirsk Oblast', Altayskiy Kray, Tyva), 2014b:516 (in key to Sphecidae s.s. of Siberia); Dunford, Turbyville, and Leavengood, 2014:11 (listed as medically important in Afghanistan); Ebrahimi, 2014:22 (Iran: Akborz, Ardebil, Āzarbāijān-e Sharghi, Bushehr, Esfahān, Ghazvin, Gilān, Golestān, Kermān, Māzandarān, and Tehrān provinces); Kazenas, 2014a:132 (Kazakhstan: Karatau Mountain Range); Mokrousov and Vafin, 2014:54 (Russia: Republic of Tatarstan: village Antonovka in Kamsko-Ust'inskiy District); Ruchin and Antropov, 2014:34 (Russia: Republic of Mordovia); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces); Gülmez and Can, 2015:12 (usability of ITS2 and 28S-D2 gene regions for species identification); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey); Samin, Bagriacik, and Monaem, 2015:195 (Iran: Kermanshah: Kermanshah); Shorenko, 2015:317 (in list of Sphecidae *sensu lato* of Crimea); Gülmez and Dizer, 2016:59 (Turkey: Tokat Province); Mokrousov and Popov, 2016:564 (Russia: Krasnodarskiy Kray); Yildirim, Ljubomirov, Özbek, and Yüksel, 2016:6 (Turkey: Antalya and Erzurum provinces); Danilov, 2017b:215 (in catalog of Sphecidae s.s. of Russia); Danilov and Mokrousov, 2017a:109 (Russia: Kalmykia: near Lagan); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:26 (Iran: known from Alborz, Ardabil, East Azerbaijan, Bushehr, Golestan, Guilan, Isfahan, Kerman, Kermanshah, Markazi, Mazandaran, Qazvin, and Tehran provinces); Shorenko, 2017:76 (in Crimea collected in May through September), 2018:127 (Crimea, including localities, habitats, and number of specimens); Can and Gülmez, 2019:349 (in key to *Prionyx* of Turkey); Danilov, 2019:78 (in review of Palearctic *Prionychini*, member of subgenus *Prionyx* s.s.); Gülmez, 2019:3 (Turkey: Amasya, Ankara, Sivas, and Tokat provinces: no specific localities), 4 (abnormal forewing venation); Ruchin and Antropov, 2019a:13221 (Russia: Mordovia State Nature Reserve); Shorenko, 2019:211 (among commonest sphecid species in Crimea); Ben Khedher, Yildirim, Braham, and Ljubomirov, 2020a:316 (in list of Tunisian Sphecidae *sensu stricto*); Bitsch, Barbier, and Jacobs *in* Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:140 (in Sphecid Fauna of Europe); Danilov, 2020:320 (specimens from Kazakhstan, Kyrgyzstan, and Russia: Astrakhan' Oblast', Dagestan, and Siberia in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Maharramov, Mokrousov, and Proshchalykin, 2020:46 (Azarbaijan: Nakichivan Autonomous Republic); Shorenko, 2020:47 (Crimea: Karadag Nature Reserve); Can and Gülmez, 2021a:3013 (Turkey: Erzincan, Giresun, and Sivas provinces); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*); Embergenov, Akhmedov, and Medetov, 2023:24 (Uzbekistan: southern Lake Aral region); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

As *Sphex mocsaryi*: Kohl, 1890b:342 (in revision of world Sphecini), corrected to *Sphex nudatus* by de Beaumont, 1957b:130; Roth, 1925:385 (in revision of North African Sphecini); Giner Marí, 1943a:83 (in Sphecid Fauna of Spain), present correction based on geographic distribution

Sphex mocsaryi. var. *denudatus* Kohl: A. Costa, 1893b:3 (Tunisia, nomen nudum?).

33. *parkeri* Bohart and Menke

Prionyx parkeri Bohart and Menke, 1963:154, ♀, ♂. Holotype: ♂, USA: California: Kern County: Mill Potrero (UCD). – Krombein, 1964c:18 (nest and prey); Lavigne and Pfadt, 1966:14 (prey and nest digging), 31 (Wyoming; preying on *Aulocara elliotti* (Thomas), Acrididae); Evans, 1975a:263 (slow colonizer of new habitats); R. Bohart and Menke, 1976:133 (in checklist of world Sphecidae); Ch. Porter, 1978:172 (Texas); Krombein, 1979b:1586 (in catalog of North American Hymenoptera); Steiner, 1981:305 (grasshopper prey is paralyzed by four stinging, exactly as in *Liris*, but stinging starts with suboesophageal nervous ganglion), 1982a:2 (grasshopper prey anti-predatory behavior), 1983a:130 (puncturing dorsolateral neck region of grasshopper prey, subsequent drinking of body fluid), 1984a:152 (prey never resumes normal activities after being stung, in contrast to *Larra*); Cornett, 1986:224 (California: Palm Springs; on flowers of *Washingtonia filifera* (Wendl), Arecaceae); Menke, 1986c:36 (Arizona: Gila County: Carrizo); Piek and Spanjer,

1986:188 (in list of Sphecidae with known prey); Steiner, 1986:97 (references to papers on nesting habits); Frommer, 1988:95 (California: Riverside County: Deep Canyon); Ahlstrom, 1995:107 (in checklist of insects of North Carolina); Hanson and Menke, 1995:637 (known from Costa Rica); Meagher and Mitchell, 1999:368 (collected in pheromone- and synthetic floral volatile-baited traps); Weissmann and Kondratieff, 1999:78 (Colorado: Great Sand Dunes National Monument); Amarante, 2002:72 (in catalog of Neotropical Sphecidae); Ruíz Cancino, Coronado Blanco, Varela Fuente, and Horta Vega, 2002:670 (in checklist of Mexican Sphecidae); Ohl and Linde, 2003:149 (number of ovarioles); Buck, 2004:24 (Canada: in checklist of Sphecidae of Ontario), 29 (distribution within Ontario), 33 (first record from Canada: Ontario: Guelph, Oakville, Pinery Provincial Park); Horta Vega, Pinson Domínguez, Barrientos Lozano, and Correa Sandoval, 2007:48 (Mexico: Tamaulipas); Dollfuss, 2008b:1414 (Mexico: Hidalgo: Metztitlán 70 km of Pachuca; Guerrero: Acahuizotla 35 km south of Chilpanzingo); Cope, Campbell, Grodsky, and Ellis, 2019:9 (Florida: Gainesville, collected in emergence traps); Danilov and Odintsev, 2023:434 (in subgenus *Priononyx*); Buck and Bennett, 2025:224 (in checklist of Sphecidae sensu lato of Canada, Alaska, and Greenland: Canada: Ontario, first record from New Brunswick and Quebec: no specific locality).

As *Prionyx pubidorsus*: Evans, 1958a:183 (nesting behavior) and Linsley, 1962:156 (sleeping aggregations), corrected to *Prionyx parkeri* by R. Bohart and Menke, 1963:156.

34. *persicus* (Mocsáry)

Sphex persicus Mocsáry, 1883:33, ♂ (as *Persicus*, incorrect original capitalization). Lectotype: ♂, Iran: no specific locality (TMB), designated by Danilov, 2012:65. – Kohl, 1885b:181 (in revision of Palearctic Sphecini); Ed. André, 1888:146 (in revision of Sphecidae of Europe and Algeria), 8* (bibliographic references); Kohl, 1889a:24 (as new synonym of *Sphex sirdariensis*). – As *Prionyx persicus*: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia); Danilov, 2012a:163 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, in Russian and in English, respectively), 2012b:65 (bibliographic references, geographic distribution); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:26 (Iran: known from Khorasan-e Razavi Province); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

Sphex hispidus F. Morawitz, 1890:576, ♀. Lectotype: Turkmenistan: Artchman (ZIL), designated by Danilov, 2012b:65. Tentatively synonymized with *Sphex persicus* by de Beaumont, 1957b:130, synonymy confirmed by Danilov, 2012b:65. – Kohl, 1890b:352 (original description copied); Dalla Torre, 1897:425 (in catalog of world Hymenoptera); F. Morawitz, 1897a:151 (description of ♂); Gussakovskij, 1933b:278 (Iran); de Beaumont, 1957b:130 (Iran: Mughan); Danilov, 2016:340 (lectotype preserved in Zoological Institute, Sankt Petersburg, Russia). – As *Priononyx hispidus*: de Beaumont, 1968b:150 (new generic assignment, member of *macula* species group; probably synonym of *Sphex persicus*). – As *Prionyx hispidus*: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae).

35. *popovi* Guichard

Prionyx popovi Guichard, 1988a:122, ♀, ♂. Holotype: ♀, Mali: Gao (BMNH). Paratypes: Qatar. – Gadallah, 2020d:86 (in list of aculeate wasps of Arabian Peninsula); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*).

36. *pseudostriatus* (Giner Marí)

Sphex pseudostriatus Giner Marí, 1944:346, ♂. Holotype: ♂, Peru: no specific locality (Mus. Barcelona). – As *Prionyx pseudostriatus*: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae, possibly a synonym of *Prionyx fervens*); Amarante, 2002:72 (in catalog of Neotropical Sphecidae), 2005a:14 (correction to his 2002 catalog); Rasmussen and Asenjo, 2009:16 (in checklist of Crabronidae of Peru); Danilov and Odintsev, 2023:434 (in subgenus *Priononyx*).

Sphex vaqueroi Giner Marí, 1944:347, ♀, ♂. Syntypes: Peru: no specific locality (Mus. Barcelona). Synonymized with *Prionyx pseudostriatus* by Menke in R. Bohart and Menke, 1976:134.

37. *pumilio* (Taschenberg)

Pseudosphex pumilio Taschenberg, 1869:420, ♀. Holotype or syntypes: ♀, Argentina: Mendoza (Halle). – Burmeister, 1872:240 (Argentina: Mendoza). – **As *Sphex pumilio***: Kohl, 1890b:369 (new generic assignment, in revision of world Sphecini); Dalla Torre, 1897:438 (in catalog of world Hymenoptera); Schrottky, 1903b:123 (in checklist of Hymenoptera of Argentina, Paraguay, and Uruguay); Berland, 1926c:204 (Argentina and Brazil: locality records); Liebermann, 1931:81 (in revision of Argentinean Sphecini). – **As *Chlorion pumilio***: Fernald, 1907:266 (new generic assignment, Argentina, variation, as *pumilo*). – **As *Neosphex pumilio***: Schrottky, 1913a:225 (new generic assignment, Argentina); Willink, 1951:95 (in revision of Argentinean Sphecini). – **As *Prionyx pumilio***: R. Bohart and Menke, 1963:151 (new generic assignment, member of *Prionyx pumilio* species group), 1976:134 (in checklist of world Sphecidae); Nascimento and Overal, 1980:8 (Chile); Sielfeld, 1980b:72 (in checklist of Chilean Sphecidae); Amarante, 2002:73 (in catalog of Neotropical Sphecidae); Dollfuss, 2008b:1414 (locality records from Chile); Vázques, Aschero, and Stevani, 2008:9 (Argentina: Central Monte desert); Buys, 2010:172 (Brazil: state of Paraíba: Santa Luzia at 06°52'20"S 36°55'0"W); Chiappa, 2012:8 (Chile: Región de Valparaíso: no specific locality); López García, Mazzitelli, Fruitos, González, Marcucci, Giusti, Alemanno, del Barrio, Portela, and Debandi, 2019:314 (Argentina: Mendoza: departamento de Tupungato: vineyards in Gualtallary); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Neosphex*); Danilov and Odintsev, 2023:433 (in subgenus *Neosphex*).

Sphex dolichoderus Kohl, 1890b:370, ♀. Syntypes: ♀, Chile: no specific locality (NHMW). Synonymized with *Chlorion pumilio* by Fernald, 1907:266. – Dalla Torre, 1897:421 (in catalog of world Hymenoptera); R. Bohart and Menke, 1976:134 (as tentative synonym of *Prionyx pumilio*).

Neosphex albospiniferus Reed, 1894:627, ♀, ♂. Syntypes: Chile: Valparaíso (depository?). – R. Bohart and Menke, 1976:134 (in checklist of world Sphecidae, as tentative synonym of *Prionyx pumilio*); Menke and Bohart, 1979:115 (as *albospinifer*, unjustified emendation); Amarante, 2005a:14 (correction to his 2002 catalog). – **As *Sphex albospinifer***: Kohl in Dalla Torre, 1897:414 (new generic assignment, in catalog of world Hymenoptera).

38. *radoszkowskyi* (Kohl)

Sphex radoszkowskyi Kohl, 1888a:151, ♀ (as *Radoszkowskyi*, incorrect original capitalization). Syntypes: Uzbekistan: Khiva (ZMHU). – Kohl, 1890b:345 (in revision of world Sphecini, as *radoszkowskii*); Dalla Torre, 1897:438 (in catalog of world Hymenoptera); Kazenas, 1969a:22 (Kazakhstan: Ili River), 1972b:113 (Kazakhstan: Ayak-Kalkan on Ili River in Alma Ata Oblast'), 1978b:41 (in key to Sphecidae of Kazakhstan and Central Asia). – **As *Priononyx radoszkowskii***: de Beaumont, 1968b:149 (new generic assignment, member of *Priononyx macula* species group). – **As *Prionyx radoszkowskyi***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Kazenas, 1992c:25 (Turkmenistan: Repetek Nature Reserve), 1998b:122 (in Sphecid Fauna of Kazakhstan); Esenbekova and Kazenas, 2000:10 (southeast Kazakhstan: near Dubun' landing place on Ili River); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:30 (geographic distribution, collecting localities in Kazakhstan), 2004d:27 (Kazakhstan: northern Caspian region), 2008a:98 (southeastern Kazakhstan: Ili River valley); Danilov, 2010b:45 (distribution of Tethyan type), 2012a:163 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:65 (bibliographic references, geographic distribution); Can and Gülmez, 2019:347 (first record from Turkey: Erzincan: Ergen Mountain at 39°38'12"N 39°30'21"E), 348 (in key to *Prionyx* of Turkey); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*); Embergenov, Akhmedov, and Medetov, 2023:24 (Uzbekistan: southern Lake Aral region); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

39. *reymondi* (Roth)

Sphex reymondi Roth, 1954:31, ♀, ♂. Holotype: ♀, Algeria: Oran Department: Ougarta (MNHN). – de Beaumont, 1968b:149 (member of *subfuscatus* species group). – **As *Prionyx reymondi***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

40. *saevus* (F. Smith)

Harpactopus saevus F. Smith, 1856:265, ♀, ♂. Syntypes: Australia: Western Australia: Swan River; and Queensland: Cape Upstart (BMNH). – de Saussure, 1867:42 (Australia: Sydney; redescription); Froggatt, 1892:211 (in catalog of Australian Hymenoptera). – **As *Sphex saevus***: Kohl, 1890b:366 (new generic assignment, in revision of world Sphecini); Dalla Torre, 1897:440 (in catalog of world Hymenoptera); R. Turner, 1910a:343 (in key to Australian Sphecini); Berland, 1926c:206 (one ♂ from Australia in MNHN); von Schulthess, 1935:304 (Indonesia: Flores: Endeh). – **As *Prionyx saevus***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Evans, Hook, and Matthews, 1982:223 (nesting habits); Baker and Pigott, 1983:67 (nesting habits); Cardale, 1985:228 (in catalog of Australian Sphecidae); Naumann, 1998:183 (Australia: northwest Queensland: Musselbrook area, approximately 18°40'S 138°23'E); Pagliano, 2003a:505 (Australia: Northern Territory: Katherine); Dollfuss, 2008b:1414 (Australia: Coopers Creek); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

ssp. harpax (Kohl)

Sphex harpax Kohl, 1898a:333, ♂. Holotype: ♂, Indonesia: Timor (TMB). – **As *Priononyx harpax***: van der Vecht, 1957c:352 (new generic assignment, description of ♀, redescription of ♂, Lesser Sunda Islands: locality records). – **As *Prionyx saevus harpax***: R. Bohart and Menke, 1976:134 (new generic assignment, new status, in checklist of world Sphecidae).

41. *semistriatus* (Schrottky)

Priononyx semistriatus Schrottky, 1920:185, ♀. Holotype or syntypes: ♀, Paraguay: Puerto Bertoni (depository?). – **As *Chlorion semistriatum***: Willink, 1948a:317 (new generic assignment, probably a synonym of *Chlorion thomae* or *Ch. striatulum*), 1951:199 (in revision of Argentinean Sphecini); not listed by R. Bohart and Menke, 1976; Amarante, 2002:73 (in catalog of Neotropical Sphecidae); Danilov and Odintsev, 2023:434 (in subgenus *Priononyx*).

42. *senegalensis* (Arnold)

Sphex senegalensis Arnold, 1951:144, ♂. Holotype: ♂, Senegal: Kaolack (BMNH). – Leclercq, 1955h:37 (bibliographic reference). – **As *Prionyx senegalensis***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*).

43. *senilis* (Morice)

Sphex senilis Morice, 1911:75, ♀, ♂. Syntypes: Algeria: Biskra (OXUM). – Roth, 1925:381 (in revision of North African Sphecini); de Beaumont, 1962c:221 (Arabia: El Riyadh), 1968b:152 (recognition characters). – **As *Prionyx senilis***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Guichard, 1988a:121 (Saudi Arabia); Gadallah and Assery, 2004a:222 (in catalog of Sphecidae of Saudi Arabia, as *senelis*); Roche, 2007a:46 (in checklist of Egyptian Sphecidae), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Dollfuss, 2008b:1415 (Jordan: Rawayshid, Mongolia: five localities; recognition characters); Gadallah, Al Dhafer, Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia); Gadallah, 2020d:86 (in list of aculeate wasps of Arabian Peninsula); Danilov and Odintsev, 2023:432 (in subgenus *Calosphex*).

As Parasphex marginatus var. *leucosoma*: W. Schulz, 1905b:34 (Algeria: Biskra), corrected to *Sphex senilis* by de Beaumont, 1968b:153.

Sphex niveatus var. *biskrensis* Roth, 1925:381. Proposed conditionally as substitute name for *Sphex senilis*, should the latter be found a synonym of *Prionyx niveatus*.

44. *sennae* (Mantero)

Sphex sennae Mantero, 1902:200, ♀ (as *Sennae*, incorrect original capitalization). Holotype: ♀, Argentina: Patagonia: Río Santa Cruz (MSG). – **As *Priononyx sennae***: Holmberg, 1903:504 (new generic assignment, listed); Schrottky, 1913a:225 (Argentina: Santa Cruz). – **As *Chlorion sennae***: Willink, 1948a:317 (new generic assignment, probably a synonym of *Chlorion thomae* or *striatulum*), 1951:291 (original description translated into Spanish). – **As *Prionyx sennae***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Amarante, 2002:73 (in catalog of Neotropical Sphecidae); Danilov and Odintsev, 2023:434 (in subgenus *Priononyx*).

45. *simillimus* (Fernald)

Chlorion simillimum Fernald, 1907:264, ♀, ♂. Syntypes: Argentina: Cordoba: Cordoba (MCZ). – Fernald, 1931a:441 (as synonym of *Chlorion neoxenum*); Willink, 1951:181 (in revision of Argentinian Sphecini). – **As *Priononyx simillimus***: Schrottky, 1913a:225 (new generic assignment, as synonym of *Priononyx neoxenus*). – **As *Sphex simillimus*** [sic]: Liebermann, 1931:23 (new generic assignment, in revision of Argentinian Sphecini). – **As *Prionyx simillimus***: Menke, 1962a:63 (new generic assignment, correction to original description); R. Bohart and Menke, 1976:134 (in checklist of world Sphecidae); Amarante, 2002:73 (in catalog of Neotropical Sphecidae), 2005a:14 (correction to his 2002 catalog); Danilov and Odintsev, 2023:434 (in subgenus *Priononyx*).

Sphex tucumanensis Strand, 1910a:133, ♂. Holotype or syntypes: ♂, Argentina: Tucumán: no specific locality (ZMHU). Synonymized with .. *simillimus* by .. – Berland, 1926c:203 (Argentina: locality records); Strand, 1927:254 (in list of species described by author); Liebermann, 1931:79 (in revision of Argentinean Sphecini); Willink, 1951:1 (possibly a synonym of *Chlorion neoxenum*).

46. *sirdariensis* (Radoszkowski)

Sphex sirdariensis Radoszkowski, 1877:9, ♂. Lectotype: ♂, Uzbekistan: on Syr-Darya River (ZMMU), designated by Danilov, 2012b:65. – Kohl, 1885b:206 (original description copied); Radoszkowski, 1886a:25 (Turkmenistan); Ed. André, 1888:145 (in revision of Sphecidae of Europe and Algeria), 8* (bibliographic references); Radoszkowski, 1888a:328 (description of male genitalia); Kohl, 1890b:347 (in revision of world Sphecini); Dalla Torre, 1897:440 (in catalog of world Hymenoptera); de Beaumont, 1968b:150 (taxonomy). – **As *Sphex occitanicus* var. *syrdariensis***: Gussakovskij, 1934a:2 (new status, possibly in error). – **As *Prionyx sirdariensis***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Kazenas, 1992c:25 (Turkmenistan: Repetek Nature Reserve), 1998b:123 (in Sphecid Fauna of Kazakhstan, unknown to author), 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:30 (geographic distribution, collecting localities in Kazakhstan); Danilov, 2010b:45 (distribution of Tethyan type), 2012a:164 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

47. *songaricus* (Eversmann)

Sphex songaricus Eversmann, 1849:368, ♀. Holotype or syntypes: ♀, "in campis Kirgisorum orientalibus", now Kazakhstan: no specific locality (ZIN). – Kohl, 1885b:206 (original description copied); Ed. André, 1888:126 (in revision of Sphecidae of Europe and Algeria, as *sougaricus*), 7* (bibliographic references, as *sougaricus*); Kohl and Handlirsch, 1889:275 (Turkmenistan); Kohl, 1890b:340 (in revision of world Sphecini); Radoszkowski, 1893a:58 (Turkmenistan); Dalla Torre, 1897:440 (in catalog of world Hymenoptera); Gussakovskij, 1933b:273 (Iran), 1935:413 (Tajikistan); de Beaumont and Bytinski-Salz, 1955:41 (Israel); de Beaumont, 1961e:2 (Iraq); Myartseva, 1963b:59 (Turkmenistan: lower Murgab River), 1964a:75 (nesting habits in Turkmenistan), 1965:84 (Turkmenistan: Akibay); de Beaumont, 1967a:273 (Turkey), 1969:81 (Turkey), 1970a:393 (Afghanistan). – **As *Prionyx songaricus***: Myartseva, 1966:48 (new generic assignment, preying on orthopterans), 1972a:84 (Turkmenistan); R. Bohart and Menke, 1976:134 (in checklist of world Sphecidae); Ebrahimi, 1993:98 (Iran); Nazarova and Shomirsaidov, 1997:23 (Tajikistan: fruit tree orchards in Vakhsh River valley); Kazenas, 1998b:123 (in Sphecid Fauna of Kazakhstan); Nazarova, 1998:40 (Tajikistan: Tigrovaya Balka Nature Reserve); Esenbekova and Kazenas, 2000:10 (southeastern Kazakhstan: 35–45 km northwest of Kapchagay); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:30 (geographic distribution, collecting localities in Kazakhstan); Nazarova, 2005:93 (alfalfa fields in southwestern Tajikistan); Dollfuss, 2008b:1415 (locality records from Kazakhstan, Syria, Turkmenistan, Turkey, and Uzbekistan); Ljubomirov and Yildirim, 2008:30 (in catalog of Sphecidae of Turkey); Danilov, 2010b:44 (distribution of Tethyan type); Ghazi-Soltani, Ebrahimi, Iranipour, and Pour Abad, 2010:797 (Iran: East Azarbaijan: Jolfa); Murai and Amr, 2011:109, 120 (first record from Syria: Al Thawrah Nature Reserve at 35°51'N 28°38'E); Danilov, 2012a:164 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and in English), 2012b:67 (bibliographic references, geographic distribution), 2014b:516 (in key to Sphecidae s.s.).

of Siberia, not yet found in Siberia); Dunford, Turbyville, and Leavengood, 2014:11 (listed as medically important in Afghanistan); Ebrahimi, 2014:23 (Iran: Khorāsān-e Razavi, Tehrān, and Zanjān provinces); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey); Yildirim, Ljubomirov, Özbek, and Yüksel, 2016:6 (Turkey: Şanlıurfa: Bozova: Yaylak); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:27 (Iran: known from Alborz, East Azerbaijan, Kerman, Khorasan-e Razavi, Markazi, Tehran, and Zanjan provinces); Augul, 2019:498 (recorded from Iraq by de Beaumont, 1961e); Can and Gülmez, 2019:349 (in key to *Prionyx* of Turkey); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Prionyx* s.s.), 2020:320 (specimens from Turkmenistan in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

Sphex tenuicornis F. Morawitz, 1890:580, ♀. Lectotype: ♀, Turkmenistan: Ashkhabad (ZIN), designated by Danilov, 2012:67. Synonymized with *Sphex songaricus* by Kohl, 1890b:340, synonymy confirmed in 1895:47. – Myartseva, 1963b:59 (Turkmenistan: lower Murgab River), 1965:84 (Turkmenistan: Akibay); Danilov, 2016:344 (lectotype preserved in Zoological Institute, Sankt Petersburg, Russia).

48. *spinolae* (F. Smith)

Sphex chilensis Spinola, 1851a:399, ♀, ♂ junior primary homonym of *Sphex chiliensis* Lepelletier de Saint Fargeau, 1845 (Article 58.15). Lectotype: ♀, Chile: no specific locality (M. Spinola collection, Torino), designated by Menke in R. Bohart and Menke, 1976:134. – Ruiz Pereira, 1924:101 (Chile: Cerro San Cristóbal); Janvier, 1926:59 (nesting habits), 1928:203 (nesting habits); Gazulla and Ruiz Pereira, 1929:299 (Chile: Hacienda de "Las Mercedes"); Ruíz Pereira and Stuardo, 1936:321 (Chile: Las Termas de Chillán); Ruíz Pereira, 1937:164 (Chile: Coquimbo Province); Soku, 1943:265 (Peru); Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino); Piek and Spanjer, 1986:189 (in list of Sphecidae with known prey); Cabrera La Rosa, 1993:71 (Peru: La Molina); Pagliano, 2008:532 (specimens in M. Spinola collection, Torino, including lectotype; some specimens are *Prionyx bifoveolatus* and *P. neoxenus*). – **As *Am-mophila chilensis***: Fraga, 1938:200 (new generic assignment, Chile: Hacienda Mauro). – **As *Priononyx chilensis***: Reed, 1894:626 (new generic assignment, in revision of Chilean Sphecidae). – **As *Prionyx chilensis***: R. Bohart and Menke 1976:134 (new generic assignment, in checklist of world Sphecidae); Amarante, 2002:72 (new generic assignment, in catalog of Neotropical Sphecidae); Dollfuss, 2008b:1408 (locality records from Argentina and Chile); Rasmussen and Asenjo, 2009:16 (in checklist of Crabronidae of Peru); Buys, 2011a:411 (Peru: Arequipa); Rodrigues and Buys, 2013:214 (Brazil; Espírito Santo: after Amarante, 2002); Buys and Rodrigues, 2014:40 (Brazil: State of Espírito Santo: Linhares); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Neosphex*); Danilov and Odintsev, 2023:433 (in subgenus *Neosphex*).

Sphex spinolae F. Smith, 1856:260 (as *Spinolae*, incorrect original capitalization). Substitute name for *Sphex chilensis* Spinola, 1851). – Kohl, 1890b:364 (in revision of world Sphecini); Dalla Torre, 1897:441 (in catalog of world Hymenoptera); Gribodo, 1895:211 (Chile); Brèthes, 1908:144 (in revision of *Sphex thomae* species group; Chile, Patagonia); Kieffer and Herbst, 1909:122 (visiting galls of *Lecanium resinatum* Kieffer and Herbst, Coccidae); Herbst, 1921a:107 (comparison with *Sphex omissus* Kohl); Berland, 1929b:312 (miscellaneous locality records); Liebermann, 1931:26 (in revision of Argentinean Sphecini). – **As *Harpactopus spinolae***: Schrottky, 1913a:225 (new generic assignment, Argentina: Santa Cruz). – **As *Chlorion spinolae***: Willink, 1948a:319 (new generic assignment, in key to *Chlorion thomae* species group); Zapata, 1974:37 (Chile: Lampa near Santiago). – **As *Priononyx spinolae***: Evans, 1958a:185 (new generic assignment, observations by Claude-Joseph, 1928). – **As *Prionyx spinolae***: R. Bohart and Menke, 1963:151 (new generic assignment, member of *Prionyx pumilio* species group), 1976:134 (in checklist of world Sphecidae); Sielfeld, 1980b:72 (in checklist of Chilean Sphecidae); Steiner, 1986:97 (references to papers on nesting habits); Amarante, 1993:19 (northeastern Brazil); Chiappa, 2012:8 (Chile: Región de Valparaíso: no specific locality); Danilov, 2020:320 (specimen from Chile in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia).

49. *stschurowskii* (Radoszkowski)

Sphex stschurowskii Radoszkowski, 1877:7, ♀ (as *Stschurowskii*, incorrect original capitalization). Lectotype: Kazakhstan or Uzbekistan: Kyzyl-Kum Desert: no specific locality (ZMMU), designated by Danilov, 2012b:67. – Kohl, 1885b:206 (original description copied); Radoszkowski, 1886a:25 (Turkmenistan); Ed. André, 1888:146 (in revision of Sphecidae of Europe and Algeria), 8* (bibliographic references); Kohl, 1890b:344 (in revision of world Sphecini), 1895:47 (Algeria); Dalla Torre, 1897:442 (in catalog of world Hymenoptera); Morice, 1897:302 (Egypt; description of ♂); Berland, 1926c:200 (specimen in MNHN); Gussakovskij, 1933b:372 (Iran); Honoré, 1944a:72 (in revision of Egyptian Sphecini); de Beaumont, 1960c:170 (Afghanistan; diagnostic characters), 1961e:2 (Iraq), , 1970c:4 (Iran: Baluchistan). – **As *Priononyx stschurowskii***: de Beaumont, 1968b:149 (new generic assignment, member of *Priononyx macula* species group). – **As *Prionyx stschurowskii***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Ebrahimi, 1993:98 (Iran); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia); Dollfuss, 2008b:1415 (Iran: North 65 km south of Tehran); Danilov, 2012a:163 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:67 (bibliographic references, geographic distribution); Dunford, Turbyville, and Leavengood, 2014:11 (listed as medically important in Afghanistan); Ebrahimi, 2014:23 (Iran: Markazi, Semnān, and Tehrān provinces); Augul, Abdul-Rassoul, and Kaddou, 2015:116 (in key to Sphecini of Iraq), 117 (locality records), 119 (illustrations); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:27 (Iran: known from Markazi, Semnan, Sistan-o Baluchestan, and Tehran provinces); Augul, 2019:498 (recorded from Iraq by Derwesh, 1965 as *Chlorion hyalipenne*); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

ssp. *hyalipennis* (Kohl)

Sphex stschurowskii var. *hyalipennis* Kohl, 1895:48, sex not stated. Holotype or syntypes: Algeria: no specific locality (NHMW). – Dalla Torre, 1897:442 (in catalog of world Hymenoptera); Morice, 1911:74 (Algeria: Biskra); Roth, 1925:385 (in revision of North African Sphecini); Berland, 1926c:201 (Algeria, Egypt, Tunisia: locality records, geographic variation); Guiglia, 1937:185 (Libya: Cyrenaica: Rus Hamra), 1942b:229 (Libya); Giner Marí, 1947:19 (Morocco: Western Sahara); Leclercq, 1955h:26 (bibliographic references, faunal records from Africa). – **As *Sphex stschurowskii hyalipennis***: de Beaumont, 1951e:268 (new status, Morocco); de Beaumont and Bytinski-Salz, 1955:42 (Israel); de Beaumont, 1956a:181 (Libya). – **As *Chlorion hyalipenne***: Derwesh, 1965:70 (new generic assignment, new status, Iraq: no specific locality). – **As *Priononyx stschurowskii hyalipennis***: de Beaumont, 1968b:149 (new generic assignment, member of *Priononyx macula* species group). – **As *Prionyx stschurowskii hyalipennis***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Guichard, 1988a:121 (Saudi Arabia); Gadallah and Assery, 2004a:222 (in catalog of Sphecidae of Saudi Arabia); Roche, 2007a:47 (in checklist of Egyptian Sphecidae, redescription), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Gadallah, Al Dhafer, Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia); Ben Khedher, Yildirim, Braham, and Ljubomirov, 2020a:316 (in list of Tunisian Sphecidae *sensu stricto*; additional record: Tunisia: Kairoun); Gadallah, 2020d:86 (in list of aculeate wasps of Arabian Peninsula).

50. *subatratus* (R. Bohart)

Priononyx subatratus R. Bohart, 1958b:90, ♀, ♂ (as *subatrata*, incorrect original termination). Holotype: ♂, USA: California: Inyo County: Deep Springs (CAS). – R. Bohart, 1958b:92, 93 (in key to North American *Prionyx*); F. Parker, 1960:206, 207 (in key to North American *Prionyx*, as *subatrata*). – **As *Prionyx subatratus***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Krombein, 1979b:1586 (in catalog of North American Hymenoptera); Rust, Hanks and Bechtel, 1983:406 (Nevada: Churchill County: Sand Mountain); Amarante, 2002:73 (in catalog of Neotropical Sphecidae); Ruíz Cancino, Coronado Blanco, Varela Fuente, and Horta Vega, 2002:670 (in checklist of Mexican Sphecidae); Dollfuss, 2008b:1415 (Arizona: south of Willcox); Danilov, 2019:78 (member of subgenus *Priononyx*); Danilov and Odintsev, 2023:434 (in subgenus *Priononyx*).

51. *subfuscatus* (Dahlbom)

Sphex subfuscatus Dahlbom, 1845:436, sex not stated (as *subfascata*, incorrect original termination). Holotype or syntypes: "Tauria" = Crimea (lost?). – Eversmann, 1849:369 (Russia: Orenburg Government, lower Volga; Kazakhstan); F. Smith, 1856:242 (in catalog of Hymenoptera in British Museum); Becker, 1880:153 (Russia: Sarepta, now Krasnoarmeysk south of Volgograd); Kohl, 1881:29 (redescription of type material), 1885b:179 (in revision of Palearctic species), 206 (Eversmann's description copied); Gasperini, 1887:18 (recorded from Dalmatia, now Croatia, by Kohl, 1885b); Ed. André, 1888:149 (in revision of Sphecidae of Europe and Algeria), 9* (bibliographic references); Kohl, 1889a:25 (comparison with *Sphex aegyptius*); Kohl and Handlirsch, 1889:275 (Turkmenistan: Chuli); F. Morawitz, 1889a:129 (China: Ordos Region); Kohl, 1890b:354 (in revision of world Sphecini); F. Morawitz, 1891a:202 (Russia: Astrakhan Government), 1893b:407 (Tajikistan: Pyandjikent, Varzaminor); De Stefani Perez, 1894:216 (Italy: Sicilia); Medina, 1894a:260 (Spain); Sickmann, 1894:216 (China: Hopei Province: Tientsin); De Stefani Perez, 1895:226 (in catalog of Sicilian Hymenoptera); Dalla Torre, 1897:442 (in catalog of world Hymenoptera); Mocsáry, 1897:79 (Kingdom of Hungary, some localities are in today's Croatia); Ferton, 1902:504 (nesting habits); Adlerz, 1904:138 (known prey: acridids); Antiga and Bofill, 1904:5 (Spain: Cataluña Province); E. Saunders, 1904c:605 (Spain: Mayorca), 636 (France: Cerbère; Spain); W. Schulz, 1904b:93 (Spain: Murcia, Cuenca; Transcaspia); Ferton, 1905:65 (prey selection), 98 (weight of prey and of female, homing); Mantero, 1905:68 (Italy: Toscana: Isola del Giglio); W. Schulz, 1905b:35 (Algeria: Chellala and Taguin in Alger Province); Dusmet and Mercet, 1906:504, 512 (in key to Spanish Sphecini); Schmiedeknecht, 1907:244 (in key to Hymenoptera of Central Europe); de Gaulle, 1908:104 (in catalog of French Hymenoptera); Ferton, 1910:177 (homing); Mantero, 1911:72 (Italy: Sardegna: Isola dell'Asinara); Morice, 1911:74 (Algeria: Biskra); Kohl, 1913b:15 (Russia: Voronezh Oblast: Valtuyki at 50°14'N 38°08'E); R. Turner, 1914b:250 (India: Tamil Nadu: Coimbatore); Dusmet y Alonso, 1915:86 (Spain: Aragón); Strand, 1915:90 (Russia: Sarepta, now Krasnoarmeysk south of Volgograd), 1916b:108 (China: Tsingtau, now Qingdao); Ferton, 1921:350 (prey paralysis incomplete); Fahringer, 1922:178 (Turkey); Maidl, 1922:67 (Croatia); Ferton, 1923:107 (prey capture and paralyzing, malaxation, nest digging, oviposition, nest closure, description of larva), 155 (incomplete paralysis of prey), 296 and 307 (easily finds nest when carrying prey), 320 (captures *Oedipoda coerulescens* (L.), Acrididae, when preferred *Caloptenus italicus* (L.), Acrididae, is in short supply); Gribodo, 1924b:49 (Libya: Apollonia); Berland, 1925d:38 (in Sphecid Fauna of France); Roth, 1925:389 (in revision of North African Sphecini); Berland, 1926c:200 (miscellaneous locality records); Coulon, 1925:116 (France: Sète; Spain: Montario); von Schulthess, 1926b:209 (Libya); Schmiedeknecht, 1930:705 (in keys to Hymenoptera of North and Central Europe); Dusmet y Alonso, 1931:7 (Portugal: Soure); Guiglia, 1932:125 (Ethiopia: Harer area; northeastern Kenya); Bischoff, 1933:5 (Morocco); Masi, 1933:197 (Italy: Toscana: Isola di Capraia); Giner Marí, 1934:130 (Spain); Grandi, 1934:130 (Italy: Lazio: Acilia); Guiglia, 1934b:293 (Libya: bibliography and summary of locality records); Gussakovskij, 1934a:3 (China: Kansu Province); Nadig, 1934:34 (France: Corse: Calanches; Italy: Sardegna: Alghero, Aritzo, Macomer); Bernard, 1935:61 (France: Var: Fréjus area); Piel, 1935:296 (nesting habits); Yasumatsu, 1935a:8, 22 (China: Jehol Region), 1938:93 (in revision of East Asian Sphecini; Korea, Manchuria); Balthasar, 1941a:105 (Czech Republic: Bzenecko area); Guiglia, 1942a:59 (Greece: Island of Rhodes: Villanova); Yasumatsu, 1942c:105 (China: Beijing); Giner Marí, 1943a:82 (in Sphecid Fauna of Spain); Guiglia, 1943d:91 (Albania: Scutari), 1944b:7 (Italy); Honoré, 1944a:70 (in revision of Egyptian Sphecini); Deleurance, 1946b:62 (prey), 67 (France: Bouche-du-Rhône: Camargue: Bois des Rièges); de Beaumont, 1947b:383 (Cyprus); Zavadil in Zavadil and Šnoflák, 1948:167 (in key to Sphecidae of Czechoslovakia); de Andrade, 1949:10 (Portugal); Berland and Bernard, 1949:3 (in revision of French *Sphex sensu lato*), 9 (review of biological data); Pittioni, 1950:20 (Cyprus); de Beaumont, 1951e:268 (Morocco); Cleu, 1953:50 (France: Ardèche River basin); de Beaumont, 1953h:195 (type material not in Lund); Nouvel and Ribaut, 1953:177 (France: Haute-Garonne: Saint-Béat); Grandi, 1954:236 (Italy); de Beaumont and Bytinski-Salz, 1955:41 (Israel); Harant and Leclercq, 1955:250 (France: Hérault: Bionne, Fontcaude, Paillade); Leclercq, 1955h:25 (bibliographic references, faunal records from Africa); Vergne, 1955:4 (France: Auvergne); Berland, 1956:1170 (in revision of African Sphecini); Ceballos, 1956:364 (in catalog of Hymenoptera of Spain); Morel, Nouvel, and Ribaut, 1956:337 (France: Département des Pyrénées-Orientales); Bajári, 1957a:8, 10 (in key to Hungarian Sphecidae); Nouvel and Ribaut, 1958:9 (France: Pyrénées-Orientales: Banyuls-sur-Mer); de Beaumont, 1959a:10 (Italy);

Diniz, 1959:27 (Portugal: five localities); Scobiola-Palade, 1959:496 (first record from Romania: Constanța Region: Agigea, description and illustration of male genitalia); de Beaumont, 1960a:5 (Greece: Island of Rhodes); Noskiewicz and Puławski, 1960:41 (in key to Polish Sphecidae); Scobiola, 1960b:232 (Romania: Medgidia Region: Valul lui Traian); Kocourek, 1963:295 (Czech Republic: Moravy: Lidéřovice; Slovakia: Šturovo); Myartseva, 1963b:59 (Turkmenistan: lower Murgab River); Ceballos, 1964:88 (in supplement to catalog of Spanish Sphecidae); Myartseva, 1964a:74 (nesting habits in Turkmenistan); de Beaumont, 1965a:65 (Greece: Crete: Heraklion); Isensee, Lesemann, and Röseler, 1965:612 (Spain: Gerona: no specific locality); Myartseva, 1965:84 (Turkmenistan: Akibay and Sakar-Chagin districts); Suárez, 1969:53 (Spain: Almería Province); Scobiola-Palade, 1960b:232 (Romania); de Beaumont, 1962b:19 (Spain); Scobiola-Palade, 1963:825 (Romania); Tsuneki, 1963b:52 (nesting habits); de Beaumont, 1965a:13 (Greece); Scobiola-Palade, 1966a:162 (Romania: Tulcea District: C.A. Rosetti, Periprava); de Beaumont, 1967a:273 (Turkey); Tsuneki, 1967e:2 (China: Manchuria); de Beaumont, 1968b:149 (member of *Sphex subfuscatus* species group); Scobiola-Palade, 1968b:141 (Romania: Island of Letea in delta of Danube); Tsuneki, 1968l:50 (South Korea: Island of Quelpart, now Island of Cheju); Kazenas, 1969a:22 (Kazakhstan: Ili River, Sharyn' River, Karatal River, Zailiyskiy Alatau); Tsuneki, 1971m:2 (China: Beijing: Tiendang); Balthasar, 1972:424 (in Sphecid Fauna of Czechoslovakia); Kazenas, 1972b:113 (Kazakhstan), 1974b:109 (feeding on flowers of *Tamarix* sp., Tamaricaceae, in Kazakhstan); Georghiou, 1977:192 (Cyprus); Kazenas, 1978b:41 (in key to Sphecidae of Kazakhstan and Central Asia); Pulawski, 1978:184 (in key to Sphecidae of European part of USSR); Radović and Krnić, 1979:unpaginated foldout (nesting in sand, foreleg structure); Scobiola-Palade, 1985:95 (Romania: delta of Danube); Meyer-Holzappel, 1986:100 (nest parasite: *Hilarella stictica* Meigen, Sarcophagidae); Piek and Spanjer, 1986:190 (in list of Sphecidae with known prey); Padr in Šedivý, 1989a:166 (in checklist of Czechoslovakian Sphecidae); Blagoveshchenskaya, 1994:88, 89 (Russia: Ul'yanovsk Oblast', as *subfasciatus* and *subfuscus*, respectively); Nagase, 2006b:2 (specimens collected by First Scientific Expedition to Manchoukou, 1933, now part of eastern China); Alieva and Humbatov, 2007:77 (nesting and prey, from literature). – **As *Harpactopus subfuscatus***: Radoszkowski, 1892:586 (new generic assignment, description of male genitalia), Vayssière, 1921:132 (nesting habits). – **As *Chlorion subfuscum***: Bischoff, 1931:8 (new generic assignment, Spain). – **As *Prionyx subfuscatus***: Diniz, 1965:4 (new generic assignment, Portugal: Lisboa, Massorra, Sines, Vale de Gaio); Myartseva, 1966:48 (preying on orthopterans), 1972a:84 (Turkmenistan); R. Bohart and Menke, 1976:134 (in checklist of world Sphecidae); Guichard, 1978:270 (first record from Greece: Kalambaka); Valetta, 1979:215 (Malta); Guichard, 1980:224 (Oman); Kazenas, 1980e:81 (Russia: Far East); Pagliano, 1980:110 (Italy: Liguria); Roche, 1981:1 (in checklist of Sphecidae of United Arab Emirates); Tsuneki, 1982b:14 (known from Korea); Mingo and Gayubo, 1983:153 (Spain); Gayubo and Tormos, 1984:10 (Spain: Valencia); Pagliano, 1984:366 (Italy); Brockmann, 1985b:312 (nest closure summary); Chevin and Chevin, 1985:38 (France: Aude); Pagliano, 1985:9 (Italy); Paik, 1985:197 (in list of Sphecidae of Korea); Radović, 1985:64 (sting apparatus analyzed); Gayubo and Tormos, 1986b:4 (Spain: Valencia); Islamov, 1986:516 (Uzbekistan: Tashkent Oblast'); Józán, 1986:367 (Hungary: Kiskunság National Park); Nemkov, 1986:92 (Russian: Siberia: Irkutsk Oblast'); Piek and Spanjer, 1986:188 (in list of Sphecidae with known prey); Steiner, 1986:97 (references to papers on nesting habits); Gayubo, 1987:107 (Spain: Provincia de Ciudad Real); Tormos and Jiménez, 1987a:122 (Spain: Valencia), 1987b:316 (Spain: Valencia Province: Dehesa de El Saler); Guichard, 1988a:121 (Arabian Peninsula); Gayubo, Asís, and Tormos, 1990a:10 (Spain); Pagliano, 1990:58 (in catalog of Italian Sphecidae); Dollfuss, 1991:29 (in key to Sphecidae of North and Central Europe); Gayubo, Borsato, and Osella, 1991:394 (Italy: Lazio, Sicilia); Gayubo and Torres, 1991:Table I (Spain: Salamanca; effects of urban pressure); Hamon, Fonfria, and Tussac, 1991:128 and 129 (in key to French Sphecini), 133 (in France north to Rhône and Loire-Atlantique Departments); Kazenas and Nasyrova, 1991:38 (Kazakhstan: preying on *Caliptamus barbarus turanicus* Serg. Tarb. and *C. barbarus cephalotes* F.W., Acrididae); Leclercq, 1991a:274 (omitted from Leclercq's, 1979, catalog of France and Benelux Sphecidae); Schembri, 1991:176 (Malta); Kazenas and Tobias, 1992:29 (sleeping aggregations); Mochi and Luchetti, 1993:104 (France: Corse); Gayubo and Borsato, 1994:202 (Italy: Sardegna); Roche and Zalut, 1994:114 (Egypt: Sinai Peninsula); Tormos, Asís, and Gayubo, 1994:187, 195 (Spain: Albacete Province); Kazenas in Nemkov, Kazenas, Budrys, and Antropov, 1995:385 (in key to Sphecidae of Russian Far East); Negrisolo in Minelli, Ruffo, and La Posta, 1995b:2 (in catalog of Italian fauna); Pagliano and Scaramozzino, 1995:730 (Italy: Island

of Lampedusa); Scharrer, 1995:22 (Croatia: Island of Krk); Minoranskiy and Shkuratov, 1996:81 (Russia: Rostov Oblast'); Wu and Zhou, 1996a:45 (in revision in Economic Insect Fauna of China); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:63 (in Sphecidae Fauna of Western Europe); Schmidt and Schmid-Egger, 1997:26 (the only German record is by Ruthe and Stein, 1857; apparently does not occur in Germany); Kazenas, 1998b:125 (in Sphecidae Fauna of Kazakhstan); Esenbekova and Kazenas, 2000:10 (southeastern Kazakhstan: near Bakanas, 60 km east of Chilik, near Kapchagay, 45 km northwest of Suzak); Vicidomini, 2000b:27 (Italy: previously recorded from Campania); Kazenas, 2001b:15 (in checklist of Sphecidae of Kazakhstan and Central Asia), 85 (nest and prey); Kazenas and Esenbekova, 2001:133 (Kazakhstan: Almatinskiy Nature Reserve); Ohl et al., 2001:142 (recorded from Germany but not occurring there); Kazenas, 2002a:30 (geographic distribution, collecting localities in Kazakhstan); Shkuratov, 2002a:383 (Russia: Rostov Oblast'), 2002b:139 (Russia: Rostov Oblast': Rostovskiy Nature Reserve at 46°27'N 42°41'E); Generani, Pagliano, Scaramozzino, and Strumia, 2003:64 (Italy: Arcipelago Toscano); S. Gess and F. Gess, 2003:92 (Namibia and South Africa: visiting flowers of *Athanasia trifurcata* (L.) L., Asteraceae, and *Zygophyllum simplex* L., Zygophyllaceae); Protsenko, 2003:68, 69 (first record from Ukraine: Odessa Oblast': island of Malyi Tataru in Danube delta at 45°21'N 29.00'E); Schmid-Egger, 2003:757 (Italy: Sicilia: Bronte, Linguaglossa); Wu, Zhou, Q. Li, and Yang, 2003:808 (China: Fujian Province); Gadallah and Assery, 2004a:217 (in key to Sphecidae of Jeddah Region, Saudi Arabia), 222 (in catalog of Sphecidae of Saudi Arabia); Kazenas, 2004b:99 (Kazakhstan: western Tien Shan Mountains), 2004d:27 (Kazakhstan: northern Caspian region); Q. Li and He, 2004:1127 (in hymenopterous fauna of Zhejiang Province, China); Shkuratov, 2004a:73 (Russia: Rostov Oblast'); Skibińska in Bogdanowicz, Chudzicka, Pilipiuk, and Skibińska, 2004:358 (in catalog of Polish Sphecidae); Wiśniowski, 2004:38 and 58 (in checklist of Polish Sphecidae); Gayubo and Özbek, 2005:9 (Turkey: Antalya: Arapsuyu; Erzurum: Dumlü; Kars: Sarýkamýp); Gülmez and Tüzün, 2005:48 (Turkey: Ankara Province); Pagliano and Negrisola, 2005:54 (in Sphecidae Fauna of Italy); Shoreenko, 2005a:162 (Ukraine: Crimea), 2005b:97 (Ukraine: Crimea: Karadagh Nature Reserve); Straka, 2005a:396 (extinct in Czech Republic); Hua, 2006:276 (in list of Chinese insects, geographic distribution); Magdalou, 2006b:109 (France: Pyrénées-Orientales: Mas-Larrieu Nature Reserve near Argelès-sur-Mer), 2006c:7 (France: Pyrénées-Orientales: Réserve Naturelle du Mas-Larrieu); Standfuss and Standfuss, 2006c:307 (Greece: Thessalia: Magnisia Peninsula at 39°N 23°E); Jacobs, 2007:42 (in key to Sphecidae of Germany, not yet found in Germany); Kazenas, 2007a:89 (Kazakhstan: Akmala Oblast': Kurgandzhin Nature Reserve and vicinity); Roche, 2007a:36 (in checklist of Egyptian Sphecidae, redescription, as *subfuscatus subfuscatus*), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae, as *subfuscatus subfuscatus*); Vepřek and Straka, 2007:199 (in catalog of Sphecidae of Czech Republic and Slovakia, known from Moravia and Slovakia only); Dollfuss, 2008b:1415 (locality records from Croatia, Greece, Italy, Kazakhstan, Kyrgyzstan, Mongolia, Russia, South Africa, Tajikistan, Turkmenistan, and Uzbekistan); Kazenas, 2008c:255 (Kazakhstan: village Koktum south of Lake Alakol'); Ljubomirov and Yildirim, 2008:30 (in catalog of Sphecidae of Turkey); Nemkov, 2008b:17 (in catalog of Sphecidae of Asiatic Russia); Baños-Picón, Asís, Gayubo, and Tormos, 2009:310 (Spain: frequency of specimens collected with hand nets and Malaise traps); Danilov, 2009:55 (Russia: Western Siberia: Kulundinskaya Steppe); Nemkov, 2009b:46 (in new catalog of Sphecidae and Crabronidae of Asiatic Russia), 2009c:235 (Russia: Primorskiy Kray: Lazovsky Nature Reserve); Pagliano, 2009:175 (Italy: Piemonte: San Benedetto Belbo); Shoreenko, 2009:366 (in list of Sphecidae *sensu lato* of Crimea); Bitsch, 2010:105 (in supplement to vol. II of Faune de France, 1997: France: first records from Lozère: Saint-Etienne-Vallée-Française; specimens from Loiret: Olivet are *Isodontia mexicana*); Burguet in Durand and Burguet, 2010:22 (France Département d'Ardèche: Ardèche Gorges); Danilov, 2010b:44 (distribution of Palearctic-Ethiopian type); Rudoiskatel', 2010:147 (Russia: southern Ural Mountains); Sakenin, Samin, and Bagriacik, 2010:17 (Iran: Khuzestan: Abadan); van der Smissen, 2010b:386 (France: Ardèche: Vogüé; Vaucluse: Bédoin); Baghirov, 2011b:140 (Russia: Altayskiy Kray: Savvushka); Cruz-Sánchez, Asís, Gayubo, Tormos, and González, 2011:497 (Spain: Salamanca and Zamora provinces: Arribes del Duero Natural Park: effects of wildfire), 501 (exclusive species of burned areas); Józán, 2011:179 (in checklist of Sphecidae *sensu lato* of Hungary); Danilov, 2012a:163, 164 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:67 (bibliographic references, geographic distribution); Kazenas, 2012b:168 (Kazakhstan: Korgalzhin State Nature Reserve); Nemkov, 2012c:434 (in catalog of Sphecidae of Russian Far East); Prisniy, 2012:46

(Russia: Belgorod Oblast', as *subfuscatus*); Protsenko, Fateryga, Ivanov, and Puzanov, 2012:58 (Ukraine: Crimea); Gadallah, Al Dhafer, Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia); Kazenas, 2013a:23 (color photograph of female, short information on geographic distribution and nesting habits); Baldock, 2014:354 (Spain: Island of Mallorca); Danilov, 2014a:424 (Russia: Siberia: Altayskiy Kray, Buryatia, Tyva), 2014b:514 (in key to Sphecidae s.s. of Siberia); Ebrahimi, 2014:24 (Iran: Ilām, Kermān, Kermānshāh, and Māzandarān provinces); Kazenas, 2014a:132 (Kazakhstan: Karatau Mountain Range); J.-K. Kim, 2014:419 (in catalog of Sphecidae *sensu lato* of Korean Peninsula); Kim, Yeo, and Kim, 2014:286 (in revision of Sphecidae *sensu stricto* of South Korea); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces, as *subfuscatus subfuscatus*); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey); Shorenko, 2015:317 (in list of Sphecidae *sensu lato* of Crimea); Levárdá and Matache, 2016:43 (in catalog of Sphecidae s.s. of Romania); Mokrousov and Popov, 2016:564 (Russia: Abkhazia, Krasnodarskiy Kray); Arens, 2017a:630 (Greece: Peloponnesus); Danilov, 2017b:215 (in catalog of Sphecidae s.s. of Russia); Danilov and Mokrousov, 2017a:109 (Russia: Dagestan, Kalmykia); Di Giovanni, Mei, and Cerretti, 2017:73 (Italy: Lombardia: Riserva Naturale Bosco della Fontana 5 km from Mantova at 45°12'N 10°44'E, found in canopy); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:27 (Iran: known from Ilam, Kerman, Kermanshah, Khuzestan, and Mazandaran provinces); Magdalou, 2017:16 (France: Pyrénées-Orientales: Réserves Naturelles Catalanes: Mas Larrieu); Shorenko, 2017:76 (in Crimea collected in June through September), 2018:127 (Crimea, including localities, habitats, and number of specimens); Augul, 2019:498 (recorded from Iraq by El-Haidari, Fattah, and Sultan, 1971 as *Sphex subfuscatus*); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Can and Gülmez, 2019:349 (in key to *Prionyx* of Turkey); Gülmez, 2019:3 (Turkey: Ankara Province: no specific locality); Bitsch, Barbier, and Jacobs *in* Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:141 (in Sphecid Fauna of Europe); Cassar and Mifsud, 2020:164 (in checklist of Sphecidae s.s. of Malta); Danilov, 2020:320 (specimens from Kazakhstan, Kyrgyzstan, Tajikistan, and Russia: south of European part and Siberia in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Gadallah, 2020d:86 (in list of aculeate wasps of Arabian Peninsula); Shorenko, 2020:47 (Crimea: Karadag Nature Reserve); Turrisi, Altadonna, Lo Cascio, Nobile, and Selis, 2020:727 (Italy: Aeolian Archipelago: island of Lipari); Cross, Baldock, and Wood, 2021:18 (in catalog of Sphecidae *sensu lato* of Portugal); Olszewski, Wiśniowski, and Ljubomirov, 2021:103 (in commented list of Sphecidae *sensu lato* of Poland); Embergenov, Élmurodova, Amirov and Kimyonazarov, 2022:47 (Uzbekistan: Khiva area); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey); Kochetkov, 2023:191 (Russia: Amur Oblast': Khingan Nature Reserve).

Sphex soror Dahlbom, 1845:436, sex not stated. Holotype: ♀, Greece: Island of Rhodes (Stockholm, coll. Hedenborg). Synonymized with *Sphex aegyptius* by Kohl, 1885b:181, and with *Sphex subfuscatus* by de Beaumont, 1949a:127 (holotype and paratype mentioned). – F. Smith, 1856:243 (in catalog of Hymenoptera in British Museum); Sichel, 1861:751 (Italy: Sicilia); Sajó, 1882:5 (Hungary); nec Honoré, 1944a:69 (= *Prionyx crudelis*); Balthasar, 1954b:281 (Palestine: Tabgha at Lake Tiberias); Casolari and Casolari Moreno, 1980:102 (specimens in M. Spinola collection, Torino); Pagliano, 2008:524, 525 (specimens in M. Spinola collection, Torino).

Sphex nigrinus Lucas, 1849:271, sex not stated (as *nigrita*, incorrect original termination), junior primary homonym of *Sphex nigrinus* Fabricius, 1781 (now in *Java*, a genus of Pompilidae), and of *Sphex nigrinus* Turton, 1802:484 (which is a lapsus or emendation of *Sphex nigrata* Gmelin, 1790:2723, a European evaniid). Holotype or syntypes: Algeria: La Calle area, now El Kala (MNHN). Synonymized with *Sphex subfuscatus* by Kohl, 1885b:179. – F. Smith, 1856:244 (in catalog of Hymenoptera in British Museum).

Sphex desertorum Eversmann, 1849:368, ♀, ♂. Syntypes: Russia: Astrakhan, Orenburg, and Saratov provinces; and Kazakhstan (ZIN). Synonymized with *Sphex subfuscatus* by Kohl, 1885b:179. – Becker, 1880:153 (Russia: Sarepta, now Krasnoarmeysk south of Volgograd); Radoszkowski, 1881:209 (Angola), 1886a:25 (Turkmenistan) and 26 (description of male genitalia).

Enodia chrysoptera Ruthe and Stein, 1857:312, ♀. Holotype: ♀, Germany: Berlin area (ZMHU). Synonymized with *Sphex subfuscatus* by Kohl, 1885b:179. – Schirmer, 1912:168 (Germany: found in Berlin area by Ruthe and Stein, 1857). – As

Sphex chrysoptera: Myartseva, 1971b:180 (new generic assignment, ranges mainly outside Palearctic Region); Casolari and Casolari Moreno, 1980:102 (specimens in M. Spinola collection, Torino). – ***As Parasphex chrysoptera***: Kirchner, 1867:217 (new generic assignment, in catalog of European Hymenoptera).

Gastrosphaeria anthracina A. Costa, 1858b:10. ♀, ♂. Syntypes: Italy: various localities (Napoli). Synonymized with *Sphex subfuscatus* by Kohl, 1885b:179. – Kirchner, 1867:217 (in catalog of European Hymenoptera); Marquet, 1881:178 (southern France); De Stefani Perez, 1882:38 (Italy: Sicilia: Sciacca). – ***As Sphex anthracina***: A. Costa, 1867b:70 and 1867c:14 (new generic assignment, in revision of Italian Sphecidae); Palma, 1869:38 (Italy: Sicilia settentrionale); A. Costa, 1882b:22 (Italy: Sardegna); Casolari and Casolari Moreno, 1980:102 (specimens in M. Spinola collection, Torino); Pagliano, 2008:524 (specimens in M. Spinola collection, Torino).

? *Sphex namkumiensis* Laidlaw, 1929:232, ♀. Holotype: ♀, India: Bihar: Namkum (Royal Scottish Mus., Edinburgh). – van der Vecht, 1957b:22 (redescription of holotype, misspelled as *nankumiensis*); R. Bohart and Menke, 1976:134 (as tentative synonym of *Prionyx subfuscatus*, in checklist of world Sphecidae).

ssp. albovillosulus (Giordani Soika)

Sphex subfuscatus albovillosulus Giordani Soika, 1942:198, ♂. Syntypes: Somalia: Ischia Baidoa (depository?). – ***As Prionyx subfuscatus albovillosulus***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae).

ssp. rhodesianum (Arnold)

Chlorion subfuscum race *rhodesianum* Arnold, 1936:28, ♀, ♂. Syntypes: Zimbabwe: Matetsi (SAM). – ***As Prionyx subfuscatus ? ssp. rhodesianus***: R. Bohart and Menke, 1976:134 (new generic assignment, status tentative, in checklist of world Sphecidae).

ssp. rukwaensis (Arnold)

Sphex rukwaensis Arnold, 1959:325, ♂. Holotype: ♂, Tanzania: Ukia in Rukwa Valley (BMNH). – ***As Prionyx subfuscatus ? rukwaensis***: R. Bohart and Menke, 1976:134 (new generic assignment, status tentative, in checklist of world Sphecidae).

52. sundewalli (Dahlbom)

Enodia sundewalli Dahlbom, 1845:439, sex not stated. Holotype or syntypes: South Africa: KwaZulu-Natal: Port Natal, now Durban (depository?) – Kohl, 1890b:453 (original description copied); Dalla Torre, 1897:443 (in catalog of world Hymenoptera, as *sundevalii*); W. Schulz, 1912:93 (type material neither in Lund nor in Berlin); Leclercq, 1955h:38 (species incertae sedis). – ***As Prionyx sundewalli***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Danilov and Odintsev, 2023:435 (tentatively in subgenus *Prionyx*).

53. thomae (Fabricius)

Sphex thomae Fabricius, 1775:346, sex not stated. Lectotype: ♂, U.S. Virgin Islands: Island of Saint Thomas (ZMUC), designated by van der Vecht, 1961a:35. – Fabricius, 1781:443 (redescription), 1787:274 (redescription); Gmelin, 1790:2725 (redescription); Christ, 1791:307 (redescription); Fabricius, 1793:199 (redescription), 1796:156 (in Index to his Entomologia Systematica, 1793); Turton, 1801:484 (redescription); Jurine, 1807:129 (in list of *Sphex*); Cresson, 1963:320 (in catalog of North American Hymenoptera), 1868:379 (New Mexico); Cameron, 1889a:36 (summary of distribution records); Kohl, 1890b:358 (in revision of world Sphecini); W. Fox, 1891d:342 (Jamaica); Radoszkowski, 1893a:58 (Turkmenistan, obviously in error); W. Fox, 1895c:266 (Mexico: Baja California Sur); Dalla Torre, 1897:443 (in catalog of world Hymenoptera); W. Fox, 1897b:378 (Brazil); Ducke, 1901:241 (Brazil: Pará: Belém); Schrottky, 1903b:123 (in checklist of Hymenoptera of Argentina, Paraguay, and Uruguay); Ducke, 1908b:82 (Brazil: Ceará State); Strand, 1910a:133 (Paraguay), 1911a:151 (Ecuador); Jörgensen, 1912:286 (Argentina: Mendoza Province); Bodkin, 1918:315 (British Guiana, nesting behavior); Campos, 1922:68 (Ecuador: Durán); Berland, 1926c:204 (miscellaneous locality records); G. Carpenter, 1930b:293 (nest closure); Liebermann, 1931:24 (in revision of Argentinean Sphecini); Bischoff and von Schulthess, 1937:168 (Argentina); Murray in Muesebeck, Krombein, and Townes, 1951:973 (in catalog of North American Hymenoptera); Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino); Pagliano, 2008:531, 532 (specimens in M. Spinola collection, Torino). – ***As Pepsis thomae***: Fabricius, 1804:209 (new generic

assignment, redescription). – **As *Harpactopus thomae***: Ashmead, 1900:229 (new generic assignment, Lesser Antilles: Island of Saint Vincent), 308 (in checklist of Caribbean Hymenoptera). – **As *Chlorion thomae***: Fernald, 1906:342 (new generic assignment, in revision of Sphecini of North America and West Indies), 1907:264 (Argentina); H. Smith, 1908b:333 (in revision of Nebraskan Sphecidae); J. Smith, 1910:677 (in new list of insects of New Jersey); Mickel, 1918b:398 (in catalog of Nebraskan Sphecidae); Fernald, 1931a:441 (study of type series), 1943a:287 (recorded from Florida, but all specimens seen by author are *pubidorsum*); Willink, 1948a:314 (history of South American records, presence in Argentina confirmed), 318 (differences between *Chlorion thomae* and *Ch. striatulus*), 320 (in key to *Chlorion thomae* species group), 1951:186 (in revision of Argentinean Sphecini); Bianchi, 1954:287 (Hawaiian Islands: Oahu: Kailua); Krombein, 1958f:191 (in supplement to catalog of North American Hymenoptera: description of larva by Evans and Lin, 1956a, reported); Yoshimoto, 1960:335 (Hawaiian Islands). – **As *Priononyx thomae***: Dahlbom, 1843:28 (new generic assignment, in revision of Sphecidae and Pompilidae), 1845:439 (in key); F. Smith, 1856:265 (in catalog of Hymenoptera in British Museum); A. Costa, 1864b:112 (two specimens from La Plata in Museo Zoologico di Napoli); Cresson, 1865a:137 (Cuba), 1865b:464 (specimens in ANSP collection); de Saussure, 1867:43 (variation); Cresson, 1868:379 (New Mexico); Taschenberg, 1869:409 (Brazil); Burmeister, 1872:239 (Argentina; Brazil: Neu-Freiburg); Cresson, 1873:213 (Texas), 1875:715 (Arizona, Nevada, New Mexico); F. Lynch Arribálzaga, 1878:328 (Argentina: Buenos Aires area); Patton, 1880a:384 (characteristics of the genus *Priononyx*); Dewitz, 1881:203 (Puerto Rico); Holmberg, 1884:226 (Uruguay); Cresson, 1887:276 (in catalog of North American Hymenoptera); Ashmead, 1890:33 (in checklist of Hymenoptera of Colorado); W. Fox, 1891c:342 (Jamaica); C. Robertson, 1892:107 (visiting flowers of *Pycnanthemum linifolium* Ph., Lamiaceae), 1894:455 (visiting flowers of *Solidago missouriensis* Nutt., Asteraceae), 456 (visiting flowers of *S. canadensis* Linnaeus, Asteraceae), 467 (visiting flowers of *Rudbeckia hirta* Linnaeus, Asteraceae), 469 (visiting flowers of *Lepachys pinnata* Torr. and Gray, Asteraceae), 472 (visiting flowers of *Coreopsis palmata* Nutt., Asteraceae), 1896:73 (visiting flowers of *Polygonum hydropiperoides* Michx., Polygonaceae); Ashmead, 1899d:353 (in checklist of North American Sphecidae); J. Smith, 1900:523 (in list of insects of New Jersey); Hartman, 1905:62 (nesting habits); F. Williams, 1914b:227 (nesting habits); Holland, 1917:294 (Cuba: Isla de Pinos, now Isla de la Juventud: Nueva Gerona); PH. Rau and N. Rau, 1918:175 (nest digging, prey: *Dissosteira carolina* (Linnaeus), Acrididae, prey transportation); G. Carpenter, 1930b:294 (nest closure); Rau, 1933:283 (Panama: Island of Barro Colorado); Richards, 1937a:101 (Guyana); Strickland, 1947:128 (Canada: Alberta: Medicine Hat); Wolcott, 1951:840 (in review of insects of Puerto Rico); Evans and Lin, 1956a:142 (description of larva); R. Bohart, 1958b:92, 93 (in key to North American *Prionyx*); Evans, 1958a:181 (nesting behavior); F. Parker, 1960:206 (in key to North American *Prionyx*). – **As *Prionyx thomae***: R. Bohart and Menke, 1963:159 (new generic assignment, in revision of Nearctic Sphecini); Lavigne and Pfadt, 1966:31 (Wyoming: preying on three grasshopper species); Alayo Dalmau, 1973:185 (in catalog of Cuban Hymenoptera), 1976:27 (in checklist of Cuban Sphecidae); R. Bohart and Menke, 1976:134 (in checklist of world Sphecidae); Kumar, Lavigne, Lloyd, and Pfadt, 1976:51 (USA: Colorado: Pawnee National Grassland); Elliott, Kurczewski, Claflin, and Salbert, 1979:357 (Bahama Islands: San Salvador Island); Heithaus, 1979:193 (Costa Rica: Guanacaste: Hacienda las Ciruelas 30.5 km northwest of Cañas); Krombein, 1979b:1586 (in catalog of North American Hymenoptera); Nascimento and Overal, 1980:8 (Brazil); de Zayas, 1981:78 (Cuba); Sielfeld, 1980b:72 (in checklist of Chilean Sphecidae); Grissell, 1981:16 (unusual nesting behavior: use of preexisting cavities); Brockmann, 1985b:312 (nest closure summary); Rust, Menke, and Miller, 1985:46 (California: Channel Islands); Parks, 1986:34 (California: Torrey Pines State Reserve); Piek and Spanjer, 1986:188 (in list of Sphecidae with known prey); Steiner, 1986:97 (references to papers on nesting habits); Ch. Porter, 1987:43 (Chile: Tarapacá Region); Yústiz, 1987:13 (Venezuela: Central Lara Depression); Spofford, Kurczewski, and Downes, 1989:258 (summary of previous records for nest parasite *Senotainia rubriventris* Macquart, a miltogrammine fly); Maes, 1989:92 (in catalog of Nicaraguan Sphecidae); Callan, 1990b:19 (in checklist of Trinidad Sphecidae); Elliott, 1992:45 (in list of wasps of Bahama Islands: occurring on Long Island and San Salvador); Snelling, 1992:14 (Virgin Islands: Island of Mona); Amarante, 1993:19 (northeastern Brazil); Snelling, 1993:18 (British Virgin Islands: Island of Guana), 19 (same: Island of Mona); Ahlstrom, 1995:107 (in checklist of insects of North Carolina); Hanson and Menke, 1995:637 (known from Costa Rica); Lecoq and Pierozzi, 1996:515 (preying on acridid *Rhammatocerus schistocercoides* (Rehn, 1906),

- Acrididae, in Brazil: Mato Grosso); Weissmann and Kondratieff, 1999:78 (Colorado: Great Sand Dunes National Monument); Fernández, 2000:142 (Colombia); Amarante, 2002:73 (in catalog of Neotropical Sphecidae); Fernández, Saríol, Vega, Ricardo, González, and Portuondo, 2002:46 (Cuba: Provincia Granma); Ruíz Cancino, Coronado Blanco, Varela Fuente, and Horta Vega, 2002:670 (in checklist of Mexican Sphecidae); Starr and Hook, 2003:22 (in catalog of Aculeata of Trinidad, West Indies); Portuondo and Fernández, 2004:135 (Cuba: Sierra Maestra and Nipe-Sagua-Baracoa mountains); Snelling, 2005:291 (British Virgin Islands: Island of Guana); Genaro, 2006:51 (in catalog of Cuban Sphecidae and Crabronidae; other countries: North America, Mexico, Nicaragua, Panama, Isla de Juventud, Little Cayman, Bahamas, Jamaica, Hispaniola, Mona Island, Puerto Rico, Island of Guana, Island of Saint Thomas, Saint Vincent, Trinidad, Guyana, Venezuela, Colombia, Ecuador, Chile, Uruguay, Paraguay, Brazil, Argentina); Horta Vega, Pinson Domínguez, Barrientos Lozano, and Correa Sandoval, 2007:48 (Mexico: Tamaulipas); Dollfuss, 2008b:1416 (locality records from Ecuador, French Guyana, and Mexico); Perez-Gelabert, 2008:241 (in list of arthropods of island of Hispaniola); Buys, 2009c:311 (nesting behavior, male behavior, larval development), 2009e:277 (Brazil: Rio de Janeiro: Duque de Caxias, Macaé, Maricá, Itatiaia, Rio de Janeiro, Reserva Biológica de Poço das Antas, Seropédica); Rasmussen and Asenjo, 2009:16 (in checklist of Crabronidae of Peru); Buys, 2011b:2 (Brazil: Rio de Janeiro: Cabo Frio, Rio de Janeiro, Seropédica), 2011c:225 (description of mature larva); Chiappa, 2012:8 (Chile: Región de Valparaíso: no specific locality); Rodrigues and Buys, 2013:214 (Brazil: Espírito Santo: Santa Teresa and Vila Velha); Buys and Rodrigues, 2014:41 (Brazil: State of Espírito Santo: several localities); Silvestre, Demétrio, Trad, de Oliveira Lima, Auko, and de Souza, 2014:70 (Brazil: Mato Grosso do Sul: dry forests in Bodoquena Mountain Range and Brazilian Chaco); Trad and Silvestre, 2017:4 (Brazil: Mato Grosso do Sul); Danilov, 2019:78 (member of subgenus *Priononyx*); Vanoye-Eligio, Meléndez-Ramírez, Ayala-Barajas, Delfin-González, and Horta-Vega, 2019:7 (Mexico: Yucatán); Buys, 2020b:80 (in cladistic analysis of larvae of Sphecidae s.s.); Danilov and Odintsev, 2023:434 (in subgenus *Priononyx*); Buck and Bennett, 2025:224 (in checklist of Sphecidae sensu lato of Canada, Alaska, and Greenland: Canada: Alberta, first record from Saskatchewan: no specific locality).
- Pepsis crucis* Fabricius, 1804:209, sex not stated. Lectotype: ♀, South American Islands: no specific locality (ZMUC), designated by R. Bohart and Menke, 1963:159. Synonymized with *Priononyx thomae* by Dahlbom, 1845:XXI. – **As *Sphex crucis***: Jurine, 1807:129 (in list of world *Sphex*); F. Smith, 1856:259 (in catalog of Hymenoptera in British Museum); Kohl, 1890b:441 (original description copied); Dalla Torre, 1897:420 (in catalog of world Hymenoptera); Ashmead, 1899d:353 (in checklist of North American Sphecidae); Casolari and Casolari Moreno, 1980:103 (specimen in M. Spinola collection, Torino); Pagliano, 2008:530 (specimen in M. Spinola collection, Torino).
- Sphex rusticus* Dahlbom, 1843:28 (as *rustica*, incorrect original termination). Not available: published as synonym of *Priononyx thomae* (Article 11.5).
- Enodia pubidorsum* A. Costa, 1862a:17 (as *pubidorsa*) and 1862b:69, ♂. Holotype: ♂, Brazil: no specific locality (Napoli). Synonymized with *Chlorion thomae* by Fernald, 1906:342 and R. Bohart and Menke, 1963:159 (as new synonym). – **As *Sphex pubidorsus***: Murray in Muesebeck, Krombein, and Townes, 1951:973 (new generic assignment, in catalog of North American Hymenoptera); K. Cooper, 1953:33 (Massachusetts: Penikese Island). – **As *Chlorion pubidorsum***: Fernald, 1931a:441 (new generic assignment, study of type), 1943a:287 (Florida); Brimley, 1938:444 (North Carolina: statewide); Dreisbach, 1944:268 (in key to Sphecinae of Michigan, as *rubidorsum*); Willink, 1948a:319 (in key to *Chlorion thomae* species group); Krombein, 1950a:268 (North Carolina: Dare County), 1953a:295 (visiting foliage of *Pinus serotina* Michx.), 296 (visiting foliage of *Quercus marilandica* Muenchh.), 298 (visiting flowers of *Cephalanthus occidentalis* Linnaeus, Rubiaceae), 332 (North Carolina), 1953b:123 (visiting foliage of *Quercus virginiana* Mill.), 124 (visiting flowers of *Pluchea* sp., Asteraceae), 133 (North Carolina: Kill Devil Hills); Krombein and Evans, 1954:233 (Florida), 1955:232 (Florida); Krombein, 1958f:191 (in supplement to catalog of North American Hymenoptera: *Priononyx canadensis* Provancher was published in 1887, not 1889); L. Davis, 1978:216 (North Carolina: Kill Devil Hills, data from Krombein, 1953a). – **As *Priononyx pubidorsum***: Strickland, 1947:128 (new generic assignment, Canada: Alberta: Cypress Hills, Lethbridge, Medicine Hat); R. Bohart, 1958b:92 (in key to North American *Prionyx*); nec Evans, 1958a:183 and Linsley, 1962:156 (= *Prionyx parkeri*); F. Parker, 1960:206, 207 (in key to North American *Prionyx*); Evans and Linsley, 1960:32 (regular member of sleeping aggregation at Southwest Research Station, Arizona).

Priononyx thomae var. *antillarum* de Saussure, 1867:43, ♀, ♂ (as *Antillarum*, incorrect original capitalization). Syntypes: Antillean Islands: no specific locality (depository?). Synonymized with ..

Priononyx thomae var. *mexicanus* de Saussure, 1867:43, ♀. Holotype: or syntypes: ♀, Mexico: Michoacán: no specific locality (NHMW). Synonymized with ..

Sphex edwardsi Cameron, 1903e:230, ♀, ♂ (as *Edwardsi*, incorrect original capitalization). Lectotype: ♂, Ecuador: Ambato (BMNH), designated by Menke in R. Bohart and Menke, 1976:134. Synonymized with ..

Sphex platensis Brèthes, 1908:146, ♀, ♂. Lectotype: ♂, Brazil: Santa Catarina: Nova Friburgo (MACN), designated by Menke in R. Bohart and Menke, 1976:134. Synonymized with *Chlorion thomae* by Willink, 1948a:315. – Jörgensen, 1912:286 (Argentina: Mendoza Province); Schrottky, 1920:187 (may be a synonym of *Priononyx thomae*); Liebermann, 1931:80 (in revision of Argentinean Sphecini); Genise, 1990:27 (type material in MACN). – **As *Priononyx platensis***: Schrottky, 1913a:225 (new generic assignment, Argentina).

? *Sphex altibia* Strand, 1911a:152, ♂. Holotype: ♂, Ecuador: Riobamba (MNH). Tentatively synonymized with *Prionyx thomae* by R. Bohart and Menke, 1976:134.

54. *trichagyrs* (Spinola)

Sphex trichagyrs Spinola, 1839:466, ♂ (as *trichagyra*, incorrect original termination). Lectotype: ♂, Egypt: no specific locality (M. Spinola collection, Torino), designated by de Beaumont, 1952e:45. – F. Smith, 1856:244 (in catalog of Hymenoptera in British Museum); Kohl, 1885b:185 (as junior synonym of *Sphex albisectus*); Honoré, 1944a:65 (in revision of Egyptian Sphecini, as *trichagyrius*); de Beaumont, 1951e:268 (Morocco: as *trichagyrius*); Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino); Pagliano, 2008:530 (lectotype in M. Spinola collection, Torino). – **As *Prionyx trichagyrs***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Guichard, 1988a:121 (Arabian Peninsula); Gadallah and Assery, 2004a:217 (in key to Sphecidae of Jeddah Region, Saudi Arabia), 222 (in catalog of Sphecidae of Saudi Arabia); Roche, 2007a:43 (in checklist of Egyptian Sphecidae), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae); Dollfuss, 2008b:1416 (Oman: Jebel Muwarrhan; Tunisia: oasis Douz; Yemen: Lawdat northeast of Aden); Murai and Amr, 2011:110, 120 (first record from Syria: Al Thawrah Nature Reserve at 35°51'N 28°38'E); Schmid-Egger, 2011b:603 (United Arab Emirates: Sharjah Desert Park); Gadallah, Al Dhafer, Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia); Ben Khedher, Yildirim, Braham, and Ljubomirov, 2020a:316 (in list of Tunisian Sphecidae *sensu stricto*); Gadallah, 2020d:86 (in list of aculeate wasps of Arabian Peninsula); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*).

Sphex leucosoma Kohl, 1890b:338, ♀. Holotype or syntypes: ♀, Egypt: Cairo (NHMW). Synonymized with *Sphex trichagyrs* by Honoré, 1944:65, synonymy confirmed by de Beaumont, 1952e:45. – Dalla Torre, 1897:428 (in catalog of world Hymenoptera); Berland, 1926b:169 (Senegal: Dakar); Rungs, 1936:24 (Morocco: Todra oasis); Berland, 1956:1169 (in revision of African Sphecini); Dollfuss, 1989:12 (type material in NHMW) – Nec Schulz, 1905b:34 (as *Parasphex marginatus* var. *leucosoma* = *Prionyx senilis*). – **As *Chlorion leucosoma***: Arnold, 1928c:353 (new generic assignment, in revision of Sphecini of southern Africa), 1930:17 (in checklist of Afrotropical Sphecidae).

55. *viduatus* (Christ)

Sphex viduatus Christ, 1791:305, sex not stated (as *viduata*, incorrect original termination). Holotype or syntypes: France: Provence Region: no specific locality (lost). – Kohl and Handlirsch, 1889:275 (Turkmenistan: Pul-i-Hatun); Kohl, 1890b:332 (in revision of world Sphecini); F. Morawitz, 1891a:202 (Russia: Astrakhan Government); Bingham, 1896a:440 (Sri Lanka: Colombo), 1897:252 (redescription); Dalla Torre, 1897:446 (in catalog of world Hymenoptera); Bingham, 1898a:105 (Yemen: Aden); Magretti, 1899:602 (Somalia: Lugh at 3°48'N 42°33'); E. Saunders, 1904c:604 (Spain: Mayorca); Dusmet and Mercet, 1906:508, 514 (in key to Spanish Sphecini); von Schulthess, 1909:441 (Libya: Dernah); Morice, 1911:74 (Algeria: Biskra); Ferton, 1912a:408 (Algeria, nest and prey); Strand, 1913a:82 (Taiwan); von Schulthess, 1914:286 (Cameroon); Strand, 1915:90 (Sri Lanka); Roth, 1925:381 (in revision of North African Sphecini); Berland, 1926b:169 (miscellaneous locality records); von Schulthess, 1926b:209 (Tunisia); Kruger, 1929a:21 and 1929b:56 (Libya: Cyrenaica: Giarabub); G. Carpenter, 1930b:290 (nest closure); Schouteden, 1930:95 (Zaire); Dusmet y

Alonso, 1931:7 (Portugal: dos Medos); Guiglia, 1932a:127 (Somalia: Mogadishu), 1932d:472 (Libya: Cufra oasis); Nadig, 1933:103 (Morocco); Guiglia, 1934b:295 (Libya: bibliography and summary of locality records, as *viduatas*); Yasumatsu, 1935b:36 (Japan: Ryukyu Archipelago: Island of Yonakuni), 1938:87 (revision; Manchuria, Ryukyus, Taiwan); Guiglia, 1939c:186 (Libya: Fezzan: Gat, Tunin), 1942b:230 (Libya; as *viduatum*); Giner Marí, 1943a:81 (in Sphecidae Fauna of Spain); Guiglia, 1943c:76 (Ethiopia: Gamo Gofa: Sagan – Omo region); Honoré, 1944a:62 (in revision of Egyptian Sphecini); Giner Marí, 1945b:362 (Morocco: Western Sahara), 1945e:220 (Morocco: Western Sahara); de Beaumont, 1947b:382 (Cyprus); Giner Marí, 1947:19 (Morocco: Western Sahara); de Andrade, 1949:9 (Portugal: Pinhal dos Medos); Berland, 1950b:295 (Niger: Aïr area); de Beaumont, 1950d:6 (Egypt: Siwa oasis), 1950f:396 (Morocco); Guiglia, 1950:248 (Ethiopia: Gamo Gofa: Asile, as *viduatum*); Pittioni, 1950:20 (Cyprus); de Beaumont, 1951e:267 (Morocco), 1953a:173 (Mauritania); de Beaumont and Bytinski-Salz, 1955:41 (Israel); Leclercq, 1955h:36 (bibliographic references, faunal records from Africa); Ceballos, 1956:365 (in catalog of Hymenoptera of Spain); de Beaumont, 1956a:181 (Libya); Berland, 1956:1167 (in revision of African Sphecini); Compte Sart, 1959:131 (Spain: Mayorca); Diniz, 1959:27 (Portugal: Mata do Urso, Pinhal dos Medos); Grandi, 1959b:287 (Algeria: Bou Megueur, Goriana in Hodna District); Suárez, 1959:53 (Spain: Almería Province); de Beaumont, 1960a:5 (Greece: Island of Rhodes), 1960b:227 (Libya); de Beaumont, 1961b:272 (Afghanistan), 1961c:45 (Greece: Island of Crete); Tsuneki, 1962a:6 (Japan: Ryukyus Islands: Island of Amami-Oshima); Ceballos, 1964:88 (in supplement to catalog of Spanish Sphecidae); de Beaumont, 1965a:13 (Greece); Iwata, 1965:106 (number of oocytes); de Beaumont, 1967a:272 (Turkey); Tsuneki, 1967i:383 (Ryukyu Islands), 1967j:4 (Taiwan); Kazenas, 1969a:21 (Kazakhstan: Ili River between Ili and Ayak-Kalkan); Robertson, 1969:480 (Tanzania: Ukiriguru); Tsuneki and Iida, 1969:4 (nesting habits), 16 (description of larva); Tsuneki, 1971f:2 (Taiwan); Haneda, 1972a:5 (Taiwan); Kazenas, 1972b:112 (Kazakhstan); Tano, 1972:22 (Ryukyu Islands); Murota, 1973a:101 (Ryukyu Islands: Amami group); Murota, 1973b:116 (Taiwan); Kazenas, 1974b:110 (feeding on flowers of *Melilotus albus* Desr., Fabaceae, in Kazakhstan); Chhotani and Ray, 1975:27 (India: Rajasthan: Gudha); Georgiou, 1977:192 (Cyprus); Kazenas, 1978b:44 (in key to Sphecidae of Kazakhstan and Central Asia); Tüzün and Yüksel, 2010:4467 (Turkey: Niğde Province). – **As *Enodia viduata***: Roth, 1924:123 (new generic assignment, Algeria: Nemours, now Ghazaouet). – **As *Chlorion viduatum***: Arnold, 1928c:349 (new generic assignment, in revision of Sphecini of southern Africa, distributed throughout Africa), 1930:17 (in checklist of Afrotropical Sphecidae); Guiglia, 1939c:186 (Libya), 1940e:292 (Somalia: Mogadishu); Arnold, 1943:79 (Zaire). – **As *Prionyx viduatus***: Iwata, 1964b:355 (new generic assignment, nesting behavior in Thailand); Myartseva, 1972a:85 (Turkmenistan; as *Prionix*); R. Bohart and Menke, 1976:134 (in checklist of world Sphecidae); Valetta, 1979:215 (Malta); Guichard, 1980:224 (Oman); Roche, 1981:2 (in checklist of Sphecidae of United Arab Emirates); Mingo and Gayubo, 1983:155 (Spain); Tsuneki, 1982g:55 (known from the Ryukyu archipelago); Brockmann, 1985b:312 (nest closure summary); Gayubo and Heras, 1986:29 (Spain: Provincia de Segovia and de Valladolid; floral records); Islamov, 1986:516 (Uzbekistan: Tashkent Oblast'); Clark, 1987:477 (Vietnam, prey: *Aiolophus thalassinus tamulus* (Fabricius), Acrididae); Gayubo, 1987:107 (Spain: Ciudad Real Province); Guichard, 1988a:121 (Arabian Peninsula); Gayubo, Asís, and Tormos, 1990a:10 (Spain); Pagliano, 1990:58 (in catalog of Italian Sphecidae); Gayubo, Borsato, and Osella, 1991:394 (first record from Italy: Sicilia: Lampedusa); Gayubo and Torres, 1991:Table I (Spain: Salamanca; effects of urban pressure); Kazenas and Nasyrova, 1991:38 (Kazakhstan: preying on *Calliptamus barbarus cephalotes* F.W., Acrididae); Schembri, 1991:176 (Malta); Sk. Yamane and Ikudome, 1990:100 (distribution in Ryukyu Islands, Japan); Gayubo, Borsato, and Osella, 1992:277 (Greece); Ebrahimi, 1993:97 (Iran); Hohmann, La Roche, Ortega, and Barquín, 1993:206 (first record from Canary Islands: La Gomera); Jha and Farooqi, 1994:13 (description and illustration of male genitalia); Negrísolo in Minelli, Ruffo, and La Posta, 1995b:2 (in catalog of Italian fauna); Pagliano and Scaramozzino, 1995:730 (Italy: Island of Lampedusa); van Vondel, 1995:29 (specimens from India in Natuurmuseum Rotterdam); Minoranskiy and Shkuratov, 1996:81 (Russia: Rostov Oblast'); Wu and Zhou, 1996a:46 (in revision in Economic Insect Fauna of China); Bitsch, Barbier, Gayubo, Schmidt, and Ohl, 1997:64 (in Sphecidae Fauna of Western Europe); Lauterbach, 1997a:255 (Canary Islands: Island of La Gomera); Kazenas, 1998b:127 (in Sphecidae Fauna of Kazakhstan); Nazarova, 1998:40 (Tajikistan: Tigrovaya Balka Nature Reserve); Porter, Stange, and Wang, 1999:5 (in checklist of Sphecidae of Taiwan); Yamane, Ikudome, and Terayama, 1999:477 (Japan: in Identification Guide to Sphecidae of Nansei =

Ryukyu Islands); Shkuratov, 2000:55 (Russia: Rostov Oblast': Vëshenskaya village area at 49°37'N 41°45'E); Kazenas, 2001b:16 (in checklist of Sphecidae of Kazakhstan and Central Asia), 86 (review of known biology); Seyoum and Pulawski, 2001:322 (potential control agent of acridid pests in Ethiopia); Shkuratov, 2001:17 (prey: *Stenobothrus lineatus* (Panzer), Acrididae); Kazenas, 2002a:31 (geographic distribution, collecting localities in Kazakhstan); Ohl and Linde, 2003:149 (number of ovarioles); Shkuratov, 2002a:383 (Russia: Rostov Oblast'); S. Gess and F. Gess, 2003:92 (Namibia and South Africa: visiting flowers of *Psilocaulon glareosum* (Berger) Dinter and Schwant., Aizoaceae, *Hermbstaedtia odorata* (Burch.) T. Cooke., Amaranthaceae, *Asclepias buchenaviana* Schinz, Apocynaceae, *Deverra denudata* (Viv.) Pfisterer and Podl., Apiaceae, and *Zygophyllum simplex* L., Zygophyllaceae); Pagliano, 2003b:130 (Italy: islands of Lampedusa and Pantelleria); Gadallah and Assery, 2004a:217 (in key to Sphecidae of Jeddah Region, Saudi Arabia), 222 (in catalog of Sphecidae of Saudi Arabia), 2004b:1396 (skeletal parts of sting apparatus); Kazenas, 2004b:99 (Kazakhstan: western Tien Shan Mountains, as *viduatus viduatus*), 2004d:27 (Kazakhstan: northern Caspian region); Gülmez and Tüzün, 2005:48 (Turkey: Ankara Province); Pagliano and Negrisolo, 2005:55 (in Sphecid Fauna of Italy); Shoreenko, 2005a:162 (Ukraine: Crimea); Hua, 2006:276 (in list of Chinese insects, geographic distribution); Terayama and Tano, 2006:7, 13, 17 (in key to Japanese Ampulicidae and Sphecidae); Kazenas, 2007a:89 (Kazakhstan: Akmala Oblast': Kurgandzhin Nature Reserve and vicinity); Roche, 2007a:43 (in checklist of Egyptian Sphecidae, redescription, as *viduatus viduatus*), 2007b:3 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae, as *viduatus viduatus*); Dollfuss, 2008b:1416 (locality records from 21 countries); Ljubomirov and Yildirim, 2008:32 (in catalog of Sphecidae of Turkey); Baños-Picón, Asís, Gayubo, and Tormos, 2009:310 (Spain: frequency of specimens collected with hand nets and Malaise traps); Danilov, 2009:55 (Russia: Western Siberia: Kulundinskaya Steppe); Gayubo, González, Tormos, and Asís, 2009:362 (Spain: Valladolid: Reserva Natural Riberas de Castronuño – Vega del Duero); González, Gayubo, Asís, and Tormos, 2009:622 (Spain: Salamanca and Zamora provinces: Arribes del Duero Natural Park); Shoreenko, 2009:366 (in list of Sphecidae *sensu lato* of Crimea); Bitsch, 2010:105 (in supplement to vol. II of Faune de France, 1997: records from Lampedusa and Pantelleria islands reported); Danilov, 2010b:44 (distribution of Palearctic-Ethiopian-Indomalayan type); Rudoiskatel', 2010:147 (Russia: southern Ural Mountains, as *viduatus viduatus*); Sakenin, Samin, and Bagriacik, 2010:17 (Iran: Sistan and Baluchistan: Nikshahr); Cruz-Sánchez, Asís, Gayubo, Tormos, and González, 2011:497 (Spain: Salamanca and Zamora provinces: Arribes del Duero Natural Park: effects of wildfire), 501 (exclusive species of burned areas); Murai and Amr, 2011:120 (recorded from Syria by Dollfuss, 2008b); Schmid-Egger, 2011b:603 (United Arab Emirates: near Qurayyah); Danilov, 2012a:164 and 2012b:62 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:67 (bibliographic references, geographic distribution); Prisniy, 2012:46 (Russia: Belgorod Oblast'); Strumia, Pagliano, and Gayubo, 2012:55 (Italy: Toscana: Riserva di San Rossore); Gadallah, Al Dhafer, Aldryhim, Fadl, and Elgharbawy, 2013:362 (in new catalog of Sphecidae of Saudi Arabia, as *viduatus viduatus*); Kazenas, 2013a:24, 25 (color photographs of adult wasps, short information on geographic distribution and nesting habits); Baldock, 2014:354 (Spain: Island of Mallorca); Danilov, 2014a:424 (Russia: Siberia: Omsk Oblast'), 2014b:516 (in key to Sphecidae s.s. of Siberia, as *viduatus viduatus*); Dunford, Turbyville, and Leavengood, 2014:11 (listed as medically important in Afghanistan); Ebrahimi, 2014:24 (Iran: Māzandarān, Sīstān-Baluchestān, and Tehrān provinces); Kazenas, 2014a:132 (Kazakhstan: Karatau Mountain Range, as *viduatus viduatus*); Schmid-Egger, 2014:623 (United Arab Emirates); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces, as *viduatus viduatus*); Augul, Abdul-Rassoul, and Kaddou, 2015:116 (in key to Sphecini of Iraq), 117 (Iraq: locality records), 119 (illustrations); Koçak and Kemal, 2015:279 (in checklist of Hymenoptera of Turkey); Pham, Kumar and Danilov, 2015:1587 (in list of Sphecidae *sensu lato* of Vietnam, as *viduatus viduatus*); Gülmez and Dizer, 2016:59 (Turkey: Tokat Province); Mokrousov and Popov, 2016:564 (Russia: Abkhazia, Krasnodarskiy Krai); Yildirim, Ljubomirov, Özbek, and Yüksel, 2016:6 (Turkey: Antalya: Kepez: Odabaşı); Arens, 2017a:631 (Greece: Peloponnesus); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:27 (Iran: known from Mazandaran, North Khorasan, Sistan-o Baluchestan, and Tehran provinces); Danilov, 2017b:215 (in catalog of Sphecidae s.s. of Russia, as *viduatus viduatus*); Danilov and Mokrousov, 2017a:108 (Russia: Kalmykia, as *viduatus viduatus*); Can and Gülmez, 2019:349 (in key to *Prionyx* of Turkey, as *viduatus viduatus*); Gülmez, 2019:3 (Turkey: Ankara, Sivas, and Tokat provinces: no specific localities); Pham, Truong, Th.T. Nguyen, Th.H. Nguyen, Q. Nguyen, and Th.M. Nguyen,

- 2019:73 (Vietnam: Hanoi and vicinity); Ben Khedher, Yildirim, Braham, and Ljubomirov, 2020a:316 (in list of Tunisian Sphecidae *sensu stricto*; additional records: Tunisia: Sfax and Tunis provinces); Bitsch, Barbier, and Jacobs *in* Bitsch, Barbier, Gayubo, Jacobs, Leclercq, and Schmidt, 2020:141 (in Sphecid Fauna of Europe, not occurring in France); Cassar and Mifsud, 2020:164 (in checklist of Sphecidae s.s. of Malta); Danilov, 2020:320 (specimens from Japan, Kazakhstan, Kyrgyzstan, Tajikistan, and Russia: Astrakhan' Oblast' and Kalmykia, as *viduatus viduatus* in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Gadallah, 2020d:86 (in list of aculeate wasps of Arabian Peninsula); Cross, Baldock, and Wood, 2021:19 (in catalog of Sphecidae *sensu lato* of Portugal); Embergenov, Élmurodova, Amirov and Kimyonazarov, 2022:47 (Uzbekistan: Khiva area); Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*); Embergenov, Seytmuratov, and Medetov, 2023:237 (Uzbekistan: first record from southern Aral region); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey, as *viduatus viduatus*).
- Sphex pubescens* Fabricius, 1793:205, sex not stated. Lectotype: ♂, Guinea: no specific locality (ZMUC), designated by van der Vecht, 1961a:33. Tentatively synonymized with *Prionyx viduatus* by Kohl, 1890b:332, synonymy confirmed by van der Vecht, 1961a:33. – Fabricius, 1796:156 (in Index to his Entomologia Systematica, 1793); Turton, 1801:487 (re-description); Dufour, 1854a:375 (Algeria: Pontéba); Lepeletier de Saint Fargeau, 1845:359 (in revision of world Hymenoptera); F. Smith, 1856:246 (in catalog of Hymenoptera in British Museum), 267 (as synonym of *Parasphex fervens*); Fairmaire, 1858:264 (Gabon); Kohl, 1885b:188 (in revision of Palearctic *Sphex*); Ed. André, 1888:131 (in revision of Sphecidae of Europe and Algeria), 10* (bibliographic references); Cameron, 1889c:106 (in list of Sphecidae of Oriental Region); Medina, 1894a:260 (Spain: Pozuelo de Calatrava), 1896:104 (Spain: Cádiz); Ceballos, 1949:101 (Spain), 1956:364 (in catalog of Hymenoptera of Spain); Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino); Pagliano, 2008:530 (specimens in M. Spinola collection, Torino, are *Prionyx pruinosus* and *Prionyx viduatus*). – **As *Enodia pubescens***: Radoszkowski, 1892:586 (description of male genitalia).
- Enodia canescens* Dahlbom, 1843:28, ♀, ♂. Syntypes: Senegal and Guinea: no specific localities (Lund). Synonymized with *Enodia fervens* by Dahlbom, 1845:439.
- Sphex micans* Eversmann, 1849:368, ♀, ♂. Syntypes: Russia: lower Volga area (ZIN). Synonymized with *Enodia lividocincta* by Radoszkowski, 1887b:91, with *Sphex lividocincta* by Kohl, 1885b:190 (tentatively), 1889a:24, and with *Sphex viduatus* by Kohl, 1890b:332. – Radoszkowski, 1871:199 (Iran: Astrabad, now Gorgan); Kohl, 1885b:205 (original description copied); nec Ed. André, 1888:133 (= *Prionyx lividocinctus*); André, 1888 9* (bibliographic references); Radoszkowski, 1887b:91 (in list of Transcaspien Hymenoptera); Pagliano, 2008:529 (specimens in M. Spinola collection, Torino, are *Sphex dorsalis*).
- Sphex leuconotus* F. Morawitz, 1890:579, ♀, junior primary homonym of *Sphex leuconotus* Brullé, 1833. Lectotype: ♀, Turkmenistan: Ashkhabad (ZIN), designated by Danilov, 2012b:68. Synonymized with *Prionyx viduatus* by Danilov, 2012b:68. – Kohl, 1890b:338 (original description copied); Dalla Torre, 1897:428 (in catalog of world Hymenoptera); Gussakovskij, 1933b:273 (Iran, as tentative synonym of *Sphex viduatus* Christ); Myartseva, 1963b:59 (Turkmenistan: lower Murgab River); Danilov, 2016:341 (lectotype preserved in Zoological Institute, Sankt Petersburg, Russia). – **As *Prionyx leuconotus***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae).
- Sphex granti* W.F. Kirby, 1900: 23, ♀, ♂. Syntypes: Yemen: Abd-el-Kuri islands: no specific locality (BMNH). Synonymized with *Sphex pollens* by Kohl, 1906a:198, and with ..
- Sphex platynotus* Matsumura, 1912:177, 178, ♀. Holotype or syntypes: ♀, Japan: Okinawa: no specific locality (depository?). Synonymized with *Sphex viduatus* by Yasumatsu, 1935b:36. – Matsumura and Uchida, 1926:39 (Okinawa).
- Sphex perezii* Berland, 1926b:170, ♀, ♂. Lectotype: ♀, Senegal: no specific locality (MNHN), designated by Menke *in* R. Bohart and Menke, 1976:134. Synonymized with *Prionyx viduatus* by Guichard, 1988a:121. – Berland, 1956:1168 (in revision of African Sphecini); de Beaumont, 1958b:56 (Algeria: Tassili des Ajjer). – **As *Chlorion perezii***: Arnold, 1928c:352 (new generic assignment, original description translated into English), 1930:17 (in checklist of Afrotropical Sphecidae). – **As *Prionyx perezii***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae).

? *Prionyx zanoni* Gribodo in Zanon, 1926:88, ♀, ♂. Syntypes: Libya: Fueihat 15 kms out of Benghazi (lost?). Synonymized with .. – Guiglia, 1934b:305 (Libya: bibliography and summary of locality records); Penati and Mariotti, 2015:131 (in list of Hymenoptera described by G. Gribodo).

ssp. *mocsaryi* (Kohl)

Enodia argentata Mocsáry, 1883:36, ♀, junior secondary homonym of *Sphex argentatus* Fabricius, 1787. Syntypes: ♀, southern Russia or Caucasus: no specific locality (TMB). – **As *Prionyx viduatus argentatus***: de Beaumont in R. Bohart and Menke, 1976:134 (new generic assignment, new status, in checklist of world Sphecidae); Tormos and Jiménez, 1987a:122 (Spain: Valencia), 1987b:316 (Spain: Valencia Province: Dehesa de El Saler); Chinin, 1991:111 (Russia: Samara Oblast'); Kazenas, 2001b:16 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:31 (geographic distribution, collecting localities in Kazakhstan); Shlyakhtenok and Skibinska, 2002:32 (Belarus': no specific locality); Shkuratov, 2004a:73 (Russia: Rostov Oblast'); Yildirim and Ljubomirov, 2005:1787 (Turkey: Erzincan: Kemah), 2007:116 (Turkey: Erzincan: Oltu; Isparta: Gökçay); Ljubomirov and Yildirim, 2008:31 (in catalog of Sphecidae of Turkey); Danilov, 2012a:164 and 2012b:62 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:68 (bibliographic references, geographic distribution); Japoshvili and Ljubomirov, 2012:96 (Turkey: Isparta: Gölcük Nature Park 8 km southwest of city of Isparta); Yildirim, 1912:74 (Turkey: Erzurum: Olur); Pankov, 2013:228 (Russia: Ivanovo Oblast'); Shlyakhtenok, 2013:131 (in annotated catalog of aculeate wasps of Belarus'); Danilov, 2014b:516 (in key to Sphecidae s.s. of Siberia); Kazenas, 2014a:132 (Kazakhstan: Karatau Mountain Range); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces); Shoreenko, 2015:317 (in list of Sphecidae *sensu lato* of Crimea), 2017:76 (in Crimea collected in July and August), 2018:127 (Crimea, including localities, habitats, and number of specimens); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

Sphex mocsaryi Kohl, 1885b:187, substitute name for *Enodia argentata* (a junior secondary homonym in *Sphex* of *Sphex argentatus* Fabricius, 1781). – Kohl, 1885b:187 (in revision of Palearctic *Sphex*); Ed. André, 1888:131 (in revision of Sphecidae of Europe and Algeria), 9* (bibliographic references); nec Kohl, 1890b:342 (= *Prionyx nudatus*); F. Morawitz, 1894:339 (Turkmenistan: Krasnovodsk); Dalla Torre, 1897:432 (in catalog of world Hymenoptera); Dusmet and Mercet, 1906:508, 514 (in key to Spanish Sphecini); von Schulthess, 1909:441 (Libya: Dernah, Gherrqn); Coulon, 1925:116 (Spain: Montario); Guiglia, 1934b:295 (Libya: bibliography and summary of locality records); Gussakovskij, 1934a:2 (China: Inner Mongolia), 1935:413 (Tajikistan); Kuntze and Noskiewicz, 1938:378 (in Poland known only from Podole region, now Ukraine); Ceballos, 1949:101 (Spain); Leclercq, 1955h:38 (bibliographic references); Ceballos, 1956:364 (in catalog of Hymenoptera of Spain); de Beaumont, 1957b:130 (as synonym of *Sphex viduatus*); Noskiewicz and Puławski, 1960:41 (in key to Polish Sphecidae, not yet found in Poland); Myartseva, 1963b:59 (Turkmenistan: lower Murgab River); Romanova, 1969:133 (Russia: North Caucasus); Kazenas, 1968a:806 (nesting habits in Kazakhstan: Mangyshlak Peninsula); Balthasar, 1972:424 (in Sphecid Fauna of Czechoslovakia: may be expected in the country. – **As *Sphex viduatus mocsaryi***: Kazenas, 1969a:21 (new status, Kazakhstan: Mangyshlak Peninsula, Golodnaya Step', Ili River), 1972b:112 (Kazakhstan), 1974b:110 (feeding on flowers of *Nitraria schoberi* L., Zygophyllaceae, and *Daucus carota* L., Apiaceae, in Kazakhstan), 1978b:44 (in key to Sphecidae of Kazakhstan and Central Asia); Pulawski, 1978:184 (in key to Sphecidae of European part of USSR). – **As *Prionyx viduatus mocsaryi***: Kazenas and Nasyrova, 1991:38 (Kazakhstan: new generic assignment, preying on *Dociostaurus ingens* (Ingen.), Acrididae); Blagoveshchenskaya, 1994:89 (Russia: Ul'yanovsk Oblast'); Nazarova, 1998:40 (Tajikistan: Tigrovaya Balka Nature Reserve); Shkuratov, 2002b:139 (Russia: Rostov Oblast': Rostovskiy Nature Reserve at 46°27'N 42°41'E); Kazenas, 2004b:99 (Kazakhstan: western Tien Shan Mountains); Protsenko, Fateryga, Ivanov, and Puzanov, 2012:58 (Ukraine: Crimea); Danilov, 2017b:215 (in catalog of Sphecidae s.s. of Russia) Danilov, 2020:320 (specimens from Kazakhstan and Russia: Volgograd Oblast' and Siberia in Institute of Systematics and Ecology of Animals, Novosibirsk, Russia); Maharramov, Mokrousov, and Proshchalykin, 2020:46 (Azarbaijan: Nakichivan Autonomous Republic); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey). *Sphex gobiensis* Tsuneki, 1971k:142, ♀. Holotype: ♀, Mongolia: Middle Gobi Aymag: Delgerhangay (TMB). Synonymized with *Prionyx viduatus argentatus* by Danilov, 2012b:68. – **As *Prionyx gobiensis***: R. Bohart and Menke, 1976:133 (new generic assignment, in checklist of world Sphecidae).

ssp. *pollens* (Kohl)

Sphex pollens Kohl, 1885b:186, ♀. Syntypes: Greece: Athens (NHMW). – Ed. André, 1888:127 (in revision of Sphecidae of Europe and Algeria), 10* (bibliographic references); Kohl, 1890b:343 (in revision of world Sphecini); Dalla Torre, 1897:437 (in catalog of world Hymenoptera); Kohl, 1906a:198 (description of ♂); Mantero, 1915:325 (Libya); Fahringer, 1922:178 (Turkey); Bischoff, 1930a:216 (Tajikistan: Pamir); Guiglia, 1934b:295 (Libya: bibliography and summary of locality records); Leclercq, 1956g:324 (Greece); Myartseva, 1964a:75 (nesting habits in Turkmenistan), 1965:82 (Turkmenistan: Akibay, Bayram-Ali; Murgab district; Mary district). – **As *Prionyx pollens***: Myartseva, 1966:48 (new generic assignment, preying on orthopterans), 1972a:84 (Turkmenistan), 1972b:106 (parasite: *Senotainia albifrons* Rondani, Sarcophagidae). – **As *Sphex viduatus pollens***: de Beaumont, 1965a:13 (new status, Greece). – **As *Prionyx viduatus pollens***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Dollfuss, 1989:12 (type material in NHMW); Ljubomirov and Yildirim, 2008:31 (in catalog of Sphecidae of Turkey); Yildirim, 2014:30 (Turkey: distribution by biogeographic provinces); Can and Gülmez, 2019:349 (in key to *Prionyx* of Turkey); Kaplan and Yildirim, 2023:1693 (in checklist of Sphecidae *sensu lato* of Turkey).

56. *xanthabdominalis* Li and Yang

Prionyx xanthabdominalis Li and Yang, 1995a:140, ♀, ♂. Holotype: ♀, China: Ningxia Province: Helanshan Mountain (Beijing Agricultural Univ.). – Danilov and Odintsev, 2023:435 (in subgenus *Prionyx*).

57. *zarudnyi* (Gussakovskij)

Sphex zarudnyi Gussakovskij, 1933b:372, ♀, ♂. Lectotype: ♀, Iran: Bazman District: Bazman-Tagab at 27°51'N 60°10'E (ZIN), designated by Danilov, 2012b:68. – de Beaumont, 1968b:149 (member of *Sphex subfuscatus* species group). – **As *Prionyx zarudnyi***: R. Bohart and Menke, 1976:134 (new generic assignment, in checklist of world Sphecidae); Danilov, 2012a:163, 164 and 2012b:61 (in keys to *Prionyx* of Russia and adjacent countries, respectively in Russian and English), 2012b:68 (bibliographic references, geographic distribution); Danilov, 2016:353 (lectotype preserved in Zoological Institute, Sankt Petersburg, Russia); Jahantigh, Rakhshani Mokhtari, and Ramroodi, 2017:27 (Iran: known from Sistan-o Baluchestan Province); Danilov, 2019:78 (in review of Palearctic Prionychini, member of subgenus *Harpactopus*); Danilov and Odintsev, 2023:433 (in subgenus *Harpactopus*).

sp.

Lavigne and Pfadt, 1966:31 (Wyoming; preying on grasshoppers *Agenotettix deorum* (Scudder), *Melanoplus gladstoni* Scudder, and *Trachyrhachys kiowa* (Thomas), all Acrididae); Kingsley, Bailowitz, and Smith, 1987:19 (Arizona: Organ Pipe Cactus National Monument: Quitobaquito Springs area); Naumann, 1993:182 (Australia: Queensland: Heathlands area in Cape York); Starr and Hook, 2003:22 (in catalog of Aculeata of Trinidad, West Indies, possibly *Prionyx fervens*); Ruíz Cancino, Coronado Blanco, and Horta Vega, 2005:170 (Mexico: recorded from Tamaulipas State); Roche, 2007a:46 (in checklist of Egyptian Sphecidae, description); Ghazi-Soltani, Ebrahimi, Iranipour, and Pour Abad, 2010:797 (Iran: East Azerbaijan: county of Tabriz, probably *Prionyx kirbii*); Mamadou, Mazih, Ghaout, and Hormatalah, 2015:59 (Niger: Agadir Region: Tafidet Valley at 18°09'16"N 9°30'52"E; use of anti-locust chemicals chlorpyrifos ethyl and fenitrothion greatly reduced this species population); Jahantigh, Rakhshani, Mokhtari, and Ramroodi, 2017:28 (Iran: East Azerbaijan Province).

NOMINA NUDA IN PRIONYX

Enodia rufipes: Casolari and Casolari Moreno, 1980:103 (specimens in M. Spinola collection, Torino).

Pseudosphex noverca Kaye, 1910:... – R. Bohart and Menke, 1976:134 (in checklist of world Sphecidae).