

Last updated: 16 May 2025

PSEUDOMICROIDES

Pseudomicroides Antropov, 2001a:810. Type species: *Belomicroides olgae* Kazenas, 1979, by original designation.

Revision: Antropov, 2001a:809.

1. *applanatus* Antropov

Pseudomicroides applanatus Antropov, 2011a:620, ♀, ♂. Holotype: ♀, United Arab Emirates: north of Ajman at 25°43'N 55°48'E (ZMHU). – Gadallah, 2020b:48 (in list of aculeate wasps of Arabian Peninsula).

2. *arabicus* Antropov

Pseudomicroides arabicus Antropov, 2011a:616, ♀, ♂. Holotype: ♀, United Arab Emirates: Al'Ayn (BMNH). – Gadallah, 2020b:48 (in list of aculeate wasps of Arabian Peninsula).

3. *desertus* (Antropov)

Belomicroides desertus Antropov, 1994a:93, ♀. Holotype: ♀, Turkmenistan: Repetek (ZMMU). – Kazenas, 2001b:35 (in checklist of Sphecidae of Kazakhstan and Central Asia). – **As *Pseudomicroides desertus***: Antropov, 2001a:811 (new combination, in revision of *Pseudomicroides*).

4. *elvirae* (Gayubo)

Belomicroides elvirae Gayubo, 1983b:39, ♀. Holotype: ♀, Canary Islands: Fuerteventura: Gran Tarajal (MNCN). – Hohmann, La Roche, Ortega, and Barquín, 1993:243 (detailed records from Canary Islands); Leclercq, 1993c:44 (in catalog of western Palearctic Crabronini and Oxybelini). – **As *Pseudomicroides elvirae***: Antropov, 2001a:812 (new combination, in revision of *Pseudomicroides*).

5. *fergusoni* (de Beaumont)

Belomicroides fergusoni de Beaumont, 1960a:24, ♂. Holotype: ♂, Greece: Rhodes Island: Kremasti (Lausanne). – R. Bohart and Menke, 1976:360 (in checklist of world Sphecidae); Pulawski, 1978:274 (in key to Sphecidae of European part of USSR); Leclercq, 1993c:43 (in catalog of western Palearctic Crabronini and Oxybelini); Józán, 2011:181 (in checklist of Sphecidae s.l. of Hungary). – **As *Pseudomicroides fergusoni***: Antropov, 2001a:814 (new combination, in revision of *Pseudomicroides*); Arens, 2002:513 (description of ♀, Greece: Pelopónnisos: Avia near Kalamata and Neokhori near Zakharo), 2009:1147 (Greece: Peloponnesus: plateau of Mantinea; preying on *Nomioides minutissimus* (Rossi), variation); Bitsch and Leclercq, 2009:227 (in supplement to vol. I of Faune de France, 1993: recently recorded from continental Greece); Leclercq, Bitsch, K. Schmidt, and Antropov *in* Bitsch, Antropov, Gayubo, Leclercq, Schmid-Egger, K. Schmidt, and Straka, 2021:356 (in revision of Sphecidae *sensu lato* of Europe).

6. *melas* (Antropov)

Belomicroides melas Antropov, 1994a:94, ♂. Holotype: ♂, Uzbekistan: 35 km southeast of Bukhara (ZMMU). – Kazenas, 2001b:35 (in checklist of Sphecidae of Kazakhstan and Central Asia); Medetov, Embergenov, Kholmatov, Elmurodova, Rakhimov, and Tajibaeva, 2024:424 (Uzbekistan: distribution). – **As *Pseudomicroides melas***: Antropov, 2001a:815 (new combination, in revision of *Pseudomicroides*).

7. *mochii* Antropov

Pseudomicroides mochii Antropov, 2001a:821, ♀. Holotype: ♀, Syria: Mezzé near Damascus (MSNT).

8. *nitidus* Antropov

Pseudomicroides nitidus Antropov, 2001:822, ♂. Holotype: ♂, Tunisia: Naftah (M. Schwarz coll., Ansfelden, Austria).
– Ben Khedher, Yildirim, Braham, and Chaaban, 2019:192 (known from Tunisia: Tozeur Province).

9. *olgae* (Kazenas)

Belomicroides olgae Kazenas, 1979c:172, ♀. Holotype: ♀, Kazakhstan: Betpak-dala Desert: 70 km northwest of Furmanovka (ZIL). – Antropov, 1994a:90 (in revision of Asian *Belomicroides*, description of ♂); Kazenas, 2001b:35 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:86 (geographic distribution, collecting localities in Kazakhstan), 2008a:103 (southeastern Kazakhstan: 17 km northwest of Bakanas). – **As *Pseudomicroides olgae***: Antropov, 2001a:816 (new combination, in revision of *Pseudomicroides*).

10. *pulawskii* Antropov

Pseudomicroides pulawskii Antropov, 2001a:823, ♀, ♂. Holotype: ♀, Mauritania: 25 km southwest of Tidjikdja (CAS).

11. *rostratus* (Krombein)

Belomicroides rostratus Krombein, 1982:31, ♀. Holotype: ♀, Sri Lanka: Hambantota District: Palatupana Tank (USNM). – Antropov, 1994a:92 (in revision of Asian *Belomicroides*). – **As *Pseudomicroides rostratus***: Antropov, 2001a:818 (new combination, in revision of *Pseudomicroides*).

12. *santschii* (von Schulthess)

Oxybelus santschii von Schulthess, 1925:187, ♀. Holotype: ♀, Tunisia: Hammamet (originally von Schulthess coll., now Zurich, may be lost). – von Schulthess, 1926b:219 (Tunisia: Hammamet); Pate, 1931a:113 (in key to *Belomicroides*). – **As *Belomicrus santschii***: von Schulthess, 1926b:219 (new combination, Tunisia: Hammamet). – **As *Belomicroides santschii***: R. Bohart and Menke, 1976:360 (in checklist of world Sphecidae); Leclercq, 1993c:43 (in catalog of western Palearctic Crabronini and Oxybelini); Roche, 2007a:104 and 2007b:7 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae). – **As *Pseudomicroides santschii***: Antropov, 2001:819 (new combination, in revision of *Pseudomicroides*).

13. *schwarzi* Antropov

Pseudomicroides schwarzi Antropov, 2001a:826, ♂. Holotype: ♂, Mali: 60 km southwest of Ségou (M. Schwarz coll., Ansfelden near Linz, Austria).

14. *sinaiticus* Antropov

Pseudomicroides sinaiticus Antropov, 2001a:827, ♂. Holotype: ♂, Egypt: Sinai: Wadi Gharandal 30 km northwest of Abu Zenima (CAS).

15. *zimini* (Gussakovskij)

Belomicrus zimini Gussakovskij, 1952:258, ♂. Lectotype: ♂, Tajikistan: Djilikul (ZIL), designated by Antropov, 1994a:90. – **As *Belomicroides zimini***: R. Bohart and Menke, 1976:360 (new combination, in checklist of world Sphecidae); Pulawski, 1978:274 (in key to Sphecidae of European part of USSR); Antropov, 1994a:90 (in revision of Asian *Belomicroides*); Leclercq, 1993c:43 (in catalog of western Palearctic Crabronini and Oxybelini); Kazenas, 2001b:35 (in checklist of Sphecidae of Kazakhstan and Central Asia), 2002a:86 (geographic distribution, collecting localities in Kazakhstan), 2008a:103 (southeastern Kazakhstan: near Kapchagay). – **As *Pseudomicroides zimini***: Antropov, 2001a:820 (new combination, in revision of *Pseudomicroides*).

sp.

Roche, 2007a:104 (in checklist of Egyptian Ampulicidae, Sphecidae, and Crabronidae).