

Last updated: 25 December 2025

## **PODAGRITUS**

*Podagritys* Spinola, 1851a:353. Type species: *Podagritys gayi* Spinola, 1851, by monotypy.

*Echuca* Pate, 1944b:354, as subgenus of *Podagritys*. Type species: *Crabro tricolor* F. Smith, 1856, by original designation.

*Parechuca* Leclercq, 1970a:91, as subgenus of *Podagritys*. Type species: *Podagritys neuqueni* Leclercq, 1957, by original designation.

*Chilichuca* Leclercq, 1981d:76, as subgenus of *Podagritys*. Type species: *Podagritys brieni* Leclercq, 1981d, by original designation.

*Ebisus* Tsuneki, 1983h:2, as subgenus of *Podagritys*. Type species: *Podagritys toxopeusi* Tsuneki, 1983, by original designation and monotypy.

Revisions: Leclercq, 1955 (Australian species), 1957f (subgenus *Echuca*), 1957h (subgenus *Podagritys* s.s.), 1981d (subgenera *Chilichuca* and *Podagritys* s.s.), 1982c (subgenus *Parechuca*), 1994b (redefinition of genus, Chilean species), 1998c (Australian species), 2000a (Latin America).

Common name in New Zealand: policemen wasps (Ferro et al., 1977:17).

### **1. *abaternus* Leclercq**

*Podagritys abaternus* Leclercq, 1982c:265, ♀. Holotype: ♀, Chile: Río Blanco, Caracautín (MCZ). – Leclercq, 1982c:252 (in key to subgenus *Parechuca*), 1994b:240, 241 (in key to *Podagritys* of Chile), 244 (bibliographic references, faunal records), 2000a:19, 28 (in key to *Podagritys* of South and Central America), 30 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae), 2005a:10 (correction to his 2002 catalog).

### **2. *acollae* Leclercq**

*Podagritys acollae* Leclercq, 1970b:275, ♀. Holotype: ♀, Peru: Junin Department: Acolla near Jauja (USNM). – R. Bohart and Menke, 1976:392 (in checklist of world Sphecidae); Leclercq, 1982c:256, 257 (in key to subgenus *Parechuca*), 266 (Peru; additional description), 2000a:19, 26 (in key to *Podagritys* of South and Central America), 30 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

*Podagritys exegetus* Leclercq, 1970:276, ♂. Holotype: ♂, Peru: Callanga (NHMW). Synonymized with *Podagritys acollae* by Leclercq, 1982c:266. – R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Dollfuss, 1989:11 (holotype in NHMW).

### **3. *aemulans* (Kohl)**

*Crabro aemulans* Kohl, 1905b:352, ♀, ♂. Syntypes: Chile: Concepción (NHMW). – Brèthes, 1910a:282 (Argentina: Mendoza Province); Jaffuel and Piri6n, 1926:375 (Chile: valle de Marga Marga). – **As *Rhopalum aemulans***: Chiappa, 2012:8 (new combination, Chile: Regi6n de Valpara6so: no specific locality, as *oemulans*). – **As *Podagritys aemulans***: Schrottky, 1913a:233 (new combination, Argentina: Mendoza); Leclercq, 1950b:9 (is a member of *Podagritys*, comments on Old World and New World *Podagritys*), 1951f:15 (Argentina, Ecuador, Peru), 1954c:193 (in catalog of world Crabronini), 1957h:3 (in key to subgenus *Podagritys* s.s.), 9 (in revision of subgenus *Podagritys* s.s., additional localities), 1970b:278 (Peru: near Jauja); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:68 (in key to subgenus *Podagritys* s.s. of South America), 79 (Brazil, Chile, Ecuador, Peru); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae); Radovi6, 1985:65 (sting apparatus analyzed); Ch. Porter, 1987:46 (Chile: Tarapac6 Region, in key to *Podagritys* of Tarapaca Region); Dollfuss, 1989:11 (type

material in NHMW); Leclercq, 1994b:243 (in key to *Podagritus* of Chile), 244 (bibliographic references, faunal records), 2000a:8, 12 (in key to *Podagritus* of South and Central America); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Dollfuss, 2004b:782 (Ecuador: Tungurahua: Baños); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

*Crabro gayi aequadoricus* Strand, 1911:154, ♂. Syntypes: Ecuador: Riobamba (MNHN). Synonymized with *Podagritus aemulans* by Leclercq, 1951f:15. – Strand, 1913c:28 (...)

#### 4. *albipes* (F. Smith)

*Rhopalum albipes* F. Smith, 1878a:7, ♂. Holotype or syntypes: ♂, New Zealand: Otago (BMNH). – Hutton, 1881:106 (in catalog of New Zealand Hymenoptera); W.F. Kirby, 1881a:41 (in checklist of Hymenoptera of New Zealand), 1884b:69 (in checklist of Hymenoptera of New Zealand); Hutton, 1904:98 (in list of New Zealand fauna); Leclercq, 1950b:6 (is a member of *Rhopalum*), 1954c:185 (in catalog of world Crabronini), 1955k:5, 8 (in key to *Rhopalum* of New Zealand), 9 (in revision of *Rhopalum* of New Zealand). – **As *Crabro albipes***: Dalla Torre, 1897:582 (new combination, in catalog of world Sphecidae). – **As *Podagritus albipes***: Leclercq, 1970a:87 (new combination); D. Miller, 1971:27 (...); R. Bohart and Menke, 1976:392 (in checklist of world Sphecidae); Callan, 1979:38 (New Zealand; prey: acalyptrate flies); Harris, 1990:55 (nesting habits); Valentine and Walker, 1991:39 (in catalog of New Zealand Hymenoptera); Harris, 1994:38 (in Fauna of New Zealand, cocoon, nest), 2000b:14 (visiting flowers of *Berberis vulgaris* L., Berberidaceae, in Bannockburn area, New Zealand), 2006:20 (nests containing Ephemeroptera and Diptera, prey ravaged by larvae of therevid fly *Anabarhynchus harrisi* Lyneborg).

#### 5. *aliciae* (R. Turner)

*Rhopalum aliciae* R. Turner, 1915a:90, ♀, ♂. Syntypes: Australia: Western Australia: Yallingup (BMNH). – R. Turner, 1915a:86 (in key to Australian *Rhopalum*). – **As *Podagritus aliciae***: Leclercq, 1950b:8 (new combination), 1954c:192 (in catalog of world Crabronini), 301 (nest and prey records), 1955f:313 (in key to Australian *Podagritus*), 327 (redescription); R. Bohart and Menke, 1976:392 (in checklist of world Sphecidae); Cardale, 1985:269 (in catalog of Australian Sphecidae); Leclercq, 1998c:290 (in key to Australian *Podagritus*), 293 (discussion of characters).

#### 6. *alisalis* Leclercq

*Podagritus alisalis* Leclercq, 2000a:30, ♀, ♂. Holotype: ♀, Argentina: Salta: El Alisal (Torino) – Leclercq, 2000a:9, 13 (in key to *Podagritus* of South and Central America); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

#### 7. *alutaceus* Leclercq

*Podagritus alutaceus* Leclercq, 1951f:4, ♀. Holotype: ♀, Ecuador: v. de Cloa (MNHN). – Leclercq, 1954c:192 (in catalog of world Crabronini), 1957h:5 (in key to subgenus *Podagritus* s.s.), 9 (in revision of subgenus *Podagritus* s.s.); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1982c:256, 257 (in key to subgenus *Parechuca*), 266 (Ecuador; description of ♂; additional description of ♀, prey), 2000a:19, 26 (in key to *Podagritus* of South and Central America), 31 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

#### 8. *anerus* Leclercq

*Podagritus anerus* Leclercq, 1955f:323, ♂. Holotype: ♂, Australia: Victoria: Scaford (National Mus. of Victoria). – Leclercq, 1955f:311 (in key to Australian *Podagritus*); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:270 (in catalog of Australian Sphecidae); Leclercq, 1998c:290 (in key to Australian *Podagritus*), 293 (synonymy).

*Podagritus alevinus* Leclercq, 1957e:3, ♀. Holotype: ♀, Australia: Victoria: Frankston (National Mus. of Victoria). Synonymized with *Podagritus anerus* by Leclercq, 1998c:293. – Leclercq, 1970b:271 (Australia); R. Bohart and Menke, 1976:392 (in checklist of world Sphecidae); Cardale, 1985:269 (in catalog of Australian Sphecidae).

### 9. *apivorus* Pulawski

*Podagritus apivorus* Pulawski, 2023:123, ♀, ♂. Holotype: ♀, Australia: Western Australia: Yanchep National Park at 31.5063°S 115.6811°E (WAM). – Houston and Stuart, 2023:132 (nesting habits).

### 10. *apostolus* Leclercq ♀

*Podagritus apostolus* Leclercq, 1981d:79, ♀, ♂ (♀ = *Podagritus ypirangae*). Holotype: ♂, Paraguay: Asunción (AMNH). Paratype: Brazil. – Leclercq, 1981d:71 and 73 (in key to subgenus *Podagritus* s.s. of South America), 2000a:9, 13 (in key to *Podagritus* of South and Central America), 31 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae, as *apostulus*), 2005a:10 (correction to his 2002 catalog).

As *Podagritus nigriventris* (♀ only): Leclercq, 1957h:6 (illustration of female clypeus), 7 (in key to subgenus *Podagritus* s.s.), 13 (bibliographic references, additional locality records), corrected to *Podagritus apostolus* by Leclercq, 2000a:31.

### 11. *arechavaletai* (Brèthes)

*Crabro arechavaletai* Brèthes, 1909b:66, ♀ (as *Arechavaletai*, incorrect original capitalization). Holotype: ♀, Uruguay: Banda Oriental: no specific locality (MACN). – Genise, 1990:26 (type material in MACN). – As *Rhopalum arechavaletai*: Leclercq, 1954c:185 (new combination, in catalog of world Crabronini). – As *Podagritus arechavaletai*: Leclercq in R. Bohart and Menke, 1976:393 (new combination, in checklist of world Sphecidae); Leclercq, 1981d:71, 75 (in key to subgenus *Podagritus* s.s. of South America), 80 (Argentina: Buenos Aires; comparison with *Podagritus joergenseni*), 2000a:10, 16 (in key to *Podagritus* of South and Central America), 31 (bibliographic references, additional locality records, color variation); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

### 12. *aricae* Leclercq

*Podagritus aricae* Leclercq, 1957h:10, ♂. Holotype: ♂, Chile: Tarapacá: Arica (USNM). – Leclercq, 1957h:2 (in key to subgenus *Podagritus* s.s.), 1970b:278 (Peru: Jauja area); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:69 (in key to subgenus *Podagritus* s.s. of South America), 80 (bibliographic references, locality records, diagnostic characters); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae); Ch. Porter, 1987:46 (Chile: Tarapacá Region, in key to *Podagritus* of Tarapaca Region); Dollfuss, 1989:11 (paratype in NHMW); Solervicens and Henry, 1989:295 (paratype in Inst. Entom. Univ. Metropol., Santiago de Chile); Leclercq, 1994b:244 (in key to *Podagritus* of Chile), 245 (bibliographic references, locality records, as *Podagritus aricae aricae*), 2000a:8, 12 (in key to *Podagritus* of South and Central America), 31 (bibliographic references); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

### *ssp. carrascoi* Fritz

*Podagritus carrascoi* Fritz, 1971a:110, ♀, ♂. Holotype: ♂, Peru: Cuzco (originally M. Fritz coll., now MACN). – As *Podagritus aricae carrascoi*: Leclercq in R. Bohart and Menke, 1976:393 (new status, in checklist of world Sphecidae); Leclercq, 1981d:69 (in key to subgenus *Podagritus* s.s. of South America), 80 (bibliographic references, locality records); Fritz, 1993:17 (holotype in M. Fritz coll.), Leclercq, 2000a:8, 12 (in key to *Podagritus* of South and Central America); Amarante, 2002:31 (in catalog of Neotropical Crabronidae), 2005a:10 (correction to his 2002 catalog); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

### 13. *australiensis* Tsuneki

*Podagritus australiensis* Tsuneki, 1977d:5, ♀, ♂. Holotype: ♂, Australia: New South Wales: Mount Victoria (TMB). – Cardale, 1985:270 (in catalog of Australian Sphecidae); Leclercq, 1998c:292 (in key to Australian *Podagritus*), 293 (analysis of diagnostic characters).

**14. bocainus Leclercq**

*Podagritus bocainus* Leclercq, 1982c:267, ♀, ♂. Holotype: ♀, Brazil: São Paulo: Serra de Bocãina (AMNH). – Leclercq, 1982c:261 (in key to subgenus *Parechuca*), 2000a:20, 28 (in key to *Podagritus* of South and Central America), 32 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Alves-dos-Santos, 2003:330 (Brazil: University of São Paulo campus).

**15. bordai Fritz**

*Podagritus bordai* Fritz, 1971a:112, ♀, ♂. Holotype: ♂, Bolivia: Cochamba (originally M. Fritz coll., now MACN). – R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:68 (in key to subgenus *Podagritus* s.s. of South America), 81 (additional description); Fritz, 1993:17 (holotype in M. Fritz coll.); Leclercq, 2000a:8, 12 (in key to *Podagritus* of South and Central America), 32 (bibliographic references, locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae), 2005a:10 (correction to his 2002 catalog).

**16. brethesi Leclercq**

*Podagritus brethesi* Leclercq, 1951f:12, ♂ (as *brèthesi*, incorrect original diacritic mark). Holotype: ♂, Brazil: Rio Grande do Sul: no specific locality (MHNG). – Leclercq, 1954c:192 (in catalog of world Crabronini), 1957h:5 (in key to subgenus *Podagritus* s.s.), 12 (in revision of subgenus *Podagritus* s.s.); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:71 and 75 (in key to subgenus *Podagritus* s.s. of South America), 81 (variation), 2000a:10, 16 (in key to *Podagritus* of South and Central America), 32 (bibliographic references, locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

**17. brieni Leclercq**

*Podagritus brieni* Leclercq, 1981d:76, ♀, ♂. Holotype: ♀, Chile: Coquimbo: Baños del Toro (originally M. Fritz coll., now MACN). – Leclercq, 1981d:67 (in key to subgenus *Chilichuca*); Fritz, 1993:17 (holotype in M. Fritz coll.); Leclercq, 1994b:243 (in key to *Podagritus* of Chile), 245 (bibliographic references, faunal records), 2000a:17 (in key to *Podagritus* of South and Central America); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**18. burnsi Leclercq**

*Podagritus burnsi* Leclercq, 1955f:317, ♀, ♂. Holotype: ♀, Australia: Victoria: Melbourne (National Mus. of Victoria). – Leclercq, 1955f:310 (in key to Australian *Podagritus*), 1957e:3 (Australia: Victoria: Kiata); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:270 (in catalog of Australian Sphecidae); Leclercq, 1998c:288 (in key to Australian *Podagritus*), 294 (collecting localities).

**19. caelebs Leclercq**

*Podagritus caelebs* Leclercq, 2000a:32, ♂. Holotype: ♂, Peru: Amazonas: Leymebamba (BMNH). – Leclercq, 2000:27 (in key to *Podagritus* of South and Central America); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

**20. carbonicolor (Dalla Torre)**

*Crabro carbonarius* F. Smith, 1856:424, ♀, junior primary homonym of *Crabro carbonarius* Dahlbom, 1838. Holotype or syntypes: ♀, New Zealand: no specific locality (BMNH). – Gourlay, 1964:49 (prey: *Cutilia truncata* Brunn., a cockroach, possibly in error). – **As *Rhopalum carbonarium***: F. Smith, 1867:484 (new combination) [citation from Dalla Torre, 1897:586, his reference is wrong]; Hutton, 1881:105 (in catalog of New Zealand Hymenoptera); W.F. Kirby, 1881a:40 (in checklist of Hymenoptera of New Zealand), 1884b:69 (in checklist of Hymenoptera of New Zealand); Hutton, 1904:98 (in list of New Zealand fauna); Leclercq, 1950b:6 (is a member of *Rhopalum*), 1955k:11 (as new synonym of *Rhopalum cora*); Davies, 1986:61 (New Zealand: Hawkes Bay).

*Crabro carbonicolor* Dalla Torre, 1897:586. Substitute name for *Crabro carbonarius* F. Smith. – **As *Rhopalum carbonicolor***: Kirkaldy, 1910:131 (new combination, Dalla Torre's substitute name recorded); Leclercq, 1954c:184 (in catalog of world Crabronini). – **As *Podagritus carbonicolor***: R. Bohart and Menke, 1976:393 (new combination,

in checklist of world Sphecidae, as synonym of *Podagritus cora*); Harris, 1994:41 (new combination, in Fauna of New Zealand, nest), 2000a:76 (preying on large Calliphoridae and Tabanidae, nest structure).

*Rhopalum swalei* Leclercq, 1955k:12, ♂. Holotype: ♂, New Zealand: Putavuru (BMNH). Synonymized with *Podagritus carbonicolor* by Harris, 1994:41. – Leclercq, 1955k:7 (in key to New Zealand *Rhopalum*). – **As *Podagritus swalei***: R. Bohart and Menke, 1976:393 (new combination, in checklist of world Sphecidae); Callan, 1979:39 (New Zealand); Valentine and Walker, 1991:40 (in catalog Hymenoptera of New Zealand).

### 21. *carolus* Leclercq

*Podagritus carolus* Leclercq, 1955f:319, ♀. Holotype: ♀, Australia: South Australia: Killalpaninna 100 mi. east of Lake Eyre (BMNH). – Leclercq, 1955f:308 (in key to Australian *Podagritus*); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:270 (in catalog of Australian Sphecidae); Leclercq, 1998c:288 (in key to Australian *Podagritus*), 294 (bibliographic reference).

### 22. *catharinae* Fritz

*Podagritus catharinae* Fritz, 1971a:109, ♂. Holotype: ♂, Brazil: Santa Catarina: no specific locality (Zurich). – R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:73 (in key to subgenus *Podagritus* s.s. of South America), 81 (in revision of South American *Podagritus*), 2000a:13 (in key to *Podagritus* of South and Central America), 33 (bibliographic references); Amarante, 2002:31 (in catalog of Neotropical Crabronidae), 2005a:10 (correction to his 2002 catalog).

### 23. *chambersi* Harris

*Podagritus chambersi* Harris, 1994:43, ♀, ♂. Holotype: ♂, New Zealand: Taupo (New Zealand Arthropod Collection, Auckland, N.Z.). – Harris, 1994:43 (nest), 1995:51 (nest structure and prey), 2002b:22 (nest structure).

### 24. *cloudatus* Leclercq

*Podagritus cloudatus* Leclercq, 1982c:267, ♀, ♂. Holotype: ♀, Colombia: Magdalena: above San Pedro de la Sierra southeast of Santa Marta (BMNH). – Leclercq, 1982c:259 (in key to subgenus *Parechuca*), 2000a:18, 24 (in key to *Podagritus* of South and Central America), 33 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

### 25. *clypeatus* Pulawski

*Podagritus clypeatus* Pulawski, 2025b:220, ♀, ♂. Holotype: ♂, Australia: Western Australia: : 5 km west-southwest of Paynes Find at 29°18'S 117°39'E (ANIC).

### 26. *colchagae* Leclercq

*Podagritus colchagae* Leclercq, 1994b:245, ♀. Holotype: ♀, Chile: Colchaga: Río Tinguiririca (FSAG). – Leclercq, 1994b:243 (in key to *Podagritus* of Chile), 2000a:17 (in key to *Podagritus* of South and Central America), 33 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Leclercq, 2007g:194 (holotype in FSAG).

### 27. *consideratus* Leclercq

*Podagritus consideratus* Leclercq, 1982c:267, ♀, ♂. Holotype: ♀, Ecuador: Tungurahua: Baños (BMNH). – Leclercq, 1982c:255 (in key to subgenus *Parechuca*), 2000a:18, 26 (in key to *Podagritus* of South and Central America), 33 (bibliographic references, variation); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

### 28. *cora* (Cameron)

*Crabro cora* Cameron, 1888b:181, ♂, actually ♀. Lectotype: ♀, New Zealand: Greymouth (BMNH), designated by Harris, 1994:45. – Dalla Torre, 1897:592 (listed incorrectly from India). – **As *Rhopalum cora***: Hutton, 1904:98 (new combination, in list of New Zealand fauna); Leclercq, 1954c:185 (in catalog of world Crabronini), 1955k:5, 7 (in key to *Rhopalum* of New Zealand), 11 (New Zealand, redescription). – **As *Podagritus cora***: Leclercq, 1970b:276 (new

combination; New Zealand: Auckland); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Callan, 1979:37 (New Zealand; available information reviewed); Harris, 1990:55 (nesting habits); Valentine and Walker, 1991:40 (in catalog Hymenoptera of New Zealand); Harris, 1994:43 (good species, in Fauna of New Zealand, nest), 2000b:14 (New Zealand: visiting flowers of *Berberis vulgaris* L., Berberidaceae, in Bannockburn area), 2002a:76 (prey: miscellaneous Diptera, nest structure, description of mature larva), 2006:20 (nests containing Ephemeroptera and Diptera prey ravaged by larvae of therevid fly *Anabarhynchus harrisi* Lyneborg).

*Crabro jocosus* Cameron, 1898a:43, sex not stated. Syntypes: New Zealand: no specific locality (BMNH). Synonymized with *Podagritus cora* by Leclercq, 1955k:11. – As *Rhopalum jocosum*: Leclercq, 1950:6 (new combination), 1954c:185 (in catalog of world Crabronini).

### 29. *corrientis* Leclercq

*Podagritus corrientis* Leclercq, 2000a:34, ♀, ♂. Holotype: ♀, Argentina: Corrientes: Ytuzaingo (originally M. Fritz coll., now FSAG). – Leclercq, 2000a:10, 16 (in key to *Podagritus* of South and Central America); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Leclercq, 2007g:194 (holotype in FSAG).

### 30. *cuencae* Leclercq

*Podagritus cuencae* Leclercq, 1982c:268, ♀, ♂. Holotype: ♀, Ecuador: Cuenca: 52 km south of Cuenca (MCZ). – Leclercq, 1982c:256, 257 (in key to subgenus *Parechuca*), 2000a:19, 26 (in key to *Podagritus* of South and Central America), 34 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

### 31. *cuevasus* Leclercq

*Podagritus cuevasus* Leclercq, 1982c:268, ♀. Holotype: ♀, Colombia: Huila: Las Cuevas de los Gacheros (BMNH). – Leclercq, 1982c:260 (in key to subgenus *Parechuca*), 2000a:18 (in key to *Podagritus* of South and Central America), 34 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

### 32. *curatus* Leclercq

*Podagritus curatus* Leclercq, 1982c:268, ♂. Holotype: ♂, Bolivia: La Paz: Chulumani (BMNH). – Leclercq, 1982c:260 (in key to subgenus *Parechuca*), 2000a:25 (in key to *Podagritus* of South and Central America), 34 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

### 33. *cuzcosus* Leclercq

*Podagritus cuzcosus* Leclercq, 1982c:268, ♂. Holotype: ♂, Peru: Cuzco (AEI). – Leclercq, 1982c:263, 264 (in key to subgenus *Parechuca*); Dollfuss, 1989:11 (paratype in NHMW, as *cuzosus*); Leclercq, 2000a:22, 29 (in key to *Podagritus* of South and Central America), 34 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

### 34. *cynericus* Leclercq

*Podagritus cynericus* Leclercq, 1981d:82, ♀, ♂. Holotype: ♀, Argentina: Catamarca: El Rodeo (IML). – Leclercq, 1981d:70, 74 (in key to subgenus *Podagritus* s.s. of South America), 2000a:9, 14 (in key to *Podagritus* of South and Central America), 34 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

### 35. *diegotus* Leclercq

*Podagritus diegotus* Leclercq, 1982c:269, ♀, ♂. Holotype: ♀, Colombia: Magdalena: Río Don Diego north of Sierra de Santa Marta (BMNH). – Leclercq, 1982c:253, 255 (in key to subgenus *Parechuca*), 2000a:18, 26 (in key to *Podagritus* of South and Central America), 35 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**36. digyalos Harris**

*Podagritus digyalos* Harris, 1994:46, ♀, ♂. Holotype: ♂, New Zealand: Fraser's Dam Creek in Obelisk Range (Otago Museum). – Harris, 1994:46 (nest), 2000a:75 (nest structure, preying on Empididae, Ephydriidae, Pseudopomyzidae, Simuliidae, and Syrphidae), 2002b:23 (preying on small Diptera, nest structure).

**37. doreeni Leclercq**

*Podagritus doreeni* Leclercq, 1955f:313, ♀. Holotype: ♀, Australia: Victoria: Trentham (National Mus. of Victoria). – Leclercq, 1955f:309 (in key to Australian *Podagritus*), 1957e:2 (Australian Capital Territory, additional description); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:270 (in catalog of Australian Sphecidae); Leclercq, 1998c:288 (in key to Australian *Podagritus*), 294 (bibliographic reference).

**38. ecuadoris Leclercq**

*Podagritus ecuadoris* Leclercq, 2000a:35, ♀. Holotype: ♀, Ecuador: Tungurahua: Baños (BMNH). – Leclercq, 2000a:29 (in key to *Podagritus* of South and Central America); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**39. edgarus Leclercq**

*Podagritus edgarus* Leclercq, 1957e:5, ♂. Australia: New South Wales: 35 mi. northwest of Nyngan (ANIC). – R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:270 (in catalog of Australian Sphecidae); Leclercq, 1998c:291 (in key to Australian *Podagritus*), 294 (original description may contain errors).

**40. erythropus (Brèthes)**

*Crabro erythropus* Brèthes, 1913:138, ♀. Holotype or syntypes: ♀, Argentina: Patagonia: Chacras de Coria (MACN). – Genise, 1990:26 (type material in MACN). – **As *Podagritus erythropus***: Leclercq, 1951f:16 (new combination, description of ♂), 1954c:193 (in catalog of world Crabronini), 1957h:2 (in key to subgenus *Podagritus* s.s.), 9 (in revision of subgenus *Podagritus* s.s., diagnostic characters), 1970b:279 (Argentina: Mendoza Province); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:69 (in key to subgenus *Podagritus* s.s. of South America), 82 (Argentina, Bolivia; as *erythropus erythropus*); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae); Leclercq, 2000a:8, 12 (in key to *Podagritus* of South and Central America), (bibliographic references, additional locality records, *Podagritus eremurus* may be an individual dark form of *P. erythropus*); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

**ssp. eremurus Leclercq**

*Podagritus erythropus eremurus* Leclercq, 1981d:83, ♀, ♂. Holotype: ♀, Argentina: Tucumán: Tafi del Valle (AEI). – Leclercq, 1981d:69 (in key to subgenus *Podagritus* s.s. of South America), 2000a:8, 12 (in key to *Podagritus* of South and Central America), 35 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

**41. ferrugineus Pulawski**

*Podagritus ferrugineus* Pulawski, 2025b:224, ♀. Holotype: ♀, Australia: Western Australia: 7 km N of Booloogooro Homestead which is at 24°39'S 113°42'E (WAM).

**42. firmatus Leclercq**

*Podagritus firmatus* Leclercq, 1998c:294, ♀. Holotype: ♀, Australia: Western Australia: 2 km west-southwest of Noongar at 31°21'S 118°57'E (ANIC). – Leclercq, 1998c:290 (in key to Australian *Podagritus*).

**43. fulvohirtus (Cameron)**

*Crabro fulvohirtus* Cameron, 1891a:153, sex not stated (as *fulvo-hirtus*, incorrect original hyphenation). Holotype or syntypes: ♀, Mexico: Guerrero: Xautipa (BMNH). – Dalla Torre, 1897:602 (in catalog of world Sphecidae). – **As *Crossocerus fulvohirtus***: Leclercq, 1950b:17 (new combination). – **As *Podagritus fulvohirtus***: Ashmead, 1899e:220 (new combination, in checklist of North American Crabronidae); Leclercq, 1954c:193 (in catalog of world

Crabronini); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1982c:262 (in key to subgenus *Parechuca*), 269 (belongs in *Podagritys*; Colombia, Costa Rica), 2000a:21, 27 (in key to *Podagritys* of South and Central America), 36 (bibliographic references, additional locality records); Hanson and Menke, 1995:644 (known from Costa Rica); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Ruíz Cancino, Coronado Blanco, Varela Fuente, and Horta Vega, 2002:664 (in checklist of Mexican Sphecidae); Cambra, 2005:102 (Panamá).

#### 44. *garcianus* Leclercq

*Podagritys garcianus* Leclercq, 1982c:269, ♀, ♂. Holotype: ♀, Peru: Cuzco (MCZ). – Leclercq, 1982c:258 (in key to subgenus *Parechuca*), 2000a:17, 23 (in key to *Podagritys* of South and Central America), 36 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

#### 45. *gastricus* Leclercq

*Podagritys gastricus* Leclercq, 1982c:269, ♀, ♂. Holotype: ♀, Argentina: Jujuy: Porto Lozano (MCZ). – Leclercq, 1982c:258 (in key to subgenus *Parechuca*), 2000a:17, 23 (in key to *Podagritys* of South and Central America), 36 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

#### 46. *gayi* Spinola

*Podagritys gayi* Spinola, 1851a:357, ♀, ♂ (as *Gayi*, incorrect original capitalization). Syntypes: Chile: "Coquimbo, etc." (Torino). – Kieffer and Herbst, 1909:122 (visiting galls of *Lecanium resinatum* Kieffer and Herbst, Kerriidae); Jaffuel and Piri6n, 1926:375 (Chile: valle de Marga Marga); Leclercq, 1951f:16 (Argentina), 1954c:193 (in catalog of world Crabronini), 301 (nest and prey records), 1957h:3 (in key to subgenus *Podagritys* s.s.), 12 (in revision of subgenus *Podagritys* s.s.; additional locality records); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Casolari and Casolari Moreno, 1980:121 (specimen in M. Spinola collection); Leclercq, 1981d:72 and 73 (in key to subgenus *Podagritys* s.s. of South America), 83 (Argentina, Chile; variation); Steiner, 1986:107 (references to publications on nesting habits); Leclercq, 1994b:243 (in key to *Podagritys* of Chile), 245 (bibliographic references, faunal records), 2000a:10, 14 (in key to *Podagritys* of South and Central America), 36 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Dollfuss, 2004b:782 (Chile: Caut6n Parque Nacional: Las Paraguas, Coquimbo: Vicu6na Elgui); Chiappa, 2012:8 (Chile: Regi6n de Valpara6so: no specific locality); Pagliano, 2014:86 (specimen in M. Spinola collection, Torino). – *As Crabro gayi*: F. Smith, 1856:393 (new combination, in catalog of Hymenoptera in British Museum); Reed, 1894:647 (in revision of Chilean Aculeata); Dalla Torre, 1897:602 (in catalog of world Sphecidae); Strand, 1911a:153 (Ecuador, determination tentative); Ruiz Pereira, 1924:103 (Chile: Cerro San Crist6bal); Janvier, 1928:69 (nesting habits), 1929:418 (sleeping habits); Gazulla and Ruiz Pereira, 1929:300 (Chile: Hacienda de "Las Mercedes"); Ru6z Pereira and Stuardo, 1936:321 (Chile: Las Termas de Chill6n); Piek and Spanjer, 1986:183 (in list of Sphecidae with known prey).

*Rhopalum patagonicum* Holmberg, 1903:470, ♀, ♂. Syntypes: Argentina: Neuqu6n: no specific locality (MACN). Synonymized with *Podagritys gayi* by Leclercq in R. Bohart and Menke, 1976:393. – Schrottky, 1913a:233 (Argentina: Neuqu6n); Leclercq, 1954c:185 (in catalog of world Crabronini).

#### 47. *gibber* Leclercq

*Podagritys gibber* Leclercq, 1998c:294, ♀, ♂. Holotype: ♀, Australia: Queensland: North Stradbroke Island: Dunwich (Kansas Univ.). – Leclercq, 1998c:290 (in key to Australian *Podagritys*).

#### 48. *guevarus* Leclercq

*Podagritys guevarus* Leclercq, 1981d:84, ♀. Holotype: ♀, Argentina: Mendoza: Potrerillos (IML). – Leclercq, 1981d:72 (in key to subgenus *Podagritys* s.s. of South America), 2000a:10, 14 (in key to *Podagritys* of South and Central America), 36 (bibliographic references, additional locality record); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

**49. heteroceru (Mantero)**

*Crabro heteroceru* Mantero, 1902:201, ♂. Syntypes: ♂, Argentina: Patagonia: Río Santa Cruz (MSNG). – **As *Corynopu heteroceru***: Holmberg, 1903:474 (new combination, Argentina); Schrottky, 1913a:234 (Argentina: Santa Cruz). – **As *Rhopalum heteroceru***: Leclercq, 1954c:186 (new combination, in catalog of world Crabronini); R. Bohart and Menke, 1976:389 (in checklist of world Sphecidae); Amarante, 2005:8 (in addendum to his 2002 catalog of Neotropical Crabronidae and Sphecidae). – **As *Podagrītus heteroceru***: Leclercq, 2000a:19, 27 (in key to *Podagrītus* of South and Central America), 37 (new combination, recognition characters); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

*Podagrītus claru* Leclercq, 1957f:21, ♀. Holotype: ♀, Argentina: Neuquén: Piedra Pintada (Museo de la Plata). Synonymized with *Podagrītus heteroceru* by Leclercq, 2000a:37. – Leclercq, 1957f:12 (in key to subgenus *Echucu*), 1982c:258 (in key to subgenus *Parechucu*), 267 (citation).

**50. imbellis (R. Turner)**

*Rhopalum tricolor imbellis* R. Turner, 1915a:92, ♀, ♂. Syntypes: Australia: Western Australia: no specific locality (BMNH). – R. Turner, 1915a:86 (in key to Australian *Rhopalum*). – **As *Rhopalum imbellis***: R. Turner, 1918a:93 (new status, diagnostic characters). – **As *Podagrītus imbellis***: Leclercq, 1950b:8 (new combination, new status, as *imbellis*), 1954c:191 (in catalog of world Crabronini), 1955f:312 (in key to Australian *Podagrītus*), 325 (redescription), 1957e:5 (Western Australia), 1970b:271 (additional localities); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:270 (in catalog of Australian Sphecidae, as *imbellis*); Leclercq, 1998c:292 (in key to Australian *Podagrītus*), 295 (locality records).

**51. joergenseni (Brèthes)**

*Crabro joergenseni* Brèthes, 1913:136, ♀ (as *Jörgenseni*, incorrect original capitalization and diacritic mark). Holotype or syntypes: ♀, Argentina: Córdoba: Arias (MACN). – Genise, 1990:26 (type material in MACN). – **As *Podagrītus joergenseni***: Leclercq, 1954c:193 (new combination, in catalog of world Crabronini), 1957h:5 (in key to subgenus *Podagrītus* s.s.), 13 (in revision of subgenus *Podagrītus* s.s., comparison with related species; additional localities), 1970b:279 (Argentina: Entre Ríos Province), 1981d:71, 75 (in key to subgenus *Podagrītus* s.s. of South America), 84 (as full species; Argentina, Brazil), 2000a:10, 16 (in key to *Podagrītus* of South and Central America), 37 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae); Dollfuss, 2004b:782 (Argentina: Entre Ríos: Pronunciamiento). – **As *Podagrītus arechavaletai joergenseni***: Leclercq in R. Bohart and Menke, 1976:393 (new status, in checklist of world Sphecidae).

**52. jordaonis Leclercq**

*Podagrītus jordaonis* Leclercq, 2000a:37, ♀, ♂. Holotype: ♀, Brazil: São Paulo: Campos de Jordão (Mus. Zoologia Universidade São Paulo). – Leclercq, 2000a:11, 15 (in key to *Podagrītus* of South and Central America); Amarante, 2002:31 (in catalog of Neotropical Crabronidae). 2005a:10 (correction to his 2002 catalog); Menezes de Jesus, Onody, dos Santos Ramos, de Oliveira Andrade, Doria dos Santos, and Ferreira Brandão, 2019:2 (holotype in Mus. Zool. Univ. São Paulo).

**53. kelseyi Leclercq**

*Podagrītus kelseyi* Leclercq, 1998c:295, ♀. Holotype: ♀, Australia: Western Australia: near Miaboolya Beach via Carnarvon (ANIC). – Leclercq, 1998c:288 (in key to Australian *Podagrītus*).

**54. kiatae Leclercq**

*Podagrītus kiatae* Leclercq, 1955f:329, ♀, ♂. Holotype: ♀, Australia: Victoria: Kiata (National Mus. of Victoria). – Leclercq, 1955f:313 (in key to Australian *Podagrītus*), 1970b:272 (South Australia, Western Australia; variation); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:271 (in catalog of Australian Sphecidae); K. Walker, Naumann, Austin, Taylor, and Cardale, 1992:48 (in catalog of insects of Tasmania); Leclercq, 1998c:293 (in key to Australian *Podagrītus*), 296 (locality records, taxonomic problems).

**55. *krombeini* Leclercq**

*Podagritus krombeini* Leclercq, 1955f:320, ♂. Holotype: ♂, Australia: New South Wales: Sydney (USNM). – Leclercq, 1955f:310 (in key to Australian *Podagritus*), 1957e:3 (Australia: New South Wales); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:270 (in catalog of Australian Sphecidae); Leclercq, 1998c:289 (in key to Australian *Podagritus*), 297 (locality records).

**56. *leptospermi* (R. Turner)**

*Rhopalum leptospermi* R. Turner, 1915a:88, ♀, ♂. Syntypes: Australia: Western Australia: Yallingup and Warren River (BMNH). – R. Turner, 1915a:86 (in key to Australian *Rhopalum*). – **As *Podagritus leptospermi***: Leclercq, 1950b:8 (new combination), 1954c:191 (in catalog of world Crabronini), 1955f:309 (in key to Australian *Podagritus*), 316 (redescription, additional localities), 1957e:3 (additional Australian localities, variation), 1970b:272 (New South Wales, Western Australia; prey: Stratiomyiidae); Evans and Matthews, 1970:154 (nesting in soil); 1971a:1 (nesting habits); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:271 (in catalog of Australian Sphecidae); K. Walker, Naumann, Austin, Taylor, and Cardale, 1992:48 (in catalog of insects of Tasmania); Leclercq, 1998c:288 (in key to Australian *Podagritus*), 297 (locality records).

**57. *longinodus* (Spinola)**

*Physoscelus longinodus* Spinola, 1851a:359, ♀, ♂ (♂ = *Podagritus valdiviae* Leclercq). Lectotype: ♀, Chile: Saturno, Elqui (Torino). – Casolari and Casolari Moreno, 1980:121 (specimens in M. Spinola collection, Torino); Pagliano, 2014:85 (specimens in M. Spinola collection, Torino, are *Podagritus longinodus* including lectotype, and *Podagritus valdiviae*). – **As *Crabro longinodus***: F. Smith, 1856:394 (new combination, in catalog of Hymenoptera in British Museum); Reed, 1894:648 (in revision of Chilean Sphecidae); Dalla Torre, 1897:609 (in catalog of world Sphecidae); Ruiz Pereira, 1924:103 (Chile: Cerro San Cristóbal); Jaffuel and Piri6n, 1926:375 (Chile: valle de Marga Marga); Janvier, 1928:75 (nesting habits); Piek and Spanjer, 1986:183 (in list of Sphecidae with known prey). – **As *Rhopalum longinodum***: Leclercq, 1954c:186 (new combination, in catalog of world Crabronini), 298 (nest and prey records); R. Bohart and Menke, 1976:389 (in checklist of world Sphecidae); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae); Steiner, 1986:106 (references to publications on nesting habits); Chiappa, 2012:8 (Chile: Regi6n de Valpara6so: no specific locality). – **As *Podagritus longinodus***: Leclercq, 1994b:240, 242 (in key to *Podagritus* of Chile), 246 (new combination, bibliographic references, locality records), 2000a:20, 27 (in key to *Podagritus* of South and Central America), 38 (bibliographic references, additional locality records); Melo, 2000:89 (use of nest by more than one female); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

*Crabro chilensis* Reed, 1894:648, ♀. Holotype: ♀, Chile: Quillota (depository unknown). Synonymized with *Podagritus longinodus* by Bohart and Menke, 1976:389, synonymy confirmed by Leclercq, 1994b:246.

*Podagritus droserus* Leclercq, 1957f:14, ♀, ♂. Holotype: ♀, Chile: no specific locality (Inst. Roy. Sci. Nat. Belgique according to Leclercq, 1957f:14, but FSAG according to Leclercq, 2007g:194). Synonymized with *Rhopalum longinodum* by Leclercq in R. Bohart and Menke, 1976:389, synonymy confirmed by Leclercq, 1994b:246. – Leclercq, 1957f:11 (in key to subgenus *Echuca*), 2007g:194 (holotype in FSAG). – **As *Rhopalum droserum***: Leclercq, 1970a:92 and 97 (in key to Latin American *Rhopalum*).

**58. *lynchii* (Holmberg)**

*Rhopalum lynchii* Holmberg, 1903:471, ♂ (as *Lynchii*, incorrect original capitalization). Syntypes: ♂, Argentina: Buenos Aires: Chacabuco (MACN). – **As *Rhopalum lynchii***: Schrottky, 1913a:233 (Argentina: Buenos Aires); Leclercq, 1954c:185 (new combination, in catalog of world Crabronini). – **As *Podagritus lynchii***: Leclercq in R. Bohart and Menke, 1976:393 (new combination, in checklist of world Sphecidae), 1981d:70, 73 (in key to subgenus *Podagritus* s.s. of South America), 84 (holotypes of *Podagritus lynchi* and *P. pamparum* examined, additional locality records), 2000a:9, 13 (in key to *Podagritus* of South and Central America), 38 (bibliographic references); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

*Crabro pamparum* Brèthes, 1913:137, ♂. Holotype or syntypes: ♂, Argentina: La Pampa: no specific locality (MACN). Synonymized with *Podagritus lynchii* by Leclercq in R. Bohart and Menke, 1976:393. – Genise, 1990:26 (type material in MACN). – **As *Podagritus pamparum***: Leclercq, 1954c:193 (new combination, in catalog of world Crabronini), 1957h:8 (in key to subgenus *Podagritus* s.s.), 14 (in revision of subgenus *Podagritus* s.s., additional localities).

### 59. *magellanus* Leclercq

*Podagritus magellanus* Leclercq, 1957f:21, ♂. Holotype: ♂, Tierra del Fuego: Río Grande, Estancia Viamonte (BMNH). – Leclercq, 1957f:11 (in key to subgenus *Echuca*), 1994b:240, 242 (in key to *Podagritus* of Chile), 248 (retransferred from *Rhopalum* to *Podagritus*, recognition characters), 2000a:20, 27 (in key to *Podagritus* of South and Central America), 38 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae). – **As *Rhopalum magellanus***: Leclercq in R. Bohart and Menke, 1976:21 (new combination, in checklist of world Sphecidae); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae).

### 60. *marcellus* Leclercq

*Podagritus marcellus* Leclercq, 1955f:323. ♀, ♂. Holotype: ♀, Australia: Western Australia: Dongarra (BMNH). – Leclercq, 1955f:312 (in key to Australian *Podagritus*), 1957e:5 (Western Australia: Yallingup); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:271 (in catalog of Australian Sphecidae); Leclercq, 1998c:291 (in key to Australian *Podagritus*), 297 (locality records, variation).

### 61. *martini* Leclercq

*Podagritus martini* Leclercq, 2000a:38, ♂. Holotype: ♂, Peru: Amazonas: Rodriguez de Mendoza (BMNH). – Leclercq, 2000a:23 (in key to *Podagritus* of South and Central America); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

### 62. *matucanae* Leclercq

*Podagritus matucanae* Leclercq, 2000a:40, ♀, ♂. Holotype: ♂, Peru: Lima: Matucana (BMNH). – Leclercq, 2000a:18, 25 (in key to *Podagritus* of South and Central America); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

### 63. *meloii* Leclercq

*Podagritus meloi* Leclercq, 2000a:40, ♀, ♂. Holotype: ♂, Costa Rica: Zurqui (G. Melo coll.). – Leclercq, 2000a:20, 23 (in key to *Podagritus* of South and Central America); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

### 64. *meridensis* Leclercq

*Podagritus meridensis* Leclercq, 1982c:270, ♀. Holotype: ♀, Venezuela: Merida: Apartaderos (AMNH). – Leclercq, 1982c:254 (in key to subgenus *Parechuca*), 2000a:19 (in key to *Podagritus* of South and Central America), 42 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

### 65. *mirandae* Leclercq

*Podagritus mirandae* Leclercq, 1981d:85, ♀, ♂. Holotype: ♀, Argentina: La Rioja: Angulos area (IML). – Leclercq, 1981d:72 and 75 (in key to subgenus *Podagritus* s.s. of South America), 2000a:11, 15 (in key to *Podagritus* of South and Central America), 42 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

### 66. *mitrensis* Leclercq

*Podagritus mitrensis* Leclercq, 1998c:297, ♀, ♂. Holotype: ♀, Australia: Victoria: 12.5 km north-northeast of Mitre at 36°37'S 141°49'E (ANIC). – Leclercq, 1998c:293 (in key to Australian *Podagritus*).

**67. mollarus Leclercq**

*Podagritus mollarus* Leclercq, 1981d:85, ♂. Holotype: ♂, Argentina: Tucumán: San Pedro Colalao (IML). – Leclercq, 1981d:75 (in key to subgenus *Podagritus* s.s. of South America), 2000a:11, 15 (in key to *Podagritus* of South and Central America), 42 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

**68. mullewanus Leclercq**

*Podagritus mullewanus* Leclercq, 1970b:273, ♀. Holotype: ♀, Australia: Western Australia: 10 mi. west of Mullewa (ANIC). – R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:271 (in catalog of Australian Sphecidae); Leclercq, 1998c:290 (in key to Australian *Podagritus*), 298 (bibliographic reference to original description).

**69. musellus Leclercq**

*Podagritus musellus* Leclercq, 1998c:298, ♀. Holotype: ♀, Australia: Southern Australia: 25 km south-southeast of Pinnaroo at 35°28'S 140°47'E (ANIC). – Leclercq, 1998c:290 (in key to Australian *Podagritus*).

**70. myrmosus Leclercq**

*Podagritus myrmosus* Leclercq, 1998c:299, ♀. Holotype: ♀, Australia: Western Australia: 13 mi west of Coolgardie (MCZ). – Leclercq, 1998c:290 (in key to Australian *Podagritus*).

**71. napieri Leclercq**

*Podagritus napieri* Leclercq, 1998c:299, ♀, ♂. Holotype: ♀, Australia: Western Australia: Napier Creek (misspelled Nappiez Creek) 21 km north-northeast of Albany at 35°40'S 117°58'E (ANIC). – Leclercq, 1998c:290 (in key to Australian *Podagritus*).

**72. nebulosus Leclercq**

*Podagritus nebulosus* Leclercq, 1982c:270, ♀. Holotype: ♀, Ecuador: Tungurahua: Baños (BMNH). – Leclercq, 1982c:254 (in key to subgenus *Parechuca*), 2000a:18 (in key to *Podagritus* of South and Central America), 42 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**73. neuqueni Leclercq**

*Podagritus neuqueni* Leclercq, 1957f:19, ♀, ♂. Holotype: ♀, Argentina: Parque Nacional Lanin (IML). – Leclercq, 1957f:11 (in key to subgenus *Echuca*); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1982c:253 (in key to subgenus *Parechuca*), 271 (Argentina, Chile; additional description), 1994b:241 (in key to *Podagritus* of Chile), 249 (bibliographic references, locality records), 2000a:20, 28 (in key to *Podagritus* of South and Central America), 42 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**74. nigriventris (Brèthes)**

*Crabro nigriventris* Brèthes, 1913:138, ♂. Holotype or syntypes: ♂, Argentina: Buenos Aires (MACN). – Genise, 1990:26 (type material in MACN). – **As *Podagritus nigriventris***: Leclercq, 1951f:11 (new combination, redescription), 1954c:192 (listed), 1957h:7 (in key to subgenus *Podagritus* s.s.), 13 (in revision of subgenus *Podagritus* s.s., Argentina, Paraguay, ♀ = *Podagritus apostolus*), 1970b:279 (Argentina: Entre Ríos Province); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:70, 71, 74 (in key to subgenus *Podagritus* s.s. of South America), 85 (description of holotype; validity of related species; Argentina, Bolivia, Brazil), 2000a:9, 14 (in key to *Podagritus* of South and Central America), 42 (bibliographic references, additional locality records, variation); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

*Podagritus concordius* Leclercq, 1970b:278, ♀, ♂. Holotype: ♀, Argentina: Entre Ríos: Concordia (FSAG). Paratypes: Brazil. Synonymized with *Podagritus nigriventris* by Leclercq, 2000a:42. – R. Bohart and Menke, 1976:393 (in

checklist of world Sphecidae); Leclercq, 1981d:71 and 74 (in key to subgenus *Podagritys* s.s. of South America), 81 (Argentina, Brazil, Uruguay), 2007g:194 (holotype in FSAG).

### 75. *noguesus* Leclercq

*Podagritys noguesus* Leclercq, 1982c:271, ♀, ♂. Holotype: ♂, Argentina: Villa Nogués (AEI). – Leclercq, 1982c:261 (in key to subgenus *Parechuca*), 2000a:21, 24 (in key to *Podagritys* of South and Central America), 43 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

### 76. *noongaris* Leclercq

*Podagritys noongaris* Leclercq, 1998c:300, ♀. Holotype: ♀, Australia: Western Australia: 2 km west-southwest of Noongar at 31°21'S 118°57'E (ANIC). – Leclercq, 1998c:292 (in key to Australian *Podagritys*) Pulawski, 2025b:230 (redescription).

### 77. *paractus* Leclercq

*Podagritys paractus* Leclercq, 1982c:272, ♂. Holotype: ♂, Bolivia: Cochabamba: Chapare, Paracti (AMNH). – Leclercq, 1982c:254 (in key to subgenus *Parechuca*), 2000a:24 (in key to *Podagritys* of South and Central America), 43 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Dollfuss, 2004b:782 (Bolivia: Cochabamba: Carrasco Biberia).

### 78. *parrotti* (Leclercq)

*Rhopalum parrotti* Leclercq, 1955k:13, ♀, ♂. Holotype: ♀, New Zealand: Wellington: Wilton Bush (originally A.W. Parrott coll., now New Zealand Natl. Mus.). – Leclercq, 1955k:8 (in key to *Rhopalum* of New Zealand); Palma, Lovis, and Tither, 1989:49 (holotype is in Natl. Mus. of New Zealand). – **As *Podagritys parrotti***: R. Bohart and Menke, 1976:393 (new combination, in checklist of world Sphecidae); Callan, 1979:39 (New Zealand); Valentine and Walker, 1991:40 (in catalog of Hymenoptera of New Zealand); Harris, 1994:48 (in Fauna of New Zealand, nest and prey), 1998:591 (preying exclusively on adult chrysomelid *Adoxia vulgaris* Broun), 2000a:74 (nest structure, prey: *Adoxia pubicollis* Broun, a galerucine chrysomelid).

### 79. *pasconus* Leclercq

*Podagritys pasconus* Leclercq, 1982c:272, ♀. Holotype: ♀, Peru: Pasco: Oxapampa (BMNH). – Leclercq, 1982c:263 (in key to subgenus *Parechuca*), 2000a:21, 27 (in key to *Podagritys* of South and Central America), 43 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

### 80. *paynesis* Leclercq

*Podagritys paynesis* Leclercq, 1998c:301, ♀, ♂. Holotype: ♀, Australia: Western Australia: 5 km west-southwest of Paynes Find at 29°18'S 117°39'E (ANIC). – Leclercq, 1998c:289 (in key to Australian *Podagritys*).

### 81. *pecunius* Leclercq

*Podagritys pecunius* Leclercq, 1982c:272, ♀, ♂. Holotype: ♂, Argentina: Tucuman: Villa Nogés (AEI). Paratypes: Bolivia, Chile. – Leclercq, 1982c:264, 265 (in key to subgenus *Parechuca*), 1994b:241 and 242 (in key to *Podagritys* of Chile), 249 (bibliographic references, Chilean locality), 2000a:22, 29 (in key to *Podagritys* of South and Central America), 43 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

As *Podagritys geraesae*: Leclercq, 1957f:12, 14 (in key to subgenus *Echuca*), corrected to *Podagritys pecunius* by Leclercq, 1982c:272.

### 82. *peratus* Leclercq

*Podagritys peratus* Leclercq, 1955f:326, ♂. Holotype: ♂, Australia: Western Australia: Yallingup (BMNH). – Leclercq, 1955f:312 (in key to Australian *Podagritys*), 1970b:273 (additional localities, comparison with *Podagritys*

*krombeini*); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:271 (in catalog of Australian Sphecidae); Leclercq, 1998c:289 (in key to Australian *Podagritus*), 301 (locality records, variation).

### 83. *picchus* Leclercq

*Podagritus picchus* Leclercq, 1982c:273, ♀, ♂. Holotype: ♀, Peru: Machu Picchu (MCZ). – Leclercq, 1982c:261, 264, 265 (in key to subgenus *Parechuca*), 2000a:22, 29 (in key to *Podagritus* of South and Central America), 44 (bibliographic references, additional locality records, prey: an empidid); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

### 84. *pilosus* Pulawski

*Podagritus pilosus* Pulawski, 2025b:226, ♀. Holotype: ♀, Australia: Western Australia: Karroun Hill Nature Reserve at 30.076°S 117.83°E (WAM).

### 85. *piscator* Leclercq

*Podagritus piscator* Leclercq, 1998c:301, ♂. Holotype: ♂, Australia: Western Australia: 9 km southeast of Carrabin at 31°22'S 118°47'E (ANIC). – Leclercq, 1998c:289 (in key to Australian *Podagritus*).

### 86. *pius* (Strand)

*Crabro pius* Strand, 1910a:156, ♀. Holotype: ♀, Paraguay: Asunción (ZMHU). – *As Rhopalum pius*: Leclercq, 1954c:186 (new combination, in catalog of world Crabronini). – *As Podagritus pius*: Leclercq in R. Bohart and Menke, 1976:393 (new combination, in checklist of world Sphecidae); Leclercq, 1982c:264 (in key to subgenus *Parechuca*), 273 (Argentina, Brazil, Colombia, Ecuador, Peru), 2000a:21, 22 (in key to *Podagritus* of South and Central America), 44 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

*Podagritus geraesae* Leclercq, 1951f:10, ♀. Holotype: ♀, Brazil: Minas Gerais: no specific locality (MNHN). Synonymized with *Podagritus pius* by Leclercq in R. Bohart and Menke, 1976:393. – Leclercq, 1954c:192 (in catalog of world Crabronini), 1957f:12 (in key to subgenus *Echuca*), nec 14 (in key = *Podagritus pecunius*), 18 (Argentina, Brazil, Peru; additional description of female).

### 87. *pixitus* Leclercq

*Podagritus pixitus* Leclercq, 1998c:302, ♀. Holotype: ♀, Australia: Northern Territory: James Ranges at 24°15'S 113°26'E (ANIC). – Leclercq, 1998c:290 (in key to Australian *Podagritus*).

### 88. *pizarrus* Leclercq

*Podagritus pizarrus* Leclercq, 1957h:14, ♂. Holotype: ♂, Argentina: Buenos Aires: San Clemente del Tuyu (IML). – Leclercq, 1957h:8 (in key to subgenus *Podagritus* s.s.); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:70, 73 (in key to subgenus *Podagritus* s.s. of South America), 86 (Brazil; color variation), 2000a:9, 13 (in key to *Podagritus* of South and Central America), 45 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

### 89. *polybia* Schrottky

*Podagritus polybia* Schrottky, 1909b:251, ♀. Holotype or syntypes: ♀, Paraguay: Puerto Bertoni (MACN). – Schrottky, 1913a:233 (Paraguay); Leclercq, 1954c:193 (new combination, in catalog of world Crabronini); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 2000a:45 (original description copied); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

### 90. *porteri* Leclercq

*Podagritus porteri* Leclercq, 1982c:273, ♀, ♂. Holotype: ♀, Argentina: Tucuman: Tafi del Valle (MCZ). – Leclercq, 1982c:257 (in key to subgenus *Parechuca*), 2000a:17, 24 (in key to *Podagritus* of South and Central America), 45 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**91. *propodealis* Pulawski**

*Podagritus propodealis* Pulawski, 2025b:228, ♀. Holotype: ♀, Australia: Western Australia: Yanchep National Park at 31.5063°S 115.6811°E (WAM).

**92. *quiaca* Leclercq**

*Podagritus quiaca* Leclercq, 1982c:274, ♀, ♂. Holotype: ♀, Argentina: Jujuy: La Quiaca (originally M. Fritz coll., now MACN). – Leclercq, 1982c:262, 263 (in key to subgenus *Parechuca*); Fritz, 1993:17 (holotype in M. Fritz coll.); Leclercq, 2000a:22, 24 (in key to *Podagritus* of South and Central America), 45 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**93. *rhopaloides* Leclercq**

*Podagritus rhopaloides* Leclercq, 1951f:2, ♂. Holotype: ♂, Ecuador: Quito (MNHN). – Leclercq, 1954c:192 (in catalog of world Crabronini), 1957h:7 (in key to subgenus *Podagritus* s.s.), 8 (transferred to subgenus *Podagritus* s.s.); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1982c:253, 256 (in key to subgenus *Parechuca*), 274 (Ecuador), 2000a:19, 25 (in key to *Podagritus* of South and Central America), 45 (additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

**94. *rieki* Leclercq**

*Podagritus rieki* Leclercq, 1957e:1, ♀, ♂. Holotype: ♀, Australia: Australian Capital Territory: Blundells (ANIC). – Leclercq, 1970b:275 (Victoria); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:271 (in catalog of Australian Sphecidae); Leclercq, 1998c:289 (in key to Australian *Podagritus*), 302 (locality records).

**95. *risettus* Leclercq**

*Podagritus risettus* Leclercq, 1982c:275, ♂. Holotype: ♂, Colombia: Bogota (ZMHU). – Leclercq, 1982c:255 (in key to subgenus *Parechuca*), 2000a:25 (in key to *Podagritus* of South and Central America), 46 (bibliographic references, additional locality records, first record from Chile); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**96. *riveti* (Strand)**

*Crabro riveti* Strand, 1911a:152, ♂ (as *Riveti*, incorrect original capitalization). Holotype: ♂, Ecuador: Casitagua (MNHN). – Strand, 1913:26 (Ecuador). – **As *Podagritus riveti***: Leclercq, 1951f:5 (new combination, additional localities), Leclercq, 1954c:191 (in catalog of world Crabronini), 1957h:7 (in key to subgenus *Podagritus* s.s.), 9 (in revision of subgenus *Podagritus* s.s.); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1982c:253, 254 (in key to subgenus *Parechuca*), 275 (Colombia, Ecuador), 2000a:19, 25 (in key to *Podagritus* of South and Central America), 46 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**97. *rozeni* Leclercq**

*Podagritus rozeni* Leclercq, 1982c:275, ♀, ♂. Holotype: ♂, Ecuador: Azuay: Santa Isabel 65 km southwest of Cuneca (AMNH). – Leclercq, 1982c:255 (in key to subgenus *Parechuca*), 2000a:17, 24 (in key to *Podagritus* of South and Central America), 46 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**98. *rufotaeniatus* (Kohl)**

*Crabro rufotaeniatus* Kohl, 1905b:353, ♂. Lectotype: ♂, Chile: no specific locality (NHMW), designated by Leclercq, 1982c:275. – Janvier, 1928:73 (nesting habits), 1929:418 (sleeping habits); Piek and Spanjer, 1986:183 (in list of Sphecidae with known prey). – **As *Rhopalum rufotaeniatum***: Leclercq, 1954c:186 (new combination, in catalog of world Crabronini), 298 (nest and prey records). – **As *Podagritus rufotaeniatus***: Leclercq in R. Bohart and Menke, 1976:393 (new combination, in checklist of world Sphecidae); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae); Leclercq, 1982c:262 (in key to subgenus *Parechuca*), 275 (Chile); Dollfuss, 1989:11 (holotype in NHMW); Steiner, 1986:107 (references to publications on nesting habits); Leclercq, 1994b:241 (in key to *Podagritus* of Chile), 249

(bibliographic references, faunal records, variation), 2000a:20, 26 (in key to *Podagritus* of South and Central America), 46 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

*Podagritus getricus* Leclercq, 1957f:16, ♀, ♂. Holotype: ♀, Chile: Bío Bío: Los Angeles (IML). Synonymized with *Podagritus rufotaeniatus* by Leclercq in R. Bohart and Menke, 1976:393. – Leclercq, 1957f:12 (in key to subgenus *Echuca*).

#### 99. *sellosus* Leclercq

*Podagritus sellosus* Leclercq, 1982c:276, ♂. Holotype: ♂, Brazil: "Sello" (ZMHU). – Leclercq, 1982c:262 (in key to subgenus *Parechuca*), 2000a:21, 27 (in key to *Podagritus* of South and Central America), 47 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

#### 100. *sericinus* Leclercq

*Podagritus sericinus* Leclercq, 1981d:86, ♀, ♂. Holotype: ♂, Argentina: Chubut: El Trebol 30 km of Comodoro Rivadavia (IML). – Leclercq, 2000a:8, 10, 14 (in key to *Podagritus* of South and Central America), 47 (bibliographic references), 1981d:69, 71, 74 (in key to subgenus *Podagritus* s.s. of South America); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

#### 101. *sombratus* Leclercq

*Podagritus nigriventris* var. *sombratus* Leclercq, 1951f:12, ♂. Holotype: ♂, Brazil, Rio de Janeiro: Itatiaya (MNHN). – Leclercq, 1954c:192 (in catalog of world Crabronini), 1957h:7 (in key to subgenus *Podagritus* s.s.), 14 (is conspecific with *Podagritus nigriventris*). – **As *Podagritus sombratus***: Leclercq, 2000a:8, 14 (in key to *Podagritus* of South and Central America), 47 (bibliographic references, additional locality records), 1981d:70, 74 (in key to subgenus *Podagritus* s.s. of South America), 86 (as full species; redescription); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

#### 102. *sorbicus* Leclercq

*Podagritus sorbicus* Leclercq, 1957f:17, ♀, ♂. Holotype: ♀, Chile: Cordillera de Nahuelbuta: Pillim Pilli (Inst. Roy. Sci. Nat. Belgique, corrected to Gembloux by Leclercq, 1982c:276). Paratypes: Argentina. – Leclercq, 1957f:12 (in key to subgenus *Echuca*), Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae); Leclercq, 1982c:260 (in key to subgenus *Parechuca*), 276 (additional description), 1994b:241, 242 (in key to *Podagritus* of Chile), 250 (bibliographic references, faunal records), 2000a:20, 28 (in key to *Podagritus* of South and Central America), 47 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Leclercq, 2007g:194 (holotype in FSAG).

#### 103. *stuartae* Pulawski

*Podagritus stuartae* Pulawski, 2023:126, ♀, ♂. Holotype: ♀, Australia: Western Australia: Creyk Park in Kelmscott (a south-eastern suburb of Perth) at 32.132°S 116.015°E (WAM). – Houston and Stuart, 2023:135 (nesting habits).

#### 104. *subandinus* Leclercq

*Podagritus subandinus* Leclercq, 1982c:277, ♂. Holotype: ♂, Peru: Atocongo (IML). – Leclercq, 1982c:259 (in key to subgenus *Parechuca*), 2000a:25 (in key to *Podagritus* of South and Central America), 47 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Rasmussen and Asenjo, 2009:9 (in checklist of Crabronidae of Peru).

#### 105. *taficus* Leclercq

*Podagritus taticus* Leclercq, 1982c:277, ♀, ♂. Holotype: ♂, Argentina: Tucuman: Tafi del Valle (AEI). – Leclercq, 1982c:263, 264 (in key to subgenus *Parechuca*), 2000a:22, 28 (in key to *Podagritus* of South and Central America), 47 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**106. *teresoides* Leclercq**

*Podagritus teresoides* Leclercq, 1982c:277, ♀. Holotype: ♀, Uruguay: Montevideo: Colón (BMNH). – Leclercq, 1982c:261 (in key to subgenus *Parechuca*), 2000a:21 (in key to *Podagritus* of South and Central America), 47 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**107. *teresus* Leclercq**

*Podagritus teresus* Leclercq, 1982c:277, ♀, ♂. Holotype: ♂, Brazil: Rio Grande do Sul: Teresopolis (AEI). – Leclercq, 1982c:260 (in key to subgenus *Parechuca*), 2000a:21, 26 (in key to *Podagritus* of South and Central America), 47 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**108. *terpenus* Leclercq**

*Podagritus terpenus* Leclercq, 1957h:16, ♀, ♂. Holotype: ♂, Argentina: Jujuy: Tumbaya (IML). – Leclercq, 1957h:3 (in key to subgenus *Podagritus* s.s.); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:72, 75 (in key to subgenus *Podagritus* s.s. of South America), 87 (additional description), 2000a:11, 15 (in key to *Podagritus* of South and Central America), 48 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

**109. *toxopeusi* Tsuneki**

*Podagritus toxopeusi* Tsuneki, 1983h:3, ♂ (subgenus *Ebibus*). Holotype: ♂, New Guinea: "Moss Forest Camp, 2700 m" (RMNH).

**110. *tricolor* (F. Smith)**

*Crabro tricolor* F. Smith, 1856:394, ♂. Holotype or syntypes: ♂, Australia: Tasmania: no specific locality (originally BMNH, now lost: R. Turner, 1915a:92). Neotype: ♂, Australia: Tasmania: Eaglehawk Neck (BMNH), designated by Leclercq, 1955f:321. – Dalla Torre, 1897:632 (in catalog of world Sphecidae); R. Turner, 1908:524 (description of ♀; New South Wales, Victoria, Tasmania). – **As *Rhopalum tricolor***: R. Turner, 1915a:86 (in key to Australian *Rhopalum*), 92 (Tasmania, southeastern Australia; nesting in sandy banks), 1915b:552 (in key to *Rhopalum* of Tasmania), 553 (Tasmania). – **As *Podagritus tricolor***: Leclercq, 1950b:7 (is a member of *Podagritus*), 1954c:191 (in catalog of world Crabronini), 1955f:311 (in key to Australian *Podagritus*), 321 (redescription), 1957e:4 (additional Australian localities), 1970b:275 (Tasmania); Riek, 1970:942 (in Insect Fauna of Australia); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:271 (in catalog of Australian Sphecidae); K. Walker, Naumann, Austin, Taylor, and Cardale, 1992:48 (in catalog of insects of Tasmania); Leclercq, 1998c:292 (in key to Australian *Podagritus*), 302 (locality records).

*Crabro militaris* R. Turner, 1908:523, ♂. Syntypes: Australia: Victoria and Tasmania: no specific localities (BMNH). Synonymized with *Podagritus tricolor* by R. Turner, 1915a:553, synonymy confirmed by Leclercq, 1950b:7 and 1955f:321. – **As *Podagritus militaris***: Pate, 1944:354 (new combination).

**111. *trifidus* Leclercq**

*Podagritus trifidus* Leclercq, 1998c:303, ♀, ♂. Holotype: ♀, Australia: Western Australia: 37 km southwest of Eucla at 31°45'S 128°33'E (ANIC). – Leclercq, 1998c:289, 290 (in key to Australian *Podagritus*).

**112. *tucumanus* Leclercq**

*Podagritus tucumanus* Leclercq, 1982c:278, ♀, ♂. Holotype: ♀, Argentina: Tucuman: Tafi del Valle (AEI). – Leclercq, 1982c:259 (in key to subgenus *Parechuca*), 2000a:18, 25 (in key to *Podagritus* of South and Central America), 48 (bibliographic references); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

**113. *valdiviae* Leclercq**

As *Physoscelus longinodus* Spinola, 1851:359, ♂ only, corrected to *Podagritus valdiviae* by Leclercq, 1994b:247.

*Podagritus valdiviae* Leclercq, 1951f:7, ♀, ♂. Holotype: ♀, Chile: Valdivia (MNHN). – Leclercq, 1954c:192 (in catalog of world Crabronini), 1957f:12 (in key to subgenus *Echuca*), 19 (diagnostic characters); R. Bohart and Menke,

1976:393 (in checklist of world Sphecidae); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae); Leclercq, 1982c:258 (in key to subgenus *Parechuca*), 278 (Argentina, Chile), 1994b:241, 242 (in key to *Podagritus* of Chile), 251 (bibliographic references, faunal records), 2000a:19, 28 (in key to *Podagritus* of South and Central America), 48 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Chiappa, 2012:8 (Chile: Región de Valparaíso: no specific locality).

*Podagritus almargus* Leclercq, 1957f:15, ♀. Holotype: ♀, Argentina: Río Negro: Puerto Radal, Isla Victoria (IML). Synonymized with *Podagritus valdiviae* by Leclercq in R. Bohart and Menke, 1976:393. – Leclercq, 1957f:12 (in key to subgenus *Echuca*).

#### 114. *valenciai* Fritz

*Podagritus valenciai* Fritz, 1971a:113, ♀, ♂. Holotype: ♂, Chile: Malleco: Lago Galletue (originally M. Fritz coll., now MACN). – R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:67 (in key to subgenus *Chilichuca*), 76 (transferred to subgenus *Chilichuca*; Argentina, Chile); Sielfeld, 1980b:74 (in checklist of Chilean Sphecidae); Fritz, 1993:17 (holotype in M. Fritz coll.); Leclercq, 1994b:242 (in key to *Podagritus* of Chile), 251 (bibliographic references, faunal records), 2000a:16 (in key to *Podagritus* of South and Central America), 48 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae), 2005a:10 (correction to his 2002 catalog).

#### 115. *venturii* (Schrottky)

*Crabro venturii* Schrottky, 1902b:114, ♂. Holotype: ♂, Argentina: Buenos Aires (depository?). – Schrottky, 1903b:177 (in checklist of Hymenoptera of Argentina, Paraguay, and Uruguay). – As *Podagritus venturii*: Schrottky, 1909:251 (new combination, coloration as in *Podagritus polybia*), 1913a:233 (Argentina: Buenos Aires); Leclercq, 1951f:14 (redescription), 1954c:192 (in catalog of world Crabronini), 1957h:5 (in key to subgenus *Podagritus* s.s.), 12 (in revision of subgenus *Podagritus* s.s., Argentina, Paraguay, comparison with *Podagritus joergenseni*), 1970b:279 (Argentina: Buenos Aires); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:71 and 75 (in key to subgenus *Podagritus* s.s. of South America), 87 (Argentina, Brazil, Uruguay), 2000a:10, 16 (in key to *Podagritus* of South and Central America), 48 (bibliographic references, additional locality records); Amarante, 2002:31 (in catalog of Neotropical Crabronidae).

#### 116. *victoriae* Leclercq

*Podagritus victoriae* Leclercq, 1998c:304, ♀, ♂. Holotype: ♀, Australia: Victoria: Healesville (National Mus. of Victoria). – Leclercq, 1998c:292 (in key to Australian *Podagritus*).

#### 117. *virtanus* Leclercq

*Podagritus virtanus* Leclercq, 1981d:78, ♀, ♂. Holotype: ♀, Chile: Malleco Province: Galletue in Cordillera Lonquimay (MCZ). Paratypes: Argentina. – Leclercq, 1981d:67 (in key to subgenus *Chilichuca*), 1994b:242 (in key to *Podagritus* of Chile), 252 (bibliographic references, faunal records), 2000a:16 (in key to *Podagritus* of South and Central America), 49 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

#### 118. *wilkersoni* Leclercq

*Podagritus wilkersoni* Leclercq, 2000a:49, ♀. Holotype: ♀, Colombia: Departamento Valle: Lago Calima (FSCA). – Leclercq, 2000a:21 (in key to *Podagritus* of South and Central America); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

#### 119. *willinki* Leclercq

*Podagritus willinki* Leclercq, 1957h:15, ♀, ♂. Holotype: ♂, Argentina: Buenos Aires: San Clemente del Tuyu (IML). – Leclercq, 1957h:3 (in key to subgenus *Podagritus* s.s.); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Leclercq, 1981d:72, 75 (in key to subgenus *Podagritus* s.s. of South America), 88 (additional locality

records), 2000a:12, 15 (in key to *Podagritys* of South and Central America), 49 (bibliographic references, additional locality records); Amarante, 2002:32 (in catalog of Neotropical Crabronidae).

#### 120. *yarrowi* Leclercq

*Podagritys yarrowi* Leclercq, 1955f:328, ♀. Holotype: ♀, Australia: Western Australia: Dongarra (BMNH). – Leclercq, 1955f:313 (in key to Australian *Podagritys*); R. Bohart and Menke, 1976:393 (in checklist of world Sphecidae); Cardale, 1985:272 (in catalog of Australian Sphecidae); Leclercq, 1998c:292 (in key to Australian *Podagritys*), 305 (locality records).

#### 121. *ypirangae* Leclercq

*Podagritys ypirangae* Leclercq, 2000a:50, ♀, ♂. Holotype: ♀, Brazil: São Paulo: Ypiranga (Museo de Zoologia, Universidade de São Paulo). – Leclercq, 2000a:9, 14 (in key to *Podagritys* of South and Central America); Amarante, 2002:32 (in catalog of Neotropical Crabronidae); Menezes de Jesus, Onody, dos Santos Ramos, de Oliveira Andrade, Doria dos Santos, and Ferreira Brandão, 2019:2 (holotype in Mus. Zool. Univ. São Paulo).

As *Podagritys apostolus*: Leclercq, 1981d:71 (in key to subgenus *Podagritys* s.s. of South America), 79 (description, ♀ only), corrected to *Podagritys ypirangae* by Leclercq, 2000a:50.

#### sp.

Aguilar, 1966:95 (Peru: hills in vicinity of Lima), 1976:59 (Peru: coastal hills); Sk. Yamane, 1983:87 (Indonesia: Krakatau Islands); Silvestre, Demétrio, Trad, de Oliveira Lima, Auko, and de Souza, 2014:68 (Brazil: Mato Grosso do Sul: dry forests in Bodoquena Mountain Range and Brazilian Chaco); Trad and Silvestre, 2017:5 (Brazil: Mato Grosso do Sul).

### SPECIES DESCRIBED IN *PODAGRITUS* AND TRANSFERRED TO OTHER GENERA

*bamendae* Leclercq, 1961 = *Rhopalum*

*cercericus* Leclercq, 1957 = *Echucoides*

*piratus* Leclercq, 1957 = *Echucoides*

### SPECIES TRANSFERRED TO AND SUBSEQUENTLY REMOVED FROM *PODAGRITUS*

*Rhopalum cornigerum* Tsuneki, 1977.

*Rhopalum cygnorum* Turner, 1915

*Crabro oceanicus* W. Schulz, 1906